



NEGATIVE PRESSURE THERAPY FOR WOUND HEALING

Dr. T. Srinivas MS	Associate Professor, Department of General Surgery, Government Medical College, Mahabubnagar.
Dr. B. Suneel Kumar MS	Assistant Professor, Department of General Surgery, Government Medical College, Mahabubnagar.
Dr. A. Nagaraju MS	Assistant Professor, Department of General Surgery, Government Medical College, Mahabubnagar.
Dr. M. Aparna* MS	Assistant Professor, Department of General Surgery, Government Medical College, Mahabubnagar. *Corresponding Author

ABSTRACT **Objective:** Wounds are a major source of morbidity, lead to considerable disability, and are associated with increased mortality; therefore, they have a significant impact on public health and the expenditure of healthcare resources. Vacuum-assisted closure (VAC) uses negative pressure to assist wound healing. Negative pressure drains fluid from the wound, thus removing the substrate for growth of microorganisms. Negative pressure may also accelerate granulation tissue formation and promote angiogenesis. The mechanical stimulation of cells by tensile forces may also play a role by increasing cellular proliferation and protein synthesis. Negative pressure wound therapy (NPWT) involves the use of a negative pressure therapy or suction device to aspirate and remove fluids, debris, and infectious materials from the wound bed to promote the formation of granulation tissue. **Methodology:** A total of 50 cases clinically presenting as ulcer between January 2019 and December 2020 in Government Medical College, Mahabubnagar were taken for study. Each case was examined clinically in a systematic manner for study of all patients presenting with ulcer. VAC dressing was done and outcome was measured by recording wound scores on days 3, 7 and 10. **Results:** In our clinical study of 50 cases managed by VAC dressing, 50% of the cases were of traumatic, 36% diabetic and 14% cases of vascular etiology. Wound healing was better in the non-diabetic group compared to diabetic and in non-smokers compared to smokers. Commonest organisms isolated were Staphylococcus, Pseudomonas and Proteus. Interpretation and conclusion: In our study VAC therapy enhanced granulation tissue formation leading to better wound healing, and faster recovery. VAC is thus a promising new technology can be used in both acute and chronic wounds, as an adjuvant therapy to improve the results of various surgical procedures.

KEYWORDS : VAC, NEGATIVE PRESSURE WOUND THERAPY, WOUND HEALING.

INTRODUCTION

Vacuum-assisted closure (VAC) is a new technique in the challenging field of management of contaminated, acute and chronic wounds. Negative pressure wound therapy (NPWT) also called vacuum-assisted wound closure and refers to wound dressing systems that continuously or intermittently apply sub-atmospheric pressure to the surface of a wound. The application of controlled levels of negative pressure has been shown to accelerate debridement and promote healing in many different types of wounds. The optimum level of negative pressure appears to be around 125 mmHg below ambient and it is believed that negative pressure assists in removal of interstitial fluid, decreasing localized edema and increasing blood flow. This in turn decreases tissue bacterial levels. Despite the significant costs involved, the technique is said to compare favorably in financial terms with conventional treatments in the management of difficult wounds.¹ It involves application of sterile, open-pore foam dressing directly on the wound. The wound is then sealed with an occlusive drape in order to create a closed, controlled environment. A fenestrated vacuum tube is connected to a vacuum source; fluid is drawn from the wound through the foam into a reservoir for subsequent disposal. Negative pressure is applied at 50-125 mm/Hg, resulting in a decrease in the local interstitial pressure, and effluent from the wound is drawn out into the collection device. Initially, the vacuum pressure is applied continuously. As the amount of drainage decreases, the vacuum may be applied on an intermittent basis. The vacuum dressing is usually changed at approximately 48-hour interval.^{2,3} Wound progress is recorded using parameters in the wound scoring system. The objectivity of assessments used to generate the wound score make this scoring system ideal for evaluating treatment and outcome of wounds.

OBJECTIVES OF THE STUDY

1. To study the outcome of vacuum assisted closure of wounds.

INTRODUCTION

A wound is defined as a discontinuity of the skin, mucous membrane or tissue caused by physical, chemical or biological insult. Surgeon's goal in wound management is to create an environment where the healing process can proceed in an optimal fashion. Wound healing is a fundamental hemostatic process in response to injury. It involves the

activation of basic cellular process of inflammation, cell proliferation, and cell growth as well as regeneration of these processes once repair is complete. The processes of wound healing take place in an overlapping and sequential manner

PHASES OF WOUND HEALING

1. Hemostasis and inflammatory phase
2. Proliferative phase
3. Maturation and remodeling phase.

MANAGEMENT OF WOUNDS

Wound healing is a highly orchestrated process, which commences with the removal of debris and control of infection. Subsequently, the wound heals through deposition of granulation tissue, wound contraction and maturation. Factors such as pressure, trauma, venous insufficiency, diabetes mellitus, vascular disease and prolonged immobilization will influence wound healing.⁹

Standard wound management consists of initial surgical debridement, then either wet-to moist (WM) gauze dressings, which need to be changed at least twice daily, can be used to cover the wound. These dressings are relatively inexpensive, readily available and easy to apply. Vacuum-assisted closure has been suggested as an alternative that may promote faster wound healing with fewer painful dressing changes.

VACUUM-ASSISTED CLOSURE

Vacuum-assisted closure is a potential alternative for treating of skin wounds. It uses negative pressure to assist wound healing.

INDICATIONS AND CONTRAINDICATIONS FOR THE USE OF VAC

The principal indications for the use of VAC are:

- Acute and traumatic wounds
- Pressure ulcers
- Chronic open wounds (stasis ulcers and diabetic ulcers)
- Meshed grafts
- Flaps
- Lower extremity diabetic ulcers

Contraindications are:
 Fistulas to organs or body cavities
 Necrotic tissue in eschar
 Osteomyelitis (untreated)
 Malignancy in the wound

MATERIALS AND METHODS

A total of 50 cases clinically presenting as ulcer between January 2019 and December 2020 were included in the study. Clinical examination of each case was done systematically for the study.

Wound Preparation:

Any dressings from the wound was removed and discarded. A culture swab for microbiology was taken before wound irrigation with normal saline. Surgical debridement was done and adequate haemostasis achieved.

METHOD OF USE OF NPWT DRESSING

The sterile hydrocolloid sheet of approximate size of the wound is placed gently into position.



RESULTS

An interventional study involving 50 cases of wounds was done in the Department of Surgery, Government Medical College, Mahabubnagar between January 2019 and December 2020.

CHART 1: AGE DISTRIBUTION OF WOUNDS




TABLE 1: AGE WISE EVALUATION OF WOUND HEALING SCORE ON DAY 7

Age in years	Day 7			
	N	Mean	SD	Median
<40	11	5.09	0.94	5
41-50	15	4.73	1.22	5
>50	15	4.87	1.06	5
Total	41	4.88	1.08	5

Kruskalwallis test, p=0.8

GENDER DISTRIBUTION OF WOUNDS

Wounds were more common in males 38 cases (76%) than in females 12 cases (24%) Male to female ratio 3.167: 1.

DURATION OF WOUNDS


Based on the duration of wounds, cases were grouped into 3 categories: <10 days, 10-30 days and >30 days. Most cases fall in the group 10-30 days 30(60%), 19 cases(38%) in the group <10 days and 1 case (2%) in the group > 30 days.

TABLE 2: DURATION OF WOUND - EVALUATION OF WOUND HEALING SCORE ON DAY 7

Duration in Years	Day 7			
	N	Mean	SD	Median
<10 days	16	4.94	0.85	5
>10 days	25	4.84	1.21	5
Total	41	4.88	1.08	5

Mann whitney test, p=0.7

CHART 2: DISTRIBUTION OF LOCATION OF WOUNDS



Wounds were most commonly located in the foot 20(40%) followed by the leg 16(32%) and the ankle and sole 4(8%) each. One case of wound in the abdomen was included.

TABLE 3: LOCATION OF WOUND - EVALUATION OF WOUND HEALING SCORE ON DAY 7

LOCATION	Wound Score					Total
	3.00	4.00	5.00	6.00	7.00	
FOREARM	0	2	3	0	0	5
ABDOMEN	0	0	0	0	1	1
LEG	1	2	7	4	0	14
ANKLE	1	0	1	1	0	3
FOOT	2	5	2	5	1	15
SOLE	1	0	2	0	0	3
TOTAL	5	9	15	10	2	41

P value – 0.024

Significant difference in wound healing depending on the location of the wound. P value- 0.024

SMOKERS AND NON-SMOKERS


27 (54%) of cases were smokers, 23 (46%) were non- smokers.

TABLE 4: SMOKING - EVALUATION OF WOUND HEALING SCORE ON DAY 7

Smoking	Day 7			
	N	Mean (18)	SD	Median
Yes	24	4.50	0.89	5
No	17	5.41	1.12	6
Total	41	4.88	1.08	5

Mann whitney test, p=0.006

CHART 3: ETIOLOGY OF WOUNDS



Based on etiology of wounds, which were determined by history and clinical examination, wounds were divided into Traumatic, Diabetic and Vascular. A major portion 25(50%) of cases fell into traumatic group and 18(36%) into diabetic and 7 (14%) into vascular group.

CHART 4: DIABETIC AND NON-DIABETIC WOUNDS





TABLE 5: ETIOLOGY - EVALUATION OF WOUND HEALING SCORE ON DAY 7


Location	N	Day 3	SD	Median
Traumatic	20	5.25	0.78	5
Diabetic	16	4.44	1.21	4
Vascular	05	4.80	1.30	5
Total	41	4.88	1.08	5

Kruskal wallis test, p=0.08 Mann-whitney test
Traumatic vs Diabetic – 0.03

CHART 5: WOUND AREA IN SQUARE CM




Wounds of area < 20 sq cm constitute maximum number of cases - 44(88%).



Most common organism cultured from the wounds was Staphylococcus aureus 21(42%).

CHART 7 : WOUND SCORES ON DAYS 3, 7 AND 10



Following VAC therapy for 3 to 7 days most of the wounds showed progress in wound healing.

CONCLUSION

VAC therapy is a recent modality of treatment of wounds. Its introduction has changed the course of management of wounds. Based on the data from the present study and other studies available, VAC does appear to result in better healing, with few serious complications, and thus looks to be a promising alternative for the management of various wounds. The application of VAC is simple, but requires training to ensure appropriate and competent use.

In patients treated with Negative Pressure Wound Therapy (NPWT) the following was noted :

- Earlier appearance of granulation and epithelialization
- Rapid decrease in wound size
- Shorter duration of hospitalization
- Early achievement complete wound score

On correlating the wound outcome variables, we also noticed that presence of Diabetes Mellitus increased the duration of formation of granulation tissue and epithelialization. This study confirms that Negative Pressure Wound Therapy is safe, has faster response in wound healing and gives better efficacy as compared to the Conventional Wound Therapy in management of lower limb ulcers. Hence it is a better choice for management of lower limb ulcers.

SUMMARY

The negative pressure wound therapy (NPWT) is a feasible and excellent option for the management of different types of wounds. Currently NPWT has become an established method in wound management.

Application of sub-atmospheric pressure decreases the bacterial colonization over the wound and increases blood flow. Increased oxygenated blood flow to the wound healing promotes the oxidative bursts in neutrophils and there by promoting the killing of microbes and preventing infection.

Most of commercially available NPWT devices are foam-based systems. In our study, We used our own NPWT by using a Romovac suction drain which is cost effective for reproducing the same results and for the benefit of patients, as majority of our patients are from the rural area.

- The present interventional study involved 50 cases of wounds.
- Patients affected were most commonly in the 5th (36%) decade followed by 6th decade (34%).
- There was a male preponderance with male: female ratio of 3.2:1.
- Most of the patients in the study presented early, presenting between 10 and 30 days of onset.
- VAC dressing was done in wounds occurring in a variety of locations like foot 20 (40%), leg 16 (32%), forearm 5 (10%), ankle and sole 4 (8%) and abdomen 1 (2%). 27 (54%) of patients were smokers
- Trauma was the most common etiology 25 (50%), followed by diabetes 18 (36%) and vascular causes 7 (14%).
- Wound area was recorded before treatment and grouped into <20 sq cm [44cases (88%)], 21-40 sq cm [4cases (8%)] and > 40 sq cm [4cases (8%)].
- Staphylococcus aureus was the most common organism cultured 21 (42%) followed by Pseudomonas aeruginosa 7 (14%). No growth was observed in 13 cases.
- Wound scores were recorded on days 3, 7 and 10 of VAC therapy. The scoring system used was based on area of granulation tissue, its color and consistency.
- Out of 18 diabetic wounds, wound scores of 5 and above were observed in 9 cases following VAC therapy. VAC dressing of longer duration are required for diabetic wounds for good outcome.
- There was significantly better outcome in non smokers compared to smokers with 75% of non smokers and 58% of smokers showing wound scores >5 on day 7 of VAC therapy.

Thus outcome of VAC therapy depends on various factors like age, aetiology of wounds and existence of comorbid conditions like diabetes and factors like smoking. The candidates for VAC therapy should be chosen after considering these factors with care. From our study, it can be concluded that VAC is a promising new technology in the field of wound healing with multiple applications in a variety of wounds.

REFERENCES

- Norman Williams, Christopher Bulstrode, Ronan O'Connell. Bailey and Love's Short Practice of Surgery. Hodder Arnold 27th Edition – 120. 1a. Stephanie R. Goldberg, Robert F. Diegelmann Wound Healing Primer Surg Clin N Am 90(2010)1133-1146.
- Courtney.M.Townsend.JR.MD,R.DanielBeauchamp.MD,B.Markevers MD. Kenneth L.Mattox,MD;Sabiston Textbook of Surgery; The Biological Basis of Modern Surgical Practice;19th edition; Vol (1) 151-177
- Leong M, Phillips LG,Wound Healing. Townsend CM, Beauchamp RD, Evers BM, Mattox KL. Sabiston Textbook of Surgery.19th ed. Philadelphia;Elsevier;2012:151-77.
- Barbul A,Efron DT. Wound healing.Brunnicardi FC.Schwartz Principles of surgery.9th ed.New York.McGraw Hill;2010:209-33.
- Galiano RD, Mustoe TA. Wound Healing.. Mulholland, Michael W, Lillmoie, Keith D, Doherty, Gerard M, Maier, Ronald V, Upchurch, Gilbert R. In Greenfield's Surgery: Scientific principles and practice.4th ed .Philadelphia; Lippincott Williams & Wilkins;2006
- Armstrong DG, Lavery LA, Frykberg RG, et al. VAC therapy appears to heal complex DFU. Abstract presented at the 2nd World Union of Wound Healing Societies' Meeting, July 8-13, 2004; Paris, France
- Joseph E, Hamori CA, Bergman S, Roaf E, Swann NF, Anastasi GW. A prospective randomized trial of vacuum-assisted closure versus standard therapy of chronic. Wounds. 2000;12(3):60-67
- Fabian TS, Kaufman HJ, Leit ED, et al. The evaluation of sub-atmospheric pressure and hyperbaric oxygen in ischemic full-thickness wound healing. Am Surg 2000;66:1136-43.
- Joseph E, Hamori CA, Bergman S, et al. A prospective randomised trial of vacuum-assisted closure versus standard therapy of chronic nonhealing wounds. Wounds 2000;3:60-7.
- Morykwas MJ, David LR, Schneider AM, et al. Use of subatmospheric pressure to prevent progression of partial-thickness burns in a swine model. J Burn Care Rehabil 1999;20:15-21.