



## STUDY OF LABORATORY PROFILE IN CHILDREN WITH ACUTE GASTROENTERITIS WITH REFERENCE TO ACID-BASE AND ELECTROLYTE IMBALANCE

<b>Jayapriya L*</b>	Junior resident, Department of Paediatrics, Dr.Vaishampayan Memorial Government Medical College, Solapur, Maharashtra, India. *Corresponding Author
<b>Ezhil Devi P</b>	Junior resident, Department of Paediatrics, Dr.Vaishampayan Memorial Government Medical College, Solapur, Maharashtra, India.
<b>Shakira Savaskar</b>	Professor & Head, Department of Paediatrics, Dr.Vaishampayan Memorial Government Medical College, Solapur, Maharashtra, India.
<b>Parmeshwar Patil</b>	Senior resident, Department of Paediatrics, Dr.Vaishampayan Memorial Government Medical College, Solapur, Maharashtra, India.

**ABSTRACT** **Background:** Diarrhoea is the third leading cause of childhood mortality in India, and responsible for 13% of all deaths/year in children under 5 year of age. Electrolyte disturbances may remain unrecognized, results in increased morbidity and mortality. The present study was conducted to assess the spectrum of electrolyte and acid base disturbances in children with acute gastroenteritis with relationship to mortality. **Material & Methods:** In this prospective clinical study total patients selected were 210, admitted in paediatric ward and paediatric intensive care unit of tertiary care hospital with diarrhoea over a period of 2 years. All patients enrolled in the study were classified according to WHO guidelines, monitored clinically and investigated for electrolytes and arterial blood gas analysis. The data collected was entered in MS excel, compiled, tabulated, analyzed and subjected to statistical tests done using SPSS. **Results:** In the present study total acute gastroenteritis patients were 210 in which 57.14% were males and 42.86% were females. Out of 210 patients 85.71% patients had some dehydration, 14.29% patients had severe dehydration. Incidence of isonatremia was 74.29%, hyponatremia was 20.47, hypernatremia was 5.24%, hypokalemia was noted in 4.76% of patients. Out of 210 patients metabolic acidosis was observed in 12.86%. Time taken for correction of electrolyte imbalance after standard treatment was 81.25% of patients in < 24 hour, 10.94% of patients in 24-48 hours, 6.25% patients in >48 hours, not corrected in 1.56% of patients. **Conclusion:** Our study concluded that most common electrolyte imbalance in acute gastroenteritis was hyponatremia followed by hypernatremia and hypokalemia. Metabolic acidosis was more common in severe dehydration compared to some dehydration. Mortality in acute gastroenteritis more commonly associated with hypernatremia. Our study shows monitoring of electrolyte and acid base levels to decrease morbidity, mortality in acute gastroenteritis.

**KEYWORDS :** Acute Gastroenteritis, Electrolytes, Acid-base Imbalance

### INTRODUCTION

Globally, there are nearly 1.7 billion cases of childhood diarrhoeal disease reported every year<sup>1</sup>. Diarrhoea is the third leading cause of childhood mortality in India, and responsible for 13% of all deaths/year in children under 5 year of age<sup>2,3</sup>. Diarrhoea is generally defined as three or more loose or watery stools within a 24-hour period, change in the consistency is more important than frequency. The main cause of death in acute diarrhoea is dehydration, which results from the loss of fluid and electrolytes in diarrheal stools and vomit. The clinical manifestations of acute gastroenteritis are related to the severity of water deficit and the type of electrolyte disturbances. Electrolyte disturbances may remain unrecognized and result in increased morbidity and mortality<sup>4</sup>. Timely recognition, a high index of suspicion and a thorough understanding of common electrolyte abnormalities is important to ensure their correction. In this regard, the biochemical derangement in children with dehydration may be hyponatraemic, isonatremic or hypernatremic<sup>5</sup>. Other biochemical abnormalities observed include hypokalaemia and metabolic acidosis. Often laboratory facilities for investigations are not available or, even if available; there is a considerable time lag in obtaining the results. Consequently, clinical recognition of water and electrolyte disturbances becomes important due to its serious consequences. Therefore purpose of study, an attempt has been made to outline the spectrum of electrolyte and acid base disturbances in children suffering from acute diarrhoea who are all admitted in paediatric ward and paediatric intensive care unit with relation to mortality.

### MATERIAL & METHODS

In this prospective clinical study was conducted in the department of paediatrics, Dr.V.M. Government Medical College, Solapur, Maharashtra, India over a period of 2 years of duration. The study was approved by the ethical committee of Dr.V.M. Government Medical College, and Shri.C.S.M. General Hospital, Solapur, Maharashtra, India. The study group was consisted of 210 paediatric patients admitted in paediatric ward and paediatric intensive care unit with diarrhoea. Patients were enrolled within the study after taking written informed consent from parents.

### Inclusion Criteria:

- All paediatric patients admitted in paediatric ward and paediatric

intensive care unit with diarrhoea age 1 month to 14 years

### Exclusion Criteria:

- Less than 1 month,
- More than 14 years,
- Acute gastroenteritis due to non infective causes,
- Chronic diarrheea,
- HIV.

Detailed clinical history including associated symptoms was noted and history regarding sensorium, activity, water intake, urine output in last 24 hours was noted. Systemic examination was done with assessment of severity of dehydration, nutritional status, sensorium and immunization status were noted in preformed proforma. Laboratory investigation like Hemogram, serum Sodium, Potassium, Creatinine, Urea and ABGA (Arterial Blood Gas Analysis) were done. All the patients were treated according to WHO guidelines for treatment of acute gastroenteritis. The electrolyte disturbances and acid base abnormalities detected were treated according to standard treatment protocol. The data collected was entered in MS excel, compiled, tabulated, analyzed and subjected to statistical tests done using SPSS. Analysis was done using SPSS. The data collected was evaluated using standardized statistical methods i.e. Mean, Standard Deviation, Pearson Chi-square test in order to derive a logical conclusion.

### RESULTS

A total of 210 children met the inclusion criteria admitted for diarrhoeal disease during the study period. Out of these, 120 (57.14%) were males and 98 (42.86%) were females. The distribution of subjects according to age 25 (11.90%), 110 (52.38%) and 75 (35.72%) of the subjects belonged to 1 month-1 year, 1 year-5 year and 5-14 year respectively. Out of 210 patients, 152 (72.38%) of the subjects had their residence in rural area whereas 58 (27.62%) of the subjects live in urban area. No malnutrition was found in 41 (19.52%) cases, MAM (Moderate Acute Malnutrition) was reported in 145 (69.05%) cases, SAM (Severe Acute Malnutrition) was reported in 24 (11.43%) cases. Out of 210 patients 85.71% patients had some dehydration, 14.29% patients had severe dehydration. Incidence of isonatremia was 74.29%, hyponatremia was 20.47, hypernatremia was 5.24%, hypokalemia was noted in 4.76% of patients. Out of 210 patients

metabolic acidosis was observed in 12.86%. Time taken for correction of electrolyte imbalance after standard treatment was 81.25% of patients in < 24 hour, 10.94% of patients in 24-48 hours, 6.25% patients in >48 hours, not corrected in 1.56% of patients. Out of 210 patients 99.52% recovered and discharged, 0.48% died.

**Table 1: Electrolyte Disturbance With Respect To Serum Sodium, Potassium And Bicarbonate Level**

Serum Sodium Level	Cases	Percentage
Isonatremia	156	74.29%
Hyponatremia	43	20.47%
Hypernatremia	11	05.24%
Serum sodium (Mean±SD)	136.69±10.27	
Potassium Level	Cases	Percentage
Normal	200	95.24%
Hypokalemia	10	04.76%
Hyperkalemia	00	00.00%
Potassium (Mean±SD)	3.96±1.46	
Bicarbonate Level	Cases	Percentage
Normal	183	87.14%
Decreased	27	12.86%

**Table 2: Distribution Of Cases With Respect To Acid Base Imbalance**

pH	Some Dehydration	Severe Dehydration	Percentage
<7.25	00	10	04.76%
7.25-7.35	00	16	07.62%
7.35-7.45	180	04	87.62%
Chi square 29.81 p value 0.0005			

**Table 3: Time Taken For Correction Of Electrolyte Imbalance After Standard Treatment**

Time In Hours	Hyponatremia	Hypernatremia	Hypokalemia	Percentage
<24	39	05	08	81.25%
24-48	03	03	01	10.94%
>48	01	02	01	06.25%
Not corrected	00	01	00	01.56%

## DISCUSSION

The diarrhoeal diseases are responsible for considerable morbidity and mortality in developing world. Dehydration, acid base and electrolyte imbalance are common complications of acute diarrhoeal disease in children<sup>5</sup>. This is due to children having major physiologic differences from adults in their total body surface area, immature renal structures, endocrine structures and higher metabolic rate. Each of these factors predisposes them to developmental variations in fluid and electrolyte balance<sup>6</sup>. Electrolyte disturbances may remain unrecognized and result in increased morbidity and mortality. Timely recognition, a high index of suspicion of common electrolyte disturbances is necessary to ensure their correction. The present study was undertaken, to outline the spectrum of electrolyte and acid base disturbances in children suffering from acute diarrhoea who are all admitted in paediatric ward and paediatric intensive care unit with relationship to mortality.

In our study acute gastroenteritis was more in males 120 (57.14%) as compared to females 90 (42.86%) with the ratio of 1.33:1, a finding which is similar to that reported by Ritika Ghosh Dastidar et al<sup>7</sup>, Srivastava et al<sup>8</sup> and Behera et al<sup>9</sup>. The type of dehydration in diarrhoea had no significant association with gender. In the present study, acute gastroenteritis was found in 25 (11.90%), 110 (52.38%) and 75 (35.72%) of the subjects belonged to 1 month- 1 year, 1 year- 5 year and 5-14 year respectively. Maximum cases were found in the age group of 1-5 year was 110 (52.38%) and minimal cases were found in the age group of 1 month-1 year was 25 (11.90%) which was approximately similar to the study done by Ritika Ghosh Dastidar et al<sup>7</sup> reported acute gastroenteritis among 11.5%, 44%, and 44.5% respectively in the infants, 1-5 yrs, and >5 yrs age group. However the results of the present study differs from those found in studies conducted by Srivastava et al<sup>8</sup> and Behera et al<sup>9</sup> which report a higher incidence of AGE in Infants. The difference may be accounted for by the fact that those with no dehydration usually not requires hospital admission were not taken into account. There was no significant association between age of the child and type of electrolyte disturbances.

152 (72.38%) of the subjects suffered from acute gastroenteritis live in

rural area while 58 (27.62%) of the subjects had acute gastroenteritis reside in urban area, indicating more acute gastroenteritis in rural area as compared to urban area. C. Karsten et al<sup>10</sup> reported that urban area people had more incidence of gastroenteritis compared to the rural area, constitutes 58.83% and 41.16% respectively. This study is contrary to our present study. This can be explained but people living in urban area had better life style and better hygiene maintenance as compared to the subjects of rural people. In this study conducted in tertiary care center, most of the patients seeking medical advice coming from rural area. Majority of the patients belong to MAM was 145 (69.05%). We found significant association in this study between malnutrition and diarrhoea which is similar to the other studies. Dr. Bela H Shah et al<sup>11</sup> showed 79% patient had malnutrition, S.K. Behara et al<sup>9</sup>, Deivanayagam et al<sup>12</sup> observed malnutrition in 89% of patients.

In our study 180 (85.71%) and 30 (14.29%) of the subjects were suffering from some and severe dehydration respectively. Most of the cases belongs to some dehydration 180 (85.71%). The findings were in contrary to study done by Mittal et al<sup>13</sup> who has reported 50.52% of moderate dehydration and 21.95% severe dehydration in their study. But similar with the study reported by Ritika Ghosh Dastidar et al<sup>7</sup> who revealed 166 (83%) of the children had moderate degree of dehydration and 34 (17%) children had severe dehydration.

Isonatremia, hyponatremia and hypernatremia was reported in 156 (74.29%), 43 (20.47%) and 11 (5.24%) of the cases respectively in the present research. Most of the patients had isonatremic dehydration 156 (74.29%). Similar findings were reported by Ritika Ghosh Dastidar et al<sup>7</sup> whom in their study found isonatremic dehydration as the commonest 71.5%, followed by Hyponatremic dehydration 22% and Hypernatremic dehydration 6.5%. Sanatha Krishnan et al<sup>14</sup> and Samadi et al<sup>15</sup> reported the incidence of Hyponatremic dehydration as 25.3% and 20.8% respectively. But study conducted by Dr. Bela H Shah et al<sup>11</sup> reported hyponatremia dehydration 62% was the predominant type of dehydration followed by isonatremic and hypernatremic dehydration was 37% and 2% respectively. Gauchan E et al<sup>16</sup> reported that isonatremic dehydration 70.66% was the predominant one, followed by hypernatremic and hyponatremic dehydration was 19.34% and 10.00% respectively.

In the current research normal, hypokalemia and hyperkalemia was reported in 200 (95.24%), 10 (4.76%) and 0 (0%) of the cases respectively. Majority of the cases in the study had normokalemia followed by hypokalemia. Approximately similar results were shown by Ritika Ghosh Dastidar et al<sup>7</sup> they found 170 children (85%) had normokalemia and 30 cases (15%) had hypokalemia and none of the cases had hyperkalemia. 6 cases of hypernatremia had associated hypokalemia. Dr. Bela H Shah<sup>11</sup> found that normokalemia is more common followed by hypokalemia. Gauchan E et al<sup>16</sup> reported that 90.66% of normal potassium, 02.66% of hyperkalemia and 06.68% of hypokalemia which cannot be explained.

Normal and decreased bicarbonate level was reported in 183 (87.14%) and 27 (12.86%) of the cases respectively in the present research, all cases whose having decreased bicarbonate level having severe dehydration. Majority of the cases had normal bicarbonate level. It is contrary to the study conducted by Teuta Faik et al<sup>17</sup> had more number of patients showed decreased bicarbonate level, 19.3%, 50.00%, 81.80% of mild dehydration, moderate dehydration and severe dehydration respectively. Dr. Bela H Shah et al<sup>11</sup> conducted a study showed that 17% cases of severe dehydration had reduction in bicarbonate level. In present study majority of patients had no acidosis, pH of about 7.35- 7.45 was 184 (87.62%), followed by pH was 7.25- 7.35 was 16 (07.62%), <7.25 was 10 (04.76%) contrary to a study by Dr. Bela H Shah et al<sup>11</sup> where 14% had pH <7.2, 7.2-7.3 was 79% and 7.35-7.45 was 7% more cases were in metabolic acidosis.

In the present study, 52 (81.25%) of dyselectrolytemia was corrected within 24 hours, 7 (10.94%) requires 24-48 hours, only 4 (06.25%) requires >48 hours and 1 (01.56%) patient's hypernatremia was not corrected and that patient expired, was comparable with the study conducted by Dr. Bela H Shah<sup>11</sup> reported 92% of hyponatremia and 81% of hypokalemia was corrected within 24 hours of starting the therapy. Very few percent of dyselectrolytemia required treatment for 48 hours. In the present study, 209 (99.52%) patients got discharged, 1 (00.48%) child who succumbed was 9month old girl who had convulsions, hypernatremia and was severely dehydrated. The child died within 24 hours of hospital stay Ritika Ghosh et al<sup>7</sup> in their study case fatality rate was 0.005%.

**CONCLUSION**

Our study concluded that most common electrolyte imbalance in acute gastroenteritis was hyponatremia followed by hypernatremia and hypokalemia. Metabolic acidosis was common in severe dehydration compared to some dehydration. Mortality in acute gastroenteritis more commonly associated with hypernatremia. Our study showed monitoring of electrolyte and acid base levels to decrease morbidity, mortality in acute gastroenteritis.

**REFERENCES**

1. World Health Organization. Diarrhoeal disease. Available at <https://www.who.int/news-room/fact-sheets/detail/diarrhoeal-disease>.
2. Lakshminarayanan S, Jayalakshmy R. Diarrheal diseases among children in India: Current scenario and future perspectives. *J Nat Sc Biol Med* 2015;6:24-8.
3. Million Death Study Collaborators, Bassani DG, Kumar R, Awasthi S, Morris SK, Paul VK, et al. Causes of neonatal and child mortality in India: A nationally representative mortality survey. *Lancet* 2010;376:1853-60.
4. Dastidar RG, Konar N. A Study of Electrolyte Disturbances in a Child Presenting with Acute Gastroenteritis, with Special Emphasis on Hyponatremic Dehydration-A Hospital based Cross-Sectional Study. *Pediatr Ther* 2017;7(2):322.
5. Webb A, Starr M. Acute gastroenteritis in children. *Aust Fam Physician* 2005; 34: 227-231.
6. Huilan S, Zhen LG, Mathan MM, Mathew MM, Olarte J, Espejo R, et al. Etiology of acute diarrhoea among children in developing countries: A multicentre study in five countries. *Bull World Health Organ.* 1991;69:549-555.
7. Ritika Ghosh Dastidar 1 Niladri Konar 2. A Study of Electrolyte Disturbances in a Child Presenting with Acute Gastroenteritis, with Special Emphasis on Hyponatremic Dehydration-A Hospital based Cross-Sectional Study. *Dastidar and Konar, Pediatr Ther* 2017, 7:2.
8. Srivastava AK, Bhatnagar JK, Prasad BG, Sharma NL. A clinical and aetiological study of diarrhoea in hospitalized children at Lucknow. *Indian J Med Res* 1973;61:596-602.
9. Behera SK, Mohapatra SS, Kar S, Das D, Panda C. Incidence and mortality of hospitalized diarrhoea cases. Part III. *Indian Pediatr* 1980;17:607-612.
10. Karsten C, Baumgarte S, Friedrich AW, Eiff CV, Becker K, Wosniok W, et al. Incidence and risk factors for community-acquired acute gastroenteritis in north-west Germany in 2004. *Eur J Clin Microbiol Infect Dis* 2009;28:935-43.
11. Shah BH, Dhara KG, Mehul RC, Jaydip SP. Study of clinical profile in acute gastroenteritis patients with special reference to dyselectrolytemia. *International Journal of Applied Research* 2016;2(8):255-257.
12. Devignam N Mala, N shafi Ahmad, Jagdish Shankar. Measles associated diarrhoea and Pneumonia in South India. *Indian pediatric* 1994;31:35.
13. Mittal SK, Saxena S, Mundkur N, Srivastava G, Gupta S. Acute diarrhea in malnourished children: Clinical, biochemical and bacteriological profile. *Indian Pediatr* 1980;17:247-254.
14. Santhanakrishnan BR, Kumaresan G, Raju VB. Fluid therapy in infants and children with AGE and dehydration. *Indian Pediatr* 1974;11:345-349.
15. Samadi AR, Wahed MA, Islam MR, Ahmed SM. Consequences of hyponatremia and hypernatremia in children with acute diarrhoea in Bangladesh. *Br Med J (Clin Res Ed)* 1983;286:671-673.
16. Gauchan E, Malla KK. Relationship of renal function tests and electrolyte levels with severity of dehydration in acute diarrhoea. *J Nepal Health Res Counc* 2015;13(29):84-9.
17. Hoxha TF, Azemi M, Avdiu M, Ismaili-jaha V, Grajqevci V, Petrela E. The usefulness of clinical and laboratory parameters for predicting severity of dehydration in children with acute gastroenteritis. *Med Arh.* 2014;68(5):304-307.