



## NON-DRUG NON-INVASIVE TREATMENT IN THE MANAGEMENT OF TENNIS ELBOW

**Harsh Pratap Shishodia**

DNB Orthopaedics, Assistant Professor, Saraswathi institute of medical science Hapur UP India.

**Ramji Lal Sahu\***

MS Orthopaedics, Professor, Saraswathi institute of medical science Hapur UP India. \*Corresponding Author

**ABSTRACT**

**Introduction:** Tennis elbow is a common disorder of upper extremity. Majority of the patients can be treated conservatively. It is the major cause of disability and time off work, after low back pain. The disease impacts upon activities of daily living ultimately leading to a loss of functional independence and quality of life. Purpose: The main purpose of this study was to assess the results of nondrug non-invasive treatment in the management of tennis elbow. **Methods:** This study was conducted in the department of Orthopaedics from December 2010 to December 2015. One hundred ten outpatients of tennis elbow with a mean age of 45 years were studied. They were managed with non-invasive treatment and were followed for twelve months. **Results:** At twelve months follow-up visits, the intensity of tennis elbow pain and disability were assessed by using Quick dash scoring system and Patient-rated elbow evaluation system. Physician global evaluations up to the age of 40 years at twelve months were excellent. At 40 to 60 years of age, it was good to excellent. Over the age of 60 years, it was good. The patient global evaluation was found very good up to the age of 40 years at twelve months follow up, good to very good between 40 to 60 years and over the age of 60 years it was good. **Conclusions:** Nondrug non-invasive interventions can reduce pain and improve function in tennis elbow.

**KEYWORDS :** Non-drug; Non-invasive; Tennis elbow; Treatment.

**INTRODUCTION:**

Tennis elbow is a common pathology of both athletes and non-athletes, affecting 1 to 3 % of the population at large [1, 2]. This condition is most often associated with overuse or a repetitive stress, as opposed to an acute inflammatory reaction. The lack of pathological evidence of inflammation in these types of injuries has led most authors to now refer to this condition as an epicondylitis, abandoning the mislabelled "itis" [3, 4, 5 and 6]. However, the choice of treatment options for this condition is even more controversial. There are many treatment options available to the clinician, but their use is often based on anecdotal evidence. Various treatments ranging from conservative to more invasive measures have been described with varying degrees of success, with no conclusive scientific evidence to support any particular treatment protocol. Although many treatment modalities may be used, few of them rest on scientific evidence and none have really been proven more effective than the others have. The paucity of evidence on treatments for lateral epicondylitis may stem from several sources, including the self-limiting nature of the condition, the lack of pathophysiological data, the methodological shortcomings of the current studies, and the existence of multiple factors which may influence the outcome [1, 7]. The purpose of this study was to find out the outcome results of nondrug non-invasive treatment in the management of tennis elbow.

**METHODS:**

This prospective study was carried out at Orthopaedics department from December 2010 to December 2015. Institutional medical ethics committee approved it. In this series, 110 patients were enrolled. The average age of patients was 45 years (ranging from 15 to 75 years) [Table 1]. The average follow up was done up to twelve months.

**Inclusion criteria**

- Age between 18 to 75 years
- No general illnesses or use of medication

A characteristic history and symptoms of tennis elbow: This is a condition characterized by pain and tenderness at the lateral epicondyle of the humerus due to non-specific inflammation at the origin of the extensor muscles of the forearm. Although, it is sometimes seen in tennis players, other activities such as squeezing clothes, carrying a suitcase etc. are frequently responsible.

Characteristic clinical signs of tennis elbow local tenderness at lateral epicondyle of humerus, Pain is aggravated by putting the extensor tendons to a stretch; for example, by palmer- flexing the wrist and fingers with the forearm pronated. Cozen's test-Painful resisted extension of the wrist with elbow in full extension elicits pain at the lateral elbow. Elbow movements are normal.

X-ray does not reveal any abnormality. CT scan and MRI are also prescribed to exclude other abnormality.

**Exclusion criteria**

Cases were excluded if there had been previous surgery or other elbow pathology such as RA, OA, or radial tunnel syndrome.

A written informed consent was obtained from all the patients; they were explained the treatment plan. Rest and watchful waiting. Sometimes taking a break from the activity(ies) that triggered tennis elbow symptoms is sufficient to alleviate the symptoms. RICE protocol, or the combination of Rest, Ice, Compression, and Elevation is often employed as a first-line treatment for tennis elbow [8-11]. In addition to rest, cloth-covered ice packs can be applied to the affected area for no more than twenty minutes at a time every two to three hours, two to three times per day. ACE bandages, compression sleeves, or other similar devices can be worn on the affected arm, and the arm can be elevated on a cushion, high table, or other type of platform. This protocol can provide pain relief while also reducing swelling and promoting healing. All patients received wrist extensor stretching, ultrasound, cross-friction massage, heat, and ice during their physical therapy visits. Additionally, the Standard Treatment Group performed isotonic wrist extensor strengthening and the Eccentric Group performed isolated eccentric wrist extensor strengthening. The strengthening and stretching exercises were also prescribed as a home exercise program. Treatments were continued until patients had resolution of symptoms or were referred back to their physician with continued symptoms. The isolated eccentric strengthening exercise was performed using a rubber bar (Thera-Band Flex Bar; The Hygenic Corporation, Akron OH) which was twisted using wrist flexion of the uninvolved limb and slowly allowed to untwist with eccentric wrist extension by the involved limb. Each eccentric wrist extensor contraction lasted approximately 4 seconds (i.e., slow release). Both upper extremities were reset for the subsequent repetitions. A 30-second rest period was timed between each set of 15 repetitions and 3 sets of 15 repetitions were performed daily. Intensity was increased by giving the patient a thicker rubber bar if the patient reported no longer experiencing discomfort during the exercise. Exercise protocols with two or three sets of 10 or 15 repetitions were commonly used. The frequency of exercise ranged from three times a week to twice a day, and the duration of intervention ranged from 2 weeks to 3 months. The results were assessed by quick dash scoring system and the patient-rated elbow evaluation system. The quick dash [12] is a shortened version of the dash scoring system. It consists of 11 items to measure physical function and symptoms in people with any or multiple musculoskeletal disorders of the upper limb. Similar to the dash, each item has five response options (1 = no difficulty; 2 = mild difficulty; 3 = moderate difficulty; 4 = severe difficulty; 5 = unable). From the item

scores, a summative score is calculated. The final score ranges between 0 (no disability) and 100 (the greatest possible disability). Only one missing item can be tolerated, and, if two or more items are missing, the score cannot be calculated [13]. The patient-rated elbow evaluation [14] (pree) consists of two sections investigating pain and function. All questions are scored on a 10-point scale. The pain section has four questions that rate pain from 'no pain' to 'worst ever'. In addition, there is a question that rates how often the patient has pain ('never' to 'always'). The scale for the function questions ranges from 'no difficulty' to 'unable to do'. The function section has 11 questions regarding specific activities of daily living, and four questions regarding personal care, household work, occupational work and recreational activities. Higher scores represent worse functioning [15, 16].

**Table 1: Age and sex variations in study group (n=110)**

| Age   | Male | Female | Total |
|-------|------|--------|-------|
| 20-40 | 20   | 22     | 42    |
| 40-60 | 18   | 16     | 34    |
| 60-75 | 16   | 18     | 34    |
| Total | 54   | 56     | 110   |

**RESULTS:**

140 elbows in 110 patients (54 males and 56 females) with tennis elbow admitted to our institute were included in present study. Fifty-six patients (56.11%) were women and fifty-four patients (49.09%) were male. 30 patients had bilateral tennis elbow and 80 patients had unilateral tennis elbow. 90 cases of tennis elbow were found on the right side and 50 cases were seen on the left side. All the patients were divided in three age groups. In the age group between 20-40 years, there were 22 females and 20 males. In the age group between 40-60 years, there were 16 females and 18 males and in the age groups between 60-75 years, there were 18 females and 16 males. The average age of patients was 45 years (ranging from 15 to 75 years) [Table 1]. All patients were followed for twelve months. At twelve months follow-up visits, the intensity of tennis elbow pain and disability were assessed by using Quick dash scoring system and Patient-rated elbow evaluation system [Table 2]. Before treatment pain, disability and unable to do were severe and worst in both the scale in 100% cases. At twelve months follow-up, in the patient-rated elbow evaluation [14] (pree) consists of two sections investigating pain and function. All questions are scored on a 10-point scale. The pain section has four questions that rate pain from 'no pain' to 'worst ever' than before the treatment with significant P value (P = 0.16, 0.73, and 0.079, respectively. After treatment, the subjective overall assessment below the age of 40 years was done, 100% of the patients were given one point. Between 40-60 years, 75% of the patient had one point, 15% had four to six points and 10% had seven to eight points. Over the age of 60 years, 50% of the patients had one point, 30% had four to six points and 20% had four to five points. In Quick dash scoring functional disability scale [12, 13] below the age of 40 years, 100% had full recovery (0% disability). Between 40-60 years, 75% of the patient had full recovery (0% disability), 25% had minor recovery (<20%disability). Above the age of 60 years, 50% had full recovery (0% disability) and 50% had minor recovery (<20%disability). (Table 2) In Global Assessment of tennis elbow, below the age of 40 years, 100% had full improvement. Between 40-60 years, 75% of the patient had full improvement. 25% had minor improvement. Above the age of 60 years, 50% had full improvement and 50% had minor improvement. In Objective Physician global evaluation, below the age of 40 years, 100% had full improvement. Between 40-60 years, 75% of the patient had full improvement. 25% had minor improvement. Above the age of 60 years, 50% had full improvement and 50% had minor improvement. In the patient global evaluation, below the age of 40 years, 100% had no difficulty. Between 40-60 years, 75% of the patient had no difficulty, 25% had minor difficulty. Above the age of 60 years, 50% had no difficulty and 50% had minor difficulty. Below the age of 40 years, at 6 months, complete subjective, functional, and clinical recovery had occurred in almost 100% of the patients. From 40 to 60 years of age at 6 months, complete subjective, functional, and clinical recovery had occurred in almost 75% of the patients. Twenty percent of the patients had minor recovery even at 24 months, but their severity became lowered significantly. Over the age of 60 years at 6 months, complete subjective, functional, and clinical recovery had occurred in almost 50% of the patients, rest 50% had minor recovery even at 24 months, but their severity became lowered significantly. Physician global evaluations up to the age of 40 years at 2 years were excellent. At 40 to 60 years of age, it was good to excellent. Over the age of 60 years, it

was good. The patient global evaluation were found very good up to the age of 40 years at 2-year follow up, good to very good between 40 to 60 years and over the age of 60 years it was good [Table 3].

**Table 2. Pre and post management evaluation of tennis elbow (n=110)**

| Age group | Quick dash scoring system                  |   | Patient-rated elbow evaluation system |   |
|-----------|--|---|---------------------------------------|---|
|           | Before treatment                           | After treatment                                   | Before treatment                      | After treatment   |
| 20-40     | 100% had Severe difficulty to unable To do | 100% had No difficulty                            | 100% had Worst pain and Unable to do  | 100% had No pain and No difficulty                                    |
| 40-60     | 100% had Severe difficulty to unable To do | 75% had no difficulty and 25% had mild difficulty | 100% had Worst pain and Unable to do  | 75% had no pain and difficulty, 25% had mild pain and mild difficulty |
| 60-75     | 100% had Severe difficulty to unable To do | 50% had no difficulty and 50% had mild difficulty | 100% had Worst pain and Unable to do  | 50% had no pain and difficulty, 50% had mild pain and mild difficulty |

**Table 3: Results in study group (n=110)**

| Age group | Subjective overall assessment [Patient-rated elbow evaluation system] | Quick dash scoring system functional disability scale | Global Assessment of tennis elbow    | Physician global evaluation | The patient global evaluation |
|-----------|---|---|--------------------------------------|-----------------------------|-------------------------------|
| 20-40     | 100% full recovery  | 100% full recovery                                    | 100% full recovery                   | Excellent                   | Very good                     |
| 40-60     | 75% full recovery 25% minor recovery                                  | 75% full recovery 25% minor recovery                  | 75% full recovery 25% minor recovery | Good-excellent              | Good-very good                |
| 60-75     | 50% full recovery 50% minor recovery                                  | 50% full recovery 50% minor recovery                  | 50% full recovery 50% minor recovery | Good                        | Good                          |

**DISCUSSION:**

Tennis elbow (Lateral epicondylitis) is an overuse injury involving the extensor muscles, especially in the extensor carpi radialis brevis. Histopathological finding is fibrous granulation tissue at the origin of the common extensor tendon [17] and vascular infiltration and degeneration of the common tendon origin [18, 19]. Others claimed that the main pathology in tennis elbow was entrapment of the anterior interosseous branch of the radial nerve and suggested surgical decompression of the nerve [20, 21]. Tennis elbow occurs most commonly in people aged 40 to 50 years with an equal distribution between men and women [3, 22, and 23]. The dominant arm is involved in 75% of patients, and the incidence most directly relates to playing time in amateur players [3]. In my study fifty-six patients (56.11%) were women and fifty-four patients (49.09%) were male. 30 patients had bilateral tennis elbow and 80 patients had unilateral tennis elbow. 90 cases of tennis elbow were found on the right side and 50 cases were seen on the left side. The average age of patients was 45 years (ranging from 15 to 75 years). It has been reported that nearly 50% of all tennis players over 35 years old and 60% of players over 50 years old suffer from tennis elbow at some point in their career [3]. Most such injuries are related to direct trauma or repetitive stress, and account for a significant amount of "down time" for the athlete in sports where the arm is utilized for throwing, catching, or swinging. Elbow biomechanics play a very important role in many overhead

sporting activities including tennis. The amount of tension and the location of the stress within the elbow joint are dependent on the stroke used and the mechanics of each stroke [24]. Electromyographic (EMG) studies of elbow function in tennis have shown that the serving motion creates a larger demand on the elbow than does the groundstroke [24]. That being said, it is well known amongst tennis players that improper backhand mechanics is one of the main causes for elbow injuries. In fact, the incidence of lateral epicondylitis has been clinically linked to a one-handed backhand, and greater wrist extension and pronation activity [24]. Some tennis instructors teach a double-hand backstroke, a stroke, which minimizes wrist pronation, and/or a stroke that avoids leading with the elbow to minimize the potential for improper mechanics. Traditionally the term tennis elbow has been synonymous with lateral epicondylitis. However, the term epicondylitis suggests an inflammatory process, and as Boyer has pointed out – there is no evidence of acute or chronic inflammation in the publications examining the pathological specimens of patients who were operated on for this condition [25]. Repetitive muscle contraction will produce tensile forces within a tendon of an involved muscle, potentially causing micro trauma. If the natural healing process fails, pathological alteration of tissue results in a fibroblastic and vascular response called Angio fibroblastic degeneration [3, 4, 5 and 6]. The pathology of tennis elbow is thus most likely to be Angio fibroblastic degeneration at the origin of the wrist extensors, and more suitably referred to as lateral epicondylitis [3, 4, 5 and 6]. The current understanding of this condition places the specific pathology at the extensor carpi radialis brevis [3, 4, 26 and 27]. The origin of the extensor carpi radialis brevis is covered by the extensor carpi radialis longus and the extensor communis origin. In fact, the common extensor origin consists of the fused tendons of extensor carpi radialis brevis, extensor digitorum, extensor digiti minimi, and extensor carpi ulnaris. Biomechanical studies of tensile force at the lateral epicondyle further indicate that stretching extensor carpi radialis brevis, extensor digitorum communis and the superficial head of the supinator produce large increases in tensile force at the epicondyle [28]. Obviously, a thorough understanding of the anatomical arrangement of these muscles and their specific actions is necessary to make a correct diagnosis. Additionally, it is important to rule out other differential diagnoses such as capitellum fracture, lateral collateral ligament injury, osteochondritis dissecans, posterior interosseus nerve syndrome, radial head fracture and synovitis [26]. Radiographic analysis of lateral epicondylitis may reveal calcification along the lateral epicondyle however, radiographs, as an initial step in diagnosing lateral epicondylitis is not necessary [29]. On the other hand, a diagnostic ultrasound of the common extensor origin can be used to confirm lateral epicondylitis in patients with elbow pain and add additional information in regards to the severity [30]. Most authors suggest that over 90% of patients will respond to conservative care, which may include rest, bracing, strengthening, therapeutic modalities, and steroid injections [3, 31, 32, 33 and 34]. In my study blow the age of 40 years, at 6 months, complete subjective, functional, and clinical recovery had occurred in almost 100% of the patients. From 40 to 60 years of age at 6 months, complete subjective, functional, and clinical recovery had occurred in almost 75% of the patients. Twenty percent of the patients had minor recovery even at 24 months, but their severity became lowered significantly. Over the age of 60 years at 6 months, complete subjective, functional, and clinical recovery had occurred in almost 50% of the patients, rest 50% had minor recovery even at 24 months, but their severity became lowered significantly. Additionally, it has been reported in cases where surgery was required that over 90% of patients responded well [35]. The attempted meta-analysis in 1992 by Labelle et al. reviewed 185 articles on the subject of tennis elbow treatment, however only a single paper was considered to be of a good quality design for controlled therapeutic trials. They concluded that there was insufficient evidence to support any single current method of treatment [36]. This conclusion was reiterated even more recently in the meta-analysis by Bisset et al. that identified 28 randomized controlled trials, which met their minimum criteria [7]. These authors suggested that there was a lack of evidence for the long-term benefit of physical interventions in general [7]. There have been a number of studies comparing therapeutic modalities with placebo for the treatment of soft tissue injuries such as lateral epicondylitis. There is insufficient evidence to support the use of most physiotherapy interventions and only weak evidence for the efficacy of therapeutic ultrasound in the treatment of tennis elbow [7, 36, 37 and 38]. Basford et al. assessed patients for pain, tenderness to palpation, grip strength, medication usage, and subjective perception of pain after a double masked, placebo controlled, randomized trial utilizing a low intensity laser. The results

of this study showed that there were no significant differences, and they concluded that there was no demonstrable beneficial effect of laser therapy [39]. The 2004 systematic review for the efficacy of splinting for lateral epicondylitis identified early positive, but not conclusive evidence supporting the effectiveness of splinting [40]. Similarly, there have been conflicting results on the use of braces and orthotic devices, which may be useful in the initial stages of therapy [41, 42 and 43]. The ability to control the pain associated with lateral epicondylitis may be achieved through acupuncture. A recent systematic review suggested that acupuncture was effective in the short-term relief of lateral epicondyle pain [44]. The Fink et al. randomized controlled trial for chronic epicondylitis also showed that real acupuncture points showed a reduction of pain and an improvement of function at early follow-up [45]. More long-term follow-up would be useful to assess whether acupuncture has a greater role than simply pain modulation. In addition to the acupuncture findings, manipulations and/or mobilizations have been suggested to have a hypoalgesic effect. The works of Strujis et al. and Paungmal et al. have shown that manipulation of the wrist and mobilization of the elbow may play a role in the management of the pain associated with lateral epicondylitis [46, 47]. The preliminary evidence does suggest that manipulations and mobilizations may have some positive effects in the reduction of pain and improvement of function [37]. Historically, a popular choice for treating tendonitis had been deep friction massages. However as evidenced by the 2002 Cochrane review there is simply not a large enough sample size to draw any conclusions in regards to control of pain or improvement in function [48]. The concepts of cross-friction techniques have since evolved into an augmented soft tissue mobilization, more commonly known as the “Graston Technique Instrument-Assisted Soft Tissue Mobilization” or simply Graston [49]. The Graston protocol for epicondylitis uses specifically designed stainless steel instruments, which are moved with multidirectional strokes around the bony prominence of the elbow. Preliminary studies utilizing this Graston technique have shown promising results when compared to a traditional physiotherapy protocol in the treatment of lateral epicondylitis [50]. Perhaps the most popular of soft tissue techniques to gain recent notoriety is Active Release Technique or ART®. This therapy is based on the observation that the anatomy of the forearm has traversing tissues situated at oblique angles to one another that are prone to reactive changes producing adhesions, fibrosis and local edema and thus pain and tenderness [51, 52]. During active release therapy, the clinician applies a combination of deep digital tension at the area of tenderness and the patient actively moves the tissue through the adhesion site from a shortened to a lengthened position [51, 52]. For example, in order to treat extensor carpi radialis brevis, the clinician applies proximal tension distal to the lateral epicondyle while the patient extends the elbow and pronates and flexes the wrist [51]. In my study physician, global evaluations up to the age of 40 years at 2 years were excellent. At 40 to 60 years of age, it was good to excellent. Over the age of 60 years, it was good. The patient global evaluation was found very good up to the age of 40 years at 2-year follow up, good to very good between 40 to 60 years and over the age of 60 years it was good.

## CONCLUSIONS:

The majority of tennis elbow patients can be treated with nondrug non-invasive forms of treatment, and only selected cases may benefit from more invasive operative treatments. Tennis elbow is certainly a challenging musculoskeletal condition to treat and this is largely due to the lack of definitive evidence for the clinical efficacy of the myriad of treatment approaches seen within the literature.

## Acknowledgement: nil

## Conflict Of Interest: nil

## REFERENCES:

- Hong QN, Durand MJ, Loisel P. Treatment of lateral epicondylitis: where is the evidence? *Joint Bone Spine*. 2004; 71(5):369–373. [PubMed]
- Walker-Bone K, Palmer KT, Reading I, Coggon D, Cooper C. Prevalence and impact of musculoskeletal disorders of the upper limb in the general population. *Arthritis Rheum*. 2004; 51(4):642–651. [PubMed]
- Ollivere CO, Nirschl RP. Tennis elbow: current concepts and rehabilitation. *Sports Med*. 1996; 22(2):133–139. [PubMed]
- Regan W, Wold LE, Conrad R, Morrey BF. Microscopic histopathology of chronic refractory lateral epicondylitis. *Am J Sports Med*. 1992; 20:746. [PubMed]
- Kraushaar BS, Nirschl RP. Tendonosis of the elbow (Tennis Elbow). Clinical features and finding of histological, immunohistochemical, and electron microscopy studies. *J Bone Joint Surg Am*. 1999; 81:259–278. [PubMed]
- Nirschl RP. Elbow tendonosis/tennis elbow. *Clin Sports Med*. 1992; 11:851–870. [PubMed]

7. Bisset L, Paungmali A, Vincenzino B, Beller E. A systematic review and meta-analysis of clinical trials on physical interventions for lateral epicondylalgia. *Br J Sports Med.* 2005; 39:411–422. [PMC free article] [PubMed]
8. Johnson GW, Cadwallader K, Scheffel SB, Epperly T. Treatment of Lateral Epicondylitis. *American Family Physician.* 2007; 76(6): 843-848.
9. Cleland JA, Whitman JM, Fritz JM. Effectiveness of Manual Physical Therapy to the Cervical Spine in the Management of Lateral Epicondylitis: A Retrospective Analysis; *Journal of Orthopaedic & Sports Physical Therapy.* 2004; 34(11), 713-724.
10. Nirschl RP, Rodin DM, Ochiai DH, Maartmann-Moe C. Iontophoretic Administration of Dexamethasone Sodium Phosphate for Acute Epicondylitis: A Randomized, Double-blinded, Placebo-controlled Study; *American Journal of Sports Medicine.* 2003; 31(2), 189-195.
11. Nirschl RP. Elbow Tendonitis/Tennis Elbow. *Clinical Sports Medicine.* 1992; 11, 851-870. ©
12. Beaton DE, Wright JG, Katz JN. Development of the Quick-DASH: comparison of three item-reduction approaches. *J Bone Joint Surg Am* 2005; 87:1038-1046. CrossRefMedline
13. Gummesson C, Ward MM, Atroshi I. The shortened disabilities of the arm, shoulder and hand questionnaire (QuickDASH): validity and reliability based on responses within the full-length DASH. *BMC Musculoskelet Disord* 2006; 7:44. CrossRefMedline
14. MacDermid JC. Outcome evaluation in patients with elbow pathology: issues in instrument development and evaluation. *J Hand Ther* 2001; 14:105-114. Medline
15. Dowrick AS, Gabbe BJ, Williamson OD, Cameron PA. Outcome instruments for the assessment of the upper extremity following trauma: a review. *Injury* 2005; 36:468-476. CrossRefMedlineWeb of Science
16. MacDermid JC, Michlovitz SL. Examination of the elbow: linking diagnosis, prognosis, and outcomes as a framework for maximizing therapy interventions. *J Hand Ther* 2006; 19:82-97. CrossRefMedlineWeb of Science
17. Goldberg EJ, Abraham E, Siegel J. The surgical treatment of chronic lateral humeral epicondylitis by common extensor release. *Clin Orthop* 1988; 233: 208-212.
18. Coonrad RW, Hooper WR. Tennis elbow : Its course, natural history, conservative and Surgical management. *J Bone Joint Surg* 1973; 55-A: 1177-1182.
19. Nirschl RP, Petrone FA. Tennis elbow: The surgical treatment of lateral epicondylitis. *J Bone Joint Surg* 1979; 61-A: 832-836.
20. Enzenauer RJ, Nordstrom DM. Anterior interosseous nerve syndrome associated with forearm band treatment of lateral epicondylitis. *Orthopedics* 1991; 14: 788-790.
21. Wilhelm A. Treatment of therapy refractory epicondylitis lateralis humeri by denervation. *Handchir Mikrochir Plast Chir* 1999; 31: 291-302.
22. Terry S. ed. *Campbell's Operative Orthopaedics*, 9th ed. St. Louis, USA: Mosby-Year Book; 1998.
23. Frostick SP, Mohammad M, Ritchie DA. Sports Injuries of the Elbow. *Br J Sports Med.* 1999; 33:301–311. [PMC free article] [PubMed]
24. Lofice J, Fleisig GS, Zheng N, Andrews JR. Biomechanics of the elbow in sports. *Clin Sports Med.* 2004; 23(4):519–530. [PubMed]
25. Boyer MI, Hastings H. Lateral tennis elbow: "Is there any science out there?" *J Shoulder Elbow Surg.* 1999; 8(5):481–491. [PubMed]
26. Conway JE. Clinical evaluation of elbow injuries in the athlete. *J Musculoskeletal Med.* 1998; 15(10):43–52.
27. Rettig AC. Elbow, forearm, and wrist injuries in the athlete. *Sports Med.* 1998; 25(2):115–130. [PubMed]
28. Erak S, Day R, Wang A. The role of the supinator in the pathogenesis of chronic lateral elbow pain: a biomechanical study. *J Hand Surg (Br)* 2004; 29(5):461–464. [PubMed]
29. Pomerance J. Radiographic analysis of lateral epicondylitis. *J Shoulder Elbow Surg.* 2002; 11(2):156–157. [PubMed]
30. Connell D, Burke F, Combes P, McNealy S, Freeman D, Pryde D, Hoy G. Sonographic examination of lateral epicondylitis. *AJR Am J Roentgenol.* 2001; 176(3):777–782. [PubMed]
31. Grundberg AB, Dobson JF. Percutaneous release of the common extensor origin for tennis elbow. *Clin Orthop.* 2000; 376:137–140. [PubMed]
32. Servier TL, Wilson JK. Treating lateral epicondylitis. *Sports Med.* 1999; 28:375–380. [PubMed]
33. Thomas DR, Plancher KD, Hawkins RJ. Prevention and rehabilitation of overuse injuries of the elbow. *Clin Sports Med.* 1995; 14(2):459–477. [PubMed]
34. Brown M. The older athlete with tennis elbow : rehabilitation considerations. *Clin Sports Med.* 1995; 14(1):267–273. [PubMed]
35. Nirschl RP, Petrone FA. Tennis Elbow: the surgical treatment of lateral epicondylitis. *J Bone Joint Surg Am.* 1979; 61:832–839. [PubMed]
36. Labelle H, Guibert R, Joncas J, Newman N, Fallaha M, Rivard CH. Lack of scientific evidence for the treatment of lateral epicondylitis of the elbow. *J Bone Joint Surg Br.* 1992; 74:646–651. [PubMed]
37. Trudel D, Duley J, Zastrow I, Kerr EW, Davidson R, MacDermid JC. Rehabilitation for patients with lateral epicondylitis: a systematic review. *J Hand Ther.* 2004; 17(2):243–266. [PubMed]
38. Smidt N, Assendelft WJ, Arola H, Malmivaara A, Greens S, Buchbinder R, van der Windt DA, Bouter LM. Effectiveness of physiotherapy for lateral epicondylitis: a systematic review. *Ann Med.* 2003; 35(1):51–62. [PubMed]
39. Basford JR, Sheffield CG, Cieslak KR. Laser Therapy: A randomized, controlled trial of the effects of low intensity Nd: YAG irradiation of lateral epicondylitis. *Arch Phys Med Rehabil.* 2000; 81:1504–1510. [PubMed]
40. Borkholder CD, Hill VA, Fess EE. The efficacy of splinting for lateral epicondylitis: a systematic review. *J Hand Ther.* 2004; 17(2):181–199. [PubMed]
41. Struijs PA, Kerkhoffs GM, Assendelft WJ, Dijk CN. Conservative treatment of lateral epicondylitis: brace versus physical therapy or a combination of both – a randomized clinical trial. *Am J Sports Med.* 2004; 32(2):462–469. [PubMed]
42. Struijs PA, Smidt N, Arola H, van Dijk CN, Buchbinder R, Assendelft WJ. Orthotic devices for tennis elbow: a systematic review. *Br J Gen Pract.* 2001; 51(472):924–929. [PMC free article] [PubMed]
43. Struijs PA, Smidt N, Arola H, van Dijk CN, Buchbinder R, Assendelft WJ. Orthotic devices for tennis elbow. *Cochrane Database Syst Rev.* 2001; (2) [PubMed]
44. Trinh KV, Phillips SD, Ho E, Damsa K. Acupuncture for the alleviation of lateral epicondyle pain: a systematic review. *Rheumatology.* 2004; 43(9):1085–1090. [PubMed]
45. Fink M, Wollenstein E, Karst M, Gehrke A. Acupuncture in chronic lateral epicondylitis: a randomized controlled trial. *Rheumatology.* 2002; 41(2):205–209. [PubMed]
46. Struijs PA, Damen PJ, Bakker EW, Blakenboort L, Assendelft WJ, van Dijk CN. Manipulation of the wrist for management of lateral epicondylitis: a randomized pilot study. *Phys Ther.* 2003; 83(7):608–616. [PubMed]
47. Paungmali A, O'Leary S, Souvlis T, Vincenzino B. Hypoalgesic and sympathoexcitatory effects of mobilization with movement for lateral epicondylalgia. *Phys Ther.* 2003; 83(4):374–383. [PubMed]
48. Brosseau L, Casimiro L, Milne S, Robinson V, Shea B, Tugwell P, Wells G. Deep transverse friction massage for treating tendinitis. *Cochrane Database Syst Rev.* 2002; (4) [PubMed]
49. Sevier TL, Wilson JK. Treating lateral epicondylitis. *Sports Med.* 1999; 28(5):375–380. [PubMed]
50. Sevier TL. Traditional physical therapy vs. graston technique augmented soft tissue mobilization in treatment of lateral epicondylitis. *J Am College Sports Med.* 1995; 27(5)
51. Mooney V. Overuse syndromes of the upper extremity: rational and effective treatment. *J Musculoskeletal Med.* 1998; 15(8):11–18.