


Spectrophotometric study of Iron, Nitrate and Phosphate in the water of river Tons at Mau (Maunath Bhanjan)



Environment

KEYWORDS : iron, nitrate, phosphate, spectrophotometer

| | |
|-------------------------|--|
| Shafqat Alauddin | Environmental Research Lab, Shibli National College, Azamgarh-276001,U.P., India |
| Shailendra Yadav | Environmental Research Lab, Shibli National College, Azamgarh-276001,U.P., India |
| Tamheed Firdaus | Environmental Research Lab, Shibli National College, Azamgarh-276001,U.P., India |

ABSTRACT

The present study specially aims to study spectrophotometrically the water quality of river Tons at Mau (Maunath Bhanjan). In the river Tons direct wastes like industrial garbage, household sewage as well as industrial wastes is dumped without treatment. River Tons is continuously been contaminated by the sewage and other types of wastes especially wastes from textile industry. In the present study three parameters were chosen viz. iron, nitrate and phosphate. Although river water of Tons would contain appreciable amount of iron, nitrate and phosphate. This study revealed that there is marked variation in the levels of these parameters during the years of study 2009, 2010 and 2011, sometimes it has been found to be more than WHO maximum contamination level.

INTRODUCTION- In continuation of our study^{1,2} on water quality of different rivers, we have now chosen river Tons of Mau (Maunath Bhanjan). Mau is situated on the fertile plains of the Ganges-Ghaghara doab. It lies between 83° 17' to 84° 52' East & 24° 47' to 26° 17' North. At its north, Ghaghara river is on the border, Ghazipur district is on the south, Ballia district is on the east & Azamgarh district is on the west side. This district represents geographical characteristics of mid Gangetic plain. "Khachari" and "Khadar" are types of soil found in the areas of north of Azamgarh - Ballia Road. In some high places "Bangar" soil is also found. In the southern part of the district, river flow is absent, due to which that area has Bangar type of soil, which is not fertile. The river system of the district is dominated by the Tons River and its tributary Choti Sarju. The Ghaghara River forms the northern border of the district. The main means of irrigation in the district are tubewells. Farmers use tubewell for irrigation purposes. A preliminary survey of the river Tonse at Maunath Bhanjan showed that there were several points of interest. It flows from the suburban areas of the district Maunath Bhanjan to the Mau city. It provides important source of water for the purpose of irrigation and drinking to the people of Maunath Bhanjan. The river water of Tonse would contain appreciable amount of iron, nitrate and phosphate.

Iron is essential nutrient reached into the river water with surface runoff during rains, because in rains iron dissolves from the soil in the runoff water. So during rainy season the level of iron contamination is higher in comparison to other season. Increase in the iron enhances the growth of the algae which can block the sunlight to reach the other aquatic bodies. High concentration of iron may increase the acidity of the water which not good for aquatic life. Iron is not hazardous to human health but it is considered as secondary aesthetic contaminant at high concentration.

Nitrate is an important mineral in the river water. Nitrate in water is present as highly soluble salt, as per the public health standard the maximum contaminant level (MCL) of nitrate for safe drinking water is 10 ppm as nitrate-N or 45 ppm as nitrate. Consumption of drinking water with high nitrate level causes hypertrophy of thyroid gland resulting in hyperthyroidism.

Phosphate enter the river water from human and animal wastes, phosphate rich rocks, laundries wastes, treatment of boiler water and agricultural practices in the form of fertilizers' residues. Organic phosphates are found primarily by biological processes, but also contributed by domestic sewage. Phosphates also occur in bottom sediments and in biological sludge, both in the precipitated inorganic forms and incorporated into organic compounds. Phosphate enhancement in the river and lakes are the cause of algal bloom (eutrophication) which reduces the dissolved oxygen in water and disrupt the natural water cycle. In general phosphate would not hurt human beings and animals unless present in very high concentration. The WHO threshold limit³ for phosphate is 0.05mg/l.

SAMPLING- Water samples were randomly collected at seven sampling stations, details of which are given in table (1). Samples were taken at a distance of 3.0 meters inside the river from the bank and depth of about 0.20 meter before about two months from the start of rainy season, during rainy season and about two months after the rainy season.

**TABLE-1
DETAILS OF SAMPLING STATIONS**

| S.No. | SAMPLING STATION | SAMPLING STATION | LOCATION |
|-------|------------------|------------------|-------------|
| 1 | BELWAN GHAT | SS1 | ENTRY POINT |
| 2 | HANUMAN GHAT | SS2 | - |
| 3 | BAM GHAT | SS3 | - |
| 4 | DHEKULIYA GHAT | SS4 | - |
| 5 | MADAIYA GHAT | SS5 | EXIT POINT |

The samples were collected from the above mentioned sampling station for last three years viz. 2009, 2010 & 2011.

MATERIALS AND METHODS- Reagents were prepared from analytical grade chemicals provided from Qualigens fine chemicals. To minimize the interference of any other ions deionized distilled water is used and all the analysis carried by UV visible Spectrophotometer (UB29).

Determination of iron- Any solution which is colored or can be made colored by adding a complexing agent can be analyzed using a spectrophotometer^{4,5}. Solutions containing iron ions are colorless, but with the addition of ortho-phenanthroline, the iron (II) ions in the sample are immediately complexed to produce a species which is orange in color. More iron (II) ions in a sample will result in a deeper orange color. From data obtained from a series of iron (II) standards, it is possible to be able to determine the amount of iron in an unknown sample.

Dissolve 0.7022 grams of ferrous ammonium sulfate hexahydrate in deionized water. Dilute to 1.00 l. This solution is 100 mg/l Fe +2 (same as 100 ppm). Prepare standard solution 100.0 ml calibrated flasks. To this flask add 5 ml of a 0.25% ortho-phenanthroline solution. Dilute with deionized water to 100.0 ml. Now the solution in calibrated flask was diluted to the mark. After 30 minutes the absorbance of the solution was measured at 510nm (λ max). Now, after pre concentration of collected samples, it was transferred to 100.0 ml calibrated flask, add 5 ml of the 0.25% ortho-phenanthroline solution to the 100.00 ml calibrated flask. Dilute to the mark with deionised water, and after 30 minutes the absorbance was measured. Then concentration of iron was determined in mg/l.

Determination of nitrate- After pre concentration of samples, it was treated with NitraVer® 5 Nitrate Reagent Powder Pillows⁶ in a 25.0 ml sample cell and shaken vigorously for 2 minutes and absorbance of the solution was measured at 500 nm (λ max).

Same procedure was repeated for known solution and absorbance observed was recorded for each solution and concentration of nitrate was determined in mg/l.

Determination of phosphate- The molybdenum blue phosphorous method in conjunction with UV-Visible spectrophotometer is a very sensitive method for the determination of phosphate at 830 nm. Phosphate in the sample can be determined at sub-ppb concentration at 830 nm. Orthophosphate and molybdate ions condense in an acidic solution to form phosphomolybdic acid, upon selective reduction (e.g. with hydrazinium sulphate) a

blue color is produced. The intensity of the blue color is proportional to the amount of phosphate initially incorporated into the heteropoly acid. If the acidity at the time of reduction is 0.5 M in sulphuric acid and hydrazinium sulphate is the reluctant then the resulting blue complex exhibits a maximum absorbance at 830 nm (λ_{max}). Intensity of the color solutions is measured on a UV visible spectrophotometer.

RESULTS AND DISCUSSION- The results obtained from the analysis of different parameters and for different years of study were recorded in table (2).

TABLE-2
DATA FROM DIFFERENT SITES OF RIVER SAI IN JAUNPUR

| S. No. | SAMPLING STATION | 2009 | | | | | | | | | 2010 | | | | | | | | | 2011 | | | | | | | | |
|--------|------------------|-----------------------|------|------|---------|-------|-------|-----------|-------|-------|-----------------------|------|------|---------|-------|-------|-----------|-------|-------|-----------------------|------|------|---------|-------|-------|-----------|-------|-------|
| | | Concentration in mg/l | | | | | | | | | Concentration in mg/l | | | | | | | | | Concentration in mg/l | | | | | | | | |
| | | iron | | | nitrate | | | phosphate | | | iron | | | nitrate | | | phosphate | | | iron | | | nitrate | | | phosphate | | |
| | BRS | DRS | ARS | BRS | DRS | ARS | BRS | DRS | ARS | BRS | DRS | ARS | BRS | DRS | ARS | BRS | DRS | ARS | BRS | DRS | ARS | BRS | DRS | ARS | BRS | DRS | ARS | |
| 1 | SS1 | 4.34 | 5.63 | 4.93 | 13.83 | 15.89 | 14.56 | 12.98 | 16.09 | 13.93 | 4.83 | 6.18 | 5.43 | 14.32 | 16.59 | 15.03 | 13.34 | 16.69 | 14.41 | 5.44 | 6.69 | 6.03 | 14.93 | 17.18 | 15.64 | 13.94 | 17.27 | 14.92 |
| 2 | SS2 | 4.69 | 5.63 | 4.93 | 14.09 | 16.13 | 15.11 | 13.17 | 16.44 | 14.21 | 5.13 | 6.49 | 5.81 | 14.63 | 16.91 | 15.43 | 13.71 | 16.98 | 14.74 | 5.76 | 6.97 | 6.37 | 15.34 | 17.87 | 15.99 | 14.26 | 17.69 | 15.21 |
| 3 | SS3 | 4.98 | 6.39 | 5.34 | 14.58 | 18.29 | 15.58 | 13.58 | 16.95 | 14.64 | 5.69 | 7.04 | 6.32 | 15.21 | 17.38 | 15.89 | 14.27 | 17.38 | 15.04 | 6.11 | 7.24 | 6.58 | 15.49 | 18.24 | 16.34 | 14.68 | 18.11 | 15.81 |
| 4 | SS4 | 5.63 | 7.09 | 5.66 | 15.09 | 18.82 | 18.07 | 14.09 | 17.48 | 15.13 | 6.12 | 7.48 | 6.89 | 15.78 | 17.89 | 16.38 | 14.81 | 17.91 | 15.61 | 6.62 | 7.75 | 7.08 | 15.99 | 18.76 | 16.84 | 13.18 | 18.59 | 16.32 |
| 5 | SS5 | 6.09 | 7.59 | 6.43 | 15.77 | 19.29 | 18.76 | 14.81 | 18.02 | 15.76 | 6.73 | 7.79 | 7.39 | 16.45 | 19.95 | 17.05 | 15.52 | 18.63 | 16.32 | 7.31 | 7.99 | 7.67 | 16.68 | 19.75 | 16.58 | 13.91 | 19.25 | 17.11 |

BRS=BEFORE RAINY SEASON; DRS=DURING RAINY SEASON; ARS= AFTER RAINY SEASON

Data in table (2) revealed that the concentration of iron, phosphate and nitrate is increasing gradually with the subsequent years. It also revealed that the concentrations of these parameters are also increasing gradually from entry point of river in the city to exit point.

The concentration of all the parameters varies seasonally, maximum concentration were reported during rainy season. The variation in the concentration of various reported parameters is gradual. As if we move from the entry point to exit point the concentration of iron is gradually increasing from 4.34 mg/l to 6.09 mg/l before rainy season, from 5.63 mg/l to 7.59 mg/l during rainy season and from 4.93mg/l to 6.43 mg/l after rainy season in the study year 2009 this trend is continued in the subsequent years.

Similar trend of variation in concentration of nitrate and phosphate is seen if move across the entry to exit point as well as from 2009 to 2011.

CONCLUSION- It has been observed during this study that there is gradual increase in concentrations of iron, nitrate and phosphate during subsequent years. The concentration of nitrate is in control as per MCL is concern. But the concentration of iron and phosphate are increasing rapidly, especially concentration of phosphate increased many fold during the years.

The concentration of phosphate is increasing due to uncontrolled use of phosphate fertilizers, dumping of household garbage which contains large percentage of phosphate and industrial wastes.

The concentrations of iron , nitrate and phosphate is increasing with every passing year and it is high time to control the dumping of untreated sewage, agricultural wastes and industrial wastes to the river water to save the river from pollution.

REFERENCE

1. Alauddin Shafqat, Shahma A., Kumar Amresh, Yadav Shailendra and Fatima Afshan , Spectrophotometric determination of Iron, Nitrate and Phosphate in the river Tamsa of Azamgarh city, Journal of Chemistry and Chemical Sciences,2012, Vol.2(4),160-164 | 2. Alauddin Shafqat, Kumar Amresh, Yadav Shailendra and Firdaus Tamheed, Spectrophotometric determination of Iron, Nitrate and Phosphate in the river Gomati of Jaunpur city, International Journal of Scientific Research,2012, Vol.1(6),7-8 | 3. Harrison R.M., Understanding Our Environment and Pollution, Cambridge Royal Society, 1992. | 4. Phillip M., Environmental Chemistry, 2nd Edition, 1996, 46-49 | 5. Skoog and West, Fundamentals of Analytical Chemistry, 2nd Ed., Chapter 29. | 6. Vogel, A Textbook of Quantitative Inorganic Analysis, 3rd Ed., p. 294,310 and 787 | 7. Bassett J., Denny R. C., Jeffery G. H. and Mendham J., Vogel's Text Book of Quantitative Inorganic Analysis, 4th Ed., U.K., 1979. | 8. APHA, AWWA, AND WEF, Standard methods for the Examination of Water and Wastewater, 19th Ed., Washington, DC, American Public Health Association, 1995. | 9. Pittman J. J., Zhang H., Schroder J. L. and Payton M. E., Soil Science and Plant Analysis, 2005, 36, | 1641-1659.