

Histopathological Study of Salivary Gland Lesions



Medical Science

KEYWORDS : Salivary gland lesions, Pleomorphic adenoma, Histopathological

Dr. Ankur N. Sarvaiya	Resident, Department of pathology, Smt. N.H.L. Municipal Medical College, Ahmedabad. Gujarat, India.
Dr. Hinal K Gajjar	Asst.Prof., Department of pathology, Smt. N.H.L. Municipal Medical College, Ahmedabad.Gujarat, India.
Dr.Sahil I. Panjvani	Resident, Department of pathology, Smt. N.H.L. Municipal Medical College, Ahmedabad. Gujarat, India.
Dr.N.R.Shah	Professor, Department of pathology, Smt. N.H.L. Municipal Medical College, Ahmedabad.Gujarat, India.

ABSTRACT

Introduction: Salivary gland tumors are a morphologically and clinically diverse group of neoplasms, which may present significant diagnostic and management challenges.

Aims: To study occurrence, age, sex and site distribution of various salivary gland lesions and compare with findings of other workers.

Methods and Material: The material required for the study was collected from January 2010 to December 2012 from the department of pathology, N.H.L.Municipal medical college, Ahmedabad for 3 years.

Results: Of the total 60 cases, 21 were diagnosed as non neoplastic lesions and 39 as neoplastic lesions of which 24 were benign and 15 were malignant. Among the all salivary gland lesions, pleomorphic adenoma is commonest, and comprises of 30% of all lesions. Of the non neoplastic lesions, sialadenitis is commonest and of malignant neoplastic lesions mucoepidermoid carcinoma is more frequent.

Conclusions: Histopathological examination of salivary gland lesions is the most important method in differential diagnosis of non-neoplastic and neoplastic lesions and in establishing the final diagnosis.

INTRODUCTION:

Salivary glands are the site of origin of many non neoplastic and neoplastic lesions. Salivary gland tumors are a morphologically and clinically diverse group of neoplasms, which may present significant diagnostic and management challenges.

Histologically, salivary gland tumors represent the most heterogeneous group of tumors of any tissue in the body¹. Although almost 40 histologic types of epithelial tumors of the salivary glands exist, some are exceedingly rare and may be the subject of only a few case reports².

AIMS AND OBJECTIVES:

1. To study occurrence of salivary lesions during a period of 3 years 2010-2012.
2. To study age, sex and site distribution of various salivary gland lesions and compare with findings of other workers.
3. To differentiate benign from malignant conditions of salivary gland.
4. We are also interested in recording the spectrum of morphological features of these tumors.

SUBJECTS AND METHODS:

The material required for the study was collected from January 2010 to December 2012 from the department of pathology, N.H.L.Municipal medical college, Ahmedabad for 3 years.

A total of 60 specimens of salivary gland lesions were analyzed, this study includes non neoplastic and neoplastic lesions of the salivary glands. In the study patient's history and clinical details were noted from the original request forms, specimens were fixed in formalin and the sections were taken from the lesion, its margins, surrounding tissues and lymph nodes if any. These sections were stained with hematoxylin and eosin, and in selected cases special stains like PAS was done after mounting on a slide.

The tumors were classified according to world Health Organization's histologic typing of salivary gland tumors.

RESULTS:

During the period from January 2010 to December 2012, a total of 12587 specimens were received for histopathological examinations, of which 60 specimens were of salivary gland lesions,

representing 0.47% as shown in the table 1.

Table 1: Incidence of salivary gland specimens

Particulars	Total	Percentage
Total number of specimens	12587	100
Other lesionsT	12527	99.53
Salivary gland specimens	60	0.47

Of the total 60 cases, 21 were diagnosed as non neoplastic lesions and 39 as neoplastic lesions of which 24 were benign and 15 were malignant as shown in table 2 and graph 2.

Table 2: Incidence of neoplastic & non neoplastic lesions

Total cases	Non neoplastic	Neoplastic benign	Neoplastic malignant
60	21	24	15
100%	25	40	15

In our study, among the all salivary gland lesions, pleomorphic adenoma is commonest, and comprises of 30% of all lesions. Of the non neoplastic lesions, sialadenitis is commonest and of malignant neoplastic lesions mucoepidermoid carcinoma is more frequent as shown in table 3 and graph 2.

Table 3: Incidence of all Salivary gland lesions and their percentage

Lesions	No.	Percentage
Sialadenitis	10	16.66
Mucocele	7	11.67
Lymphoepithelial cyst	1	1.67
Hydatid cyst	1	1.67

Parotid fistula	1	1.67
Necrotising Sialometaplasia	1	1.67
Pleomorphic adenoma	18	30
Warthin's tumour	4	6.66
Myoepithelioma	1	1.67
Lymphangioma	1	1.67
Mucoepidermoid carcinoma	5	8.33
Adenoid cystic carcinoma	3	5
Acinic cell carcinoma	1	1.67
Malignant mixed tumor	2	3.33
Squamous cell carcinoma	2	3.33
Malignant lymphoma	1	1.67
Sarcoma	1	1.67
Total	60	100

Table 4. Age wise distribution of salivary gland lesions

Lesions	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79
Sialadenitis	0	0	2	3	4	1	0	0
Mucocele	0	3	3	1	0	0	0	0
Lymphoepithelial cyst	0	0	1	0	0	0	0	0
Necrotising Sialometaplasia	0	0	0	0	1	0	0	0
Hydatid cyst	1	0	0	0	0	0	0	0
Parotid fistula	0	0	0	0	1	0	0	0
Non neoplastic lesions	1	3	6	4	6	1	0	0
Pleomorphic adenoma	0	3	4	6	3	2	0	0
Warthin's tumour	0	0	0	1	2	1	0	0
Myoepithelioma	0	0	0	0	0	1	0	0
Lymphangioma	1	0	0	0	0	0	0	0
Benign neoplastic lesions	1	3	4	7	5	4	0	0
Mucoepidermoid carcinoma	0	1	0	0	1	2	1	0
Adenoid cystic carcinoma	0	0	0	0	2	1	0	0
Acinic cell carcinoma	0	0	0	0	0	1	0	0
Malignant mixed tumor	0	0	0	0	1	0	0	1
Squamous cell carcinoma	0	0	0	0	1	0	0	1
Malignant lymphoma	0	0	0	0	0	0	1	0
Sarcoma	0	0	1	0	0	0	0	0
Malignant neoplastic lesions	0	1	1	0	5	4	2	2
Total	2	7	11	11	16	9	2	2

From age wise distribution, it is noted that non neoplastic lesions were commonest in 3rd decade of life, benign tumors were common in 4th decade and malignant tumors were common from 5th decade onwards.

Table 5. Gender wise distribution of salivary gland lesions

LESIONS	MALES	FEMALES
Sialadenitis	5	5
Mucocele	5	2
Lymphoepithelial cyst	1	0
Hydatid cyst	1	0
Parotid fistula	1	0
Necrotising Sialometaplasia	1	0
NON NEOPLASTIC LESIONS	14	7
Pleomorphic adenoma	7	11
Warthin's tumour	4	0
Myoepithelioma	0	1
Lymphangioma	1	0
BENIGN NEOPLASTIC LESIONS	12	12
Mucoepidermoid carcinoma	2	3
Adenoid cystic carcinoma	1	2
Acinic cell carcinoma	1	0
Malignant mixed tumor	1	1
Squamous cell carcinoma	2	0
Malignant lymphoma	1	0
Sarcoma	0	1
MALIGNANT NEOPLASTIC LESIONS	8	7
Total	34	26

In our study, male preponderance is seen for all salivary gland lesions with M:F ratio of 1.30:1. For Non neoplastic lesions M:F ratio is 2:1, for benign neoplastic lesions M:F ratio is 1:1 and malignant neoplastic lesions M:F ratio is 1.14:1.

Table 6. Site wise distribution of cases

Sr. no.	Site	Total	Non neo plastic	Neoplastic			% of all tumors	% of all lesions
				Benign	Malignant	Total		
1	Parotid gland	37	6	20	11	31	79.5	61.6
2	Submandibular gland	13	9	3	1	4	10.25	21.6
3	Minor salivary gland	10	6	1	3	4	10.25	16.6
4	Total	60	21	24	15	39	100.00	100.0

From above result, it is evident that common sites of all lesions are parotid(61.67%), submandibular(21.66%) and minor salivary glands (16.67%) in order of frequency. Of all salivary gland tumors parotid gland is commonest (79.5%) whereas non neoplastic lesions are more common in submandibular and minor salivary glands.

DISCUSSION

The results obtained were compared with those of previous studies of well known workers in this field and the significant differences and similarities in results are discussed below.

Among Non neoplastic lesions, chronic sialadenitis is the commonest , among benign tumors, pleomorphic adenoma was common and mucoepidermoid was the commonest in malignant conditions as comparable to G C Fernanes et al³.

Table 7. Frequency of benign and malignant tumors

Series	Total	Benign	Malignant
Ito et al ⁴	335	67.5	32.5
Edda et al ⁵	125	53.4	46.6
Ahmed et al ⁶	100	86	14
Nagarkar et al ⁷	36	75	25
Present study	39	62	38

Benign tumors are more common over the malignant tumors as observed in present as well previous studies. In terms of relative proportions, present study correlates with the Ito et al⁴ study.

Table 8. Age distribution in different series

Series	Benign	Malignant
Thomas et al ⁸	39	47
Ahmed et al ⁶	35.7	42.4
Edda et al ⁵	38	44
Rewusuwan et al ⁹	72	49
Agarwal et al ¹⁰	35	42
Present study	37.25	48.2

Benign tumors are seen at lower age than malignant tumors. Present study correlates with most of the Indian studies.

Table 9. Sex distribution in different series

Series	M:F
Das DK et al ¹¹	1.28:1
Erik G et al ¹²	1.18:1
Ahmed et al ⁶	1.17:1
Edda et al ⁵	1:1.3
Present study	1.04:1

In our study, male to female ratio in all salivary gland tumor is 1.04:1 suggesting slight male preponderance. These findings are consistent with other studies.

Table 10. Sex distribution in benign and malignant tumors in various studies

Series	Benign (M:F)	Malignant (M:F)
Ahmed et al ⁶	1.1:1	1.1:1
Edda et al ⁵	1:1.4	1:1.1
Mohd Ayub ¹³	1.9:1	1:3.3
Present study	1:1	1.1:1

In benign conditions there is equal sex distribution while male were slightly more affected in malignant conditions. Our study is comparable with other studies.

In pleomorphic adenoma M:F ratio is 1:1.57, similar to the findings of Masanja et al (2003)¹⁴ and in Warhins tumor all the cases were noted in males.

In mucoepidermoid carcinoma, M:F ratio is 1:1.5, comparable to findings of Ethunandan M et al¹⁵, and in adenoid cystic carcinoma M:F ratio is 1:2, similar to Ethunandan M et al¹⁵.

Table 11. Site distribution of tumors in different series

Series	Parotid gland	Submandibular gland	Minor salivary glands
Rewusuwan et al ⁹	79%	18%	2%
Edda et al ⁵	34.0%	33.2%	32.8%
Ahmed et al ⁶	70%	18%	12%
Tanaka et al ¹⁶	75%	25%	-
Buddhraj et al ¹⁷	82.8%	13.8%	3.4%
Our study	79.5%	10.25%	10.25%

Among the salivary gland tumors, parotid gland was the most commonly involved in 79.5% of cases, consistent with the study of Rewusuwan et al⁹.

CONCLUSION:

In conclusion from the present study, it is evident that histopathological examination of salivary gland lesions is the most important method in differential diagnosis of nonneoplastic and neoplastic lesions and in establishing the final diagnosis as well as in predicting prognosis, typing, staging and grading of all salivary neoplasms.

REFERENCE

- Brandwein MS, Ferlito A, Bradley PJ, et al. (2002): Diagnosis and classification of salivary neoplasms: pathologic challenges and relevance to clinical outcomes. *Acta Otolaryngol* 7 (122): , 758-64. | 2. Speight PM, Barrett AW (2002): Salivary gland tumours. *Oral Dis* 5 (8): 229-40. | 3. Fernandes GG and Pandit AA (2002). Diagnosis of salivary gland tumours by FNAC. *Bombay Hospital Journal*; 4: 201-206. | 4. Ito F A, Ito K, Vargas P A, Almeida O P and Lopes M A (2005), Salivary gland tumors in a Brazilian population: a retrospective study of 496 cases. *International journal of oral and maxillofacial surg* ;5(34):533-536. | 5. Edda A M Vuhahula (2004), salivary gland tumors in Uganda: clinical pathological study. *African health sciences*. April ;1(4):15-23. | 6. Ahmad S, Lateef M, Ahmad R (2002). Clinicopathological study of primary salivary gland tumors in Kashmir. *JK Practitioner* ;4(9):231-233. | 7. Nagarkar M N, Bansal S, Dass A, Singhal K S, Mohan H (2004). Salivary gland tumours: Our Experience. *Indian J Otolaryngol Head and Neck Surg*; 1 (56): 31-34. | 8. Thomas K M, Hutt M S R and Borgestein J(1980). Salivary gland tumors in Malawi. *Cancer*;46: 2328-2334. | 9. Rewusuwan S, Settakorn J, Mahanupab P (2006). Salivary gland tumors in Maharaj Nakorn Chiang Mai hospital: A retrospective study of 198 cases. *Chiang Mai Med Bull* ; 2(45):45-43. | 10. Agarwal RV, Solanki BR, Junnarkar RV(1967).Salivary gland tumor. *Ind J Cancer* ;4:209-213. | 11. Das K D, Petkar A M, Al Mane M N (2004). Sheik A Z and Malik K M. Role of fine needle aspiration cytology in the diagnosis of swellings in the salivary gland regions: a study of 712 cases. *Med Princ Pract*; 13: 95-106. | 12. Erik G. Cohen, MD; Patel S G, MD; Lin O, MD; Boyle J O, MD; Kraus D H, MD (2004). Fine-Needle Aspiration Biopsy of Salivary Gland Lesions in a Selected Patient Population; *Arch Otolaryngol Head Neck Surg*;130:773-778. | 13. Mohammed Ayub M, Zahid S, Abbas Z and Shoukat M (2008). Morphological pattern of parotid tumors. *Journal of the College of Physicians and Surgeons*;5(18):274-277 | 14. Masanja MI, Kalyanyama BM and Simon ENM (2003). Salivary gland tumours in Tanzania. *East African Medical Journal*; 8(80): 429-431. | 15. Ethunandan M, Pratt CA, Macpherson DW (2002). Changing frequency of parotid gland neoplasms: analysis of 560 tumours treated in a district general hospital. *Ann R Coll Surg Engl*; 84: 1-6. | 16. Tanaka K, Masuda M, Shinden S, Ogata A, Suzuki M (1998). Fine-needle aspiration cytology of tumors of major salivary glands. *Nippon Jibiinkoka Gakkai Kaiho*. Oct;10(101):1283-91. | 17. Buddhraj SN, Pasupathy, Perianayagam (1974). Salivary gland tumors in Pondicherry. *Ind J Surg*;36:235-239. |