


## Phytoremediation of Cadmium From Contaminated Soil by *Vetiveria Zizanioides L.*



### Botany

**KEYWORDS :** Phytoremediation, Arbuscular Mycorrhizae (AM), Vetiver, Cadmium, Hyperaccumulator

\* Gunwal, I

Department of Botany, University of Rajasthan, Jaipur, Rajasthan. \* Corresponding Author

Sharma, K.C

Department of Botany, University of Rajasthan, Jaipur, Rajasthan.

Mago, P

Department of Botany, Sri Aurobindo College, University of Delhi, New Delhi

### ABSTRACT

Industrial waste is one of the most important sources of contamination in the surface environment. The impact of heavy metals in soils, plants, animals and humans is due to the unabated expansion of toxic effects. The toxic wastes generated are treated by physicochemical processes in which "Bioremediation" is the microbial clean up approach. Plant based remediation (Phytoremediation) is an emerging biotechnological tool, for cleaning metal polluted or contaminated soil. The present study deals with phytoremediation of metal polluted soil collected from Delhi and surrounding areas utilizing Vetiver grass, which is one of the best "Hyperaccumulators" for the phytoremediation of metal polluted sites. *Vetiveria zizanioides* (Linn) Nash) were grown in three types of experimental plots containing sterilized soil with a known quantity of Cd salts as present in polluted soils (i) Control without any kind of AM spores, (ii) AM inoculum prepared from spores extracted from non Cd polluted soils, (iii) AM inoculum prepared from spores extracted from Cd polluted soils. Plants were observed for their growth and harvested after 3 months. The soil and plant samples were analyzed for cadmium (Cd) using Atomic Absorption Spectrophotometry (AAS) at regular intervals i.e. every 15 days till harvesting of experimental plant. The impact of cadmium on the biomass of the plant was investigated. Phytoremediation is found to be highly efficient in remediating the heavy metal polluted sites.

### INTRODUCTION:-

Industrial waste is one of the most important sources of contamination in the surface environment. Numerous studies of environmental contamination due to industrial wastage activities have been undertaken to further the understanding of the impacts of heavy metals and metalloids in soils, plants, animals and humans. The unabated expansion of chemical industries and technologies indeed left a trail of anxiety and global concern. The toxic metals may be absorbed by plants grown in contaminated soil which then accumulate in animals eating those plants perhaps reaching to chronic toxic levels. Most heavy metals become quite insoluble in soil at pH 6 or more. Cadmium, being highly soluble than other heavy metals, is a frequent contaminant. Other metals such as Ni, Cu, Mo and Zn are also soluble but to a lesser extent [21]. Soil pollution by heavy metals such as mercury, cadmium, chromium and lead are of great concern to public health [17]. The source of heavy metal in plant is the environment in which they grow and their growth medium (soil) from which heavy metals are then taken up by roots or foliage of plants [24]. Plants growing in polluted environment can accumulate heavy metals at high concentration causing serious risk to human health when consumed [2,4,32].

Cadmium is Natural. Cadmium (elemental symbol Cd and CAS registry number 7440-43-9) is a member, along with zinc and mercury, of Group 12 (CAS IIB) of the Periodic Table of the Elements. It is generally characterized as a soft, ductile, silver-white or bluish-white metal, and is listed as 64<sup>th</sup> in relative abundance amongst the naturally occurring elements. Cadmium is often described as a "heavy metal", but the term has no meaning in fact since it possesses an average density, atomic number and atomic weight compared to other metals

Phytoremediation is the use of certain plants to clean up soil, sediment, water and polyaromatic hydrocarbons (PAHs). Phytoremediation is an aesthetically pleasing mechanism that can reduce remedial costs, restore habitat and clean up contamination in place rather than entombing it in place or transporting the problem to another site. Phytoremediation can be used to clean up contamination in several ways such as Phytovolatilization, Microorganism stimulation, Phytostabilization, Phytoaccumulation or extraction and Phytodegradation by plants [12, 15]. The primary motivation behind the development of phytoremediative technologies is the potential for low cost remediation [14]. Use of the "hyperaccumulators" for the phytoremediation of metalpolluted sites was first suggested by Chaney [8] However, hyperaccumulators were later believed to have limited po-

tential in this area because of their small size and slow growth, which limit the speed of metal removal [10, 11, 13]. By definition, a hyperaccumulator must accumulate at least 100 mg g<sup>-1</sup> (0.01% dry wt [25,33]. Vetiver grass (*Vetiveria zizanioides* L. Nash) has been one among the well documented grasses to have strong resistance to the execrable environment and to be able to survive in high concentrations of heavy metals [34].

Although certain plants like *Brassica juncea* (cv. Vardan), *Vetiveria zizanioides*, *Viola sp.* possess a higher potential for removing Cd from moderately contaminated soil and could be used as a hyper accumulator for contaminated soil remediation. Recently vetiver grass, due to its ecofriendly nature, found a new use for phytoremediation of contaminated sites. Vetiver grass is both a xerophyte and a hydrophyte and, once established, is not affected by droughts or floods [18]. It is highly tolerant to droughts and water logging, frost, heat, extreme soil pH, solidity, salinity, Al and Mn toxicity. It is also highly tolerant to a range of trace elements such as As, Cd, Cu, Cr, and Ni [29]. It is suitable for the stabilization and rehabilitation and reclaiming of acid sulfate and trace metals contaminated soils, i.e. phytoremediation

The present study hypothesises that *Vetiver sp.*, a plant which shows high mycorrhizal colonization, is a hyper accumulator, with a high biomass, has the potential that mycorrhizal *Vetiver* could enhance Cd uptake due to more access to soil beyond rhizosphere with the help of mycorrhizal hyphae.

### MATERIALS AND METHODS:

Plants of *Vetiveria zizanioides* were grown under natural conditions in experimental plots of Department of Botany, University of Rajasthan. The soil samples of all the experimental plots were used for analysis of cadmium in soil as well as plant parts at regular intervals. Biomass and growth parameters of plant in all the experimental plots were noted at regular intervals.

### Experimental setup

The experiment was conducted in the Botanical Garden, Sri Aurobindo College, University of Delhi in 3 micro-plots of 10 m<sup>2</sup> area each. The top soil (up to 30 cm depth) was fumigated twice with 0.1% formaldehyde at an interval of 15 days. Then the soil was allowed to dry and the fumigant was dissipated. *Vetiver sp.* will be grown in three types of experimental plots containing sterilized soil with a known quantity of Cd salts as present in polluted soils (i) Control without any kind of AM spores, (ii) AM inoculum prepared from spores extracted from non Cd polluted soils, (iii) AM inoculum prepared from spores extracted from Cd

polluted soils. Clums rate was kept uniform for all treatments and when clums were 15 days old, thinning was done to maintain spacing of 10 cm between the plants and 20 cm within the rows. The plants were allowed to grow and no fertilizer or pesticide was added to the soil during the course of the experiment. Weeding was done mechanically at regular intervals and plots were irrigated with tap water.

T1 – Control Vetiver plant + Cd + without any spores

T2 – Cadmium contaminated soil + AM fungi spores from non contaminated soil + Vetiver plant

T3 – Cadmium contaminated soil +AM fungi spores from contaminated soil+ Vetiver plant

### Description of sampling points

Three different points (A, B and C) 10 km around Delhi were sampled for soil and plant while the fourth sampling point is located in garden of Sri Aurobindo college, Delhi. The samples from this point served as control and the point was designated as D.

### Collection of plant and soil samples

Soil and plant samples each were collected from points A, B, C and D. Plant samples were collected carefully using hand trowel to dig the soil around the plant and the plants were pulled out carefully, ensuring that no part of the root was lost. The different plant samples were kept in different polythene bags and properly labeled. Soil samples were collected from the same point where the plant samples were uprooted. The soil samples were collected to a depth of 15cm using a soil auger. The soil samples were kept in polythene bags and labeled to avoid a mix-up of the different soil samples.

Pilot study has been carried out in the laboratory, Department of Botany, Sri Aurobindo College, University of Delhi, New Delhi-Clums of Vetiver were procured from Central Soil Salinity And Research Institute, Karnal, Haryana and planted in prepared plots (4x4inch) of 15 kg soil capacity containing sandy soil from the different locations in and around New Delhi, India (28.38N, 77.11E, 228 m altitude) during the winter season. Three plots for each treatment were used for cultivating the plants. Cadmium supplied in the form of Cadmium nitrate dissolved in double distilled water (DDW), was mixed thoroughly in the soil. The prepared plots were placed in field conditions (in Sri Aurobindo College) to expose the growing plants to natural environment. Ten plants per plot were maintained. Ninety-day-old plants were analyzed at 15 days interval for investigating the uptake potential. The culms of vetiver grass with root (10 cm) and shoot (20 cm) were selected for the study. This experiment was conducted over a 12-week growth period with three treatments to *Vetiveria zizanioides* [22].

### Estimation of Cadmium in soil and plant parts

#### Soil and plant collection and Preparation-

The soil (loamy sand) used in this study was collected from the experimental plots at regular intervals. The soil was sieved (4 mm) and steam sterilized (100\_C for 1 hour for 3 consecutive days) to eliminate naturally occurring AMF propagules. Root samples along with surrounding soil were taken in polythene bags and stored at 4\*c until they were processed further. The root samples were preserved in Formalin Alcohol Acetone (FAA) in the ratio of 90:5:5 (v/v/v) before clearing and staining.

#### Sample preparation:

Each plant sample was separated into leaves, roots, and stems and then dried at 50°C for 8 hours using an oven. The dried plant samples were milled using a laboratory blender and kept for digestion. Unwanted materials such as stones, leaves and debris were removed from the soil samples by handpicking. The soil sample was further broken down into finer particles using a laboratory mortar and pestle. The soil samples were dried for 8 hours at 80°C using an oven.

#### Sample digestion:

Soil and plant samples were digested before analysis to reduce organic matter interference and allow for the conversion of the metal into a form that can be analyzed by the Atomic Absorption Spectrophotometer (AAS) [28].

#### Plant sample digestion:

Plant samples were digested following the Mixed Acid Procedure of Allen et. al. [3]

#### Reagents-

1. Perchloric Acid, 60%
2. Nitric Acid, concentrated.
3. Sulphuric Acid, concentrated.

#### Procedure-

1. Weigh 0.20-0.50g of air-dried or oven-dried milled plant sample into a 50 ml Kjeldahl flask using a digital weighing balance.
2. Add 1 ml of 60% Perchloric acid, 5 ml nitric acid and 0.5 ml Sulphuric acid to the weighed milled plant sample.
3. Swirl gently and digest slowly at moderate heat on a laboratory hot plate, increasing the heat later, until the white fume evolving from the Kjeldahl flask turned brown.
4. Digest for 10-15 minutes after the appearance of white fumes.
5. The digest was allowed to cool (the cold digest is usually colourless or occasionally pink).
6. Dilute to about 10 ml and boil for a few minutes and then filtered through a Whatman's filter paper (No. 44 paper) into 50 ml volumetric flask, leaving a whitish residue.
7. The filtrate was then made up to 25 ml using distilled water and kept for further analysis.
8. The residual acid is now about 1% (v/v).

#### Soil sample digestion:


The same procedure for the digestion of plant sample was used according to the method of Allen et al. (1974)

### RESULT AND DISCUSSION:

#### RESULTS

The results of the chemical analysis of cadmium (Cd) in soil and plant samples collected are presented in **Figure 1**. The concentration of cadmium is higher in soil samples obtained from the experimental plot I when compared with those obtained from the experimental plots II & III. The concentrations of the heavy metals in the plant tissue were higher in the experimental plots II & III than the experimental plot I. Irrespective of the concentration of the metals and the presence or absence of AM spores the bioaccumulation of the selected metal was higher in the roots than shoots. It was 140% and 101% respectively in cadmium exposed *vetiver* plant. These findings suggested that the soil type could cause the difference in cadmium accumulation in *vetiver* plant. [5] reported that plant species as well as cultivars differ widely in their uptake ability and accumulation of heavy metals.

The average highest cadmium concentration in shoots and roots of vetiver grass was 3.99 ppm & 11.83 ppm. It was lower than the toxic threshold level [32].Cd was found to accumulate more in roots than in shoots (the accumulation rate in shoot/root is from 1.85% to 13.32%). This finding is similar to the results of Vousta. They found that a small amount of Cd was translocated to the shoot.




**Fig 1- Mean Concentration of Cadmium mg/k or ppm in soil**

### samples and plant tissues of *Vetiveria zizanioides* at regular intervals.

Plants of *V. zizanioides* were harvested at regular intervals (15 days) from experimental plots. Roots and shoots of vetiver were taken at regular interval (15 days) and analyzed for Cd uptake every year on 15 DAS (Initial), 45 DAS, 90 DAS (maturity). It was observed that shoot started uptake on 45 DAS and uptake was slow and much less than roots in all the 3 plots studied because shoot showed stunted growth, chlorosis of shoot occurred, no. of fronds decreased and Stability in uptake was observed on 75 DAS and 90 DAS in all the 3 plots studied because the plant attain maturity and started dying

- At the end of cropping it was observed that there is depletion in cadmium from soil in each plot and roots had accumulated more cadmium in comparison to shoots because early chlorosis, stunted growth, decrease in no. of fronds of shoots were observed due to high concentration of cadmium in soil in each plot. Stability in uptake was observed on 75 DAS and 90 DAS in all the 3 plots studied because the plants attain maturity and started dyeing.
- It was observed that roots and shoots taken from plot 2 showed maximum uptake in all years in comparison to roots and shoots taken from plot 1 and plot 3.
- From the above result it is concluded that roots have maximum uptake potential than shoots. ( Fig. 1 )



**Fig. 2 - Comparison of Biomass accumulation in all experimental plots every year**

The plants were harvested and washed by deionised water followed by proper blotting between filter papers. Plants were dried separately in a hot air oven at  $65^{\circ}\text{C} \pm 2^{\circ}\text{C}$  for 72 h. The samples were weighed on an electronic top pan balance so as to obtain the biomass accumulation, which was expressed in g per plant.

Biomass increased with increasing age of plants. A linear increase in the biomass was evident with increase in time period with the maximum biomass accumulation evident at 90 DAS (Days After Sowing) in each plot and at every year. Plants grown in plot 1 produced significantly had less root and shoot biomass than plants in both the plots 2 and 3. The dry weight values indicated that there were no significant differences between plot 1 and plot 3 for shoot weight. However, the total biomass of treatment plants of plot 1 was significantly less than the plot 2 and plot 3.

Note: It was observed that the biomass of root increased consecutively in all plots and every year and it is more than shoot biomass in all treatments. It was observed that the total biomass showed gradual increase in three consecutive treatment and yearly. The results showed that root accumulate more cadmium than shoot. (Fig. 2)

### DISCUSSION:

The concentration of Cd in the soil was higher in soil samples from the industrial site compared with the non-industrial site. There is no doubt that heavy metals are present in soil naturally and non-degradable, and can be accumulated in the plant tissues [3,18] as shown by the concentrations of heavy metals obtained in soils from non-industrial site, but their concentrations can be increased by industrial activities[22]. *Vetiver* is native to India, its environmental application for soil and water conservation is traditionally practiced for longer time. Systematic efforts to develop the *vetiver* grass technology for mitigation of soil erosion and water conservation was first initiated in India; however it was not practiced seriously. Several countries on the other hand, taking cues from the Indian initiative extensively implemented environmental applications of this grass [20]. Currently two main methods for treating contaminated water namely, 'engineering' and 'biological' are being used. The biological method consists of land irrigation, wetland and hydroponics system [9].

As *vetiver* has been found to be highly tolerant to extreme soil condition including heavy metal contamination, the present study was conducted to check the reduction level of Cd contaminants of soil on the plant's ability to tolerate toxic levels of Cd and on the ability to accumulate these heavy metals in roots and shoots. [31] have proved the similar results. It might be concluded that heavy metals in soil even at the higher level to plant growth have no negative effect on *vetiver* root growth but showed negative effect on the shoots and the growth of shoots sustained after some days so as the growth of plant. Our study did not in coordination with the mentioned results. [26]

For successful phytoremediation, the first requisite is high production of plant biomass. Since metal removal is a function of metal concentration in the harvestable biomass, the plant should be able to produce enormous biomass so that it can grow successfully at the contaminated site. Cr uptake is an important parameter in understanding the cellular responses of high HM concentration in plants and is one of the requisites contributing to the success of phytoremediation. The ambient metal concentration in the soil was the major factor influencing the metal-uptake efficiency as the metal uptake was observed to increase with increase in treatment doses Considering the definition of hyperaccumulator given by Foy [6,7,16,25] Indian mustard behaves as a potential Cd hyperaccumulator. The present study is not in accordance with the above mentioned results because plant chosen in the present study has a higher biomass of roots as compared to aerial parts and due to higher biomass of its roots the plant is used for keeping a check on soil erosion. On the other hand, Cd accumulation was enhanced in various plant parts with time. In general, when metal concentration was increased, the amount of metal accumulation in plants increased. Similar results were obtained for *Lemna minor*[19] and water hyacinth [35]

In conclusion, our test plants displayed enhanced tolerance towards a range of Cd concentration besides accumulating high Cd in its various parts, thus serving as a suitable contender for phytoremediation. It is suggested that the remediation potential of *Vetiver* can be tapped to the utmost by periodically harvesting the plant from the site being remediated, avoiding chances of the plant attaining lethal concentration of metal that could lead to oversaturation and hence the damage. The harvested biomass could then be incinerated and disposed off or the accumulated metal could also be recovered for commercial uses and thus recycled and reused.

The results of the present investigation showed that the cd uptake by roots was more in comparison to the shoot and the concentration of cadmium in soil decreased slowly initially but pick

up more later and steadily in three consecutive year of cropping and slightly reached below the permissible level and continuously present study showed is not in accordance with the following studies.

## REFERENCE

- Adeyeye, E.I., 2005. Trace metals in soils and plants from Fadama farms in Ekiti state, Nigeria. *Bulletin of Chemical Society of Nigeria*, 2(1): 1-5. | 2. Akinola, M.O. and T.A. Ekiyoyo, 2006. Accumulation of lead, cadmium and chromium in some plants cultivated along the bank of river Ribila at Odo-nla area of Ikorodu, Lagos state, Nigeria. *Journal of Environmental Biology*, 27(3): 597-599. | 3. Allen S.E., Grimshaw H.W., Parkinson J.A. and Quarmby C. 1974. *Chemical Analysis of Ecological Materials*, 1st edn. Blackwell Scientific Publications, Oxford. | 4. Alloway, B.J., 1990. *Heavy Metal in Soils*. 2nd ed. Glasgow Blackie and Son Publishers, England, pp: 339. | 5. Alloway, B.J., 1995. *Heavy Metals in Soils*. Blackie Academic & Professional, London. | 6. Baker A. J. M. and Brooks R. R., 1989: Terrestrial higher plants which hyperaccumulate metallic elements. A review of their distribution, ecology and phytochemistry. - *Biorecovery* 1; 81-126. | 7. Baker, A. J. M., McGrath, S. P., Reeves, R. D. and J. A. C. Smith, 2000. Metal hyperaccumulator plants: A review of the ecology and physiology of a biological resource for phytoremediation of metal polluted soils p 85-107. In N. Terry and G. Banuelos (ed) *Phytoremediation of contaminated soil and water*. Lewis publishers, Boca Raton, FL. | 8. Chaney, R.L. 1983. Plant uptake of inorganic waste constituents. *Land treatment of hazardous wastes*. Park Ridge, NJ, Noyes Data Corp, pp 50-76. | 9. Chomchalow, N., 2003. The role of vetiver in controlling water quality and treating water quality. An overview with special reference to Thailand. *Au.J.T* 6 (3), 145-161. | 10. Comis, D. 1996. Green remediation: Using plants to clean the soil. *Journal of soil and water conservation*, 51(3), pp 184-187. | 11. Cunningham, S.D., Berti, W.R., and Huang, J.W. 1995. Phytoremediation of contaminated soils. *Trends in Biotechnology*, 13(9), pp 393-397. | 12. Cunningham, S.D., Shann, J.R., Crowley, D.E., and Anderson, T.A. 1997. Phytoremediation of contaminated water and soil. In: Kruger, E.L., Anderson, T.A. and Coats, J.R. eds. *Phytoremediation of soil and water contaminants*. ACS symposium series, Washington, DC, American Chemical Society, 664, Pp 2 - 19. | 13. Ebbs, S.D., Lasat, M.M., Brandy, D.J., Cornish, J., Gordon, R., and Kochian, L.V. 1997. Heavy metals in the environment: Phytoextraction of cadmium and zinc from a contaminated soil. *Journal of Environmental Quality*, 26, Pp 1424-1430. | 14. Ennsley, B.D. 2000, rational for use of phytoremediation. In: Raskin, I. and Ennsley, B.D. eds. *Phytoremediation of toxic metals: using plants to cleanup The environment*. New York, John Wiley & Sons, pp 312. | 15. Flathman, P.E., and Lanza, G.R. 1998. Phytoremediation: Current views on an emerging green technology. *Journal of Soil Contamination*, 7(4), pp 415-432. | 16. Foy CD, Chaney RL, White MC. The physiology of metal toxicity in plants. *Ann Rev Plant Physiol* 1978; 29:511 -66. | 17. Goyer, R.A., 1996. Toxic and essential metal interaction. *Annual Review of Nutrition*, 17: 37-50. | 18. Greenfield, J.C. 1989. Vetiver grass: The ideal plant for vegetative soil and moisture conservation, The World Bank, Washington D.C., USA | 19. Jain R, Srivastava S, Madan VK, Jain R. 2000. Influence of chromium on growth and cell division of sugarcane. *Indian J Plant Physiol*; 5:228-31. | 20. Lavania, U.C., 2004. Vetiver system ecotechnology for water quality improvement and environmental enhancement. *Curr.Sci*, 86, 11-14. | 21. Mathur, A.K., Iijima, K.K., Tandon, S.K., and Dikshith, T.S. 1977. effect of nickel sulfate on male rats. *Bull. Environ. Contam. Toxicol*, 17, pp 241-248. | 22. McDonald, S. 2006. Phytoremediation of Lead-Contaminated Soil using mycorrhizal fungi on heavy-metal uptake by soybeans. *Soil Biol. Biochem.*, 1990, 22, 865-869. | 23. Nriagu, J.O., 1990. Global metal pollution: Poisoning the biosphere. *Environment ENVIRONMENTAL*, 32(9): 7-11, 28-33. | 24. Okonkwo, N.C., J.C. Igwe and E.C. Onwuchekwa, 2005. Risk and health implications of polluted soils for crop production. *African Journal of Biotechnology*, 4(13): 1521-1524. | 25. Reeves, R. D. and Baker, A.J.M. 2000. Metal accumulating plants In: Raskin I and Ennsley B.D. (Eds) *Phytoremediation of toxic metals. Using Plants to Clean Up the Environment*. John Wiley and Sons, Inc. New York pp 193 - 230. | 26. Roongtanakiat, N. and P. Chairaj, 2001. Uptake potential of some heavy metals by vetiver grass. *J.Nat.Sci*.35: 46-50. | 27. Singh, B., 2001. Heavy metals in soils: Sources, Chemical Reactions and Forms in Geo environment. *Proceedings of the 2nd Australia and New Zealand conference on environment geo techniques*. New Castle, New South Wales. | 28. Tandon, H.L.S., 1993. *Methods of analysis of soils, plant, waters and fertilizers*. Fertilizer development and consultation organization. New Delhi. | 29. Truong P, and Claridge J. 1996. Effect of heavy metals toxicities on vetiver growth. Bangkok, Thailand: Vetiver Newsletter, Number 15 | 30. Truong P.N.V. 1999. Vetiver Grass Technology for Mine Rehabilitation. Office of Royal Development Projects Board, Bangkok. *Technical Bulletin* 1999/2 (1999) 12. | 31. Truong, P.N and D. Baker. 1998. Vetiver grass system for environmental protection. *Technical Bulletin No. 1998/1*. Pacifica Rim Vetiver Network. Office of the Royal Development Projects Board, Bangkok, Thailand. | 32. Vousta, D., A. Grimanis and C. Samara, 1996. Trace elements in vegetables grown in an industrial area in relation to soil and air particulate matter. *Environmental Pollution*, 94(3): 325-335 | 33. Watanabe, M.E., 1997. Phytoremediation on the brink of commercialization. *Environ Sci Technol*, 31, pp 182-186. | 34. Xia, H.P., Ao, H.X., and Liu, S.Z. 1999. Application of the Vetiver Ecoengineering for the prevention of highway slippage in South China. In: International Erosion Control Association (eds), *Proceedings of the First Asia Pacific conference on Ground and Water Bioengineering Erosion Control and Slope Stabilization*. Manila, The Philippines, pp 522-527. | 35. Zhu, Y. L., Pilon-Smits, E.A.H., Tarun, A.S., Weber, S.U., Jouanin, L. and Terry, N. 1999: Cadmium tolerance and accumulation in Indian mustard is enhanced by [overexpressing glutamylcysteine synthetase. - *Plant Physiology*. 121; 1169-1177. | | |