

## Prevalence of Hepatitis A in Clinically Suspected Cases of Acute Viral Hepatitis in Tertiary Care Hospital, Rajkot



### Microbiology

**KEYWORDS:** Hepatitis A, Acute viral hepatitis, IgM ELISA, Seroprevalence

**Dr. Nirav Pandya**

Consultant microbiologist, Max Super Speciality Hospital, Dehradun, India.

**Dr. Anil Chaudhary**

Tutor, Microbiology department, P.D.U Medical College, Rajkot, Gujarat, India

### ABSTRACT

*Objective: Hepatitis A is one of the most common infections during childhood, transmitted by faeco oral route in Developing countries. This study was performed to detect the profile of Hepatitis A virus in clinically suspected cases of acute viral Hepatitis attending Max hospital, Dehradun.*

*Material & Methods: Study was conducted in our institute from Jan 2013 to March 2014. Total 310 blood samples of patient suspected of acute viral hepatitis were collected and tested for anti HAV IgM antibody by ELISA method.*

*Findings: Out of 310 samples, total 65 patients (20.97%) were positive for anti HAV IgM antibody among this. 30 (46.15%) were < 5 years of age, 23 (35.38%) were between 6-10 years of age, 5 (7.69%) were between 11-18 years of age and only 7 (10.77%) were between > 18 years of age. 39 (60%) were Male and 26 (40%) were Female. Negative history of HAV vaccine in all cases.*

*Conclusion: Study confirms that, HAV most commonly cause acute viral hepatitis in < 5yr, followed by 6-10 yr. In this present study there is low evidence of HAV infection in adults. HAV is more common in Male compare to female. It is concluded that in addition to other preventive measures, if children are to be vaccinated against hepatitis A they should be immunized against HAV by 1 year of age when the maternal antibodies disappear.*

### Introduction:

Hepatitis A is a disease caused by a RNA virus that is classified as a member of the picornavirus family. Hepatitis A virus (HAV) is usually transmitted by the fecal-oral route, most frequently via contaminated food or water or close contact. The virus is also associated with poor sanitary conditions and low socioeconomic status.[1] HAV causes an acute benign disease that is generally self-limited, rarely fatal and normally limited to the liver. Seventy percent of infected children less than 6 years old are asymptomatic and even when illness occurs, it is usually subclinical. [2] On the other hand, in children above the age of 6, infection is usually symptomatic and accompanied by jaundice in more than 70% of cases. [3] Typical symptoms last less than 2 months, yet prolonged or relapsing disease accounts for 10-15% of cases. [4] The case fatality ratio of Hepatitis A is estimated to be 0.3-0.6% and reaches up to 1.8% in adults older than 50 years of age. [5] Patients with chronic liver disease who acquire hepatitis A are at an increased risk of acute liver failure disease. [6],[7] At this point there is no specific treatment for HAV infection. The Centres for Disease Control and Prevention (CDC) recommends the vaccination of all children at age 1 year as it is the most effective method to prevent hepatitis A infection. [8]

Previous vaccination attempts to reduce the incidence of HAV infection were very successful. Universal vaccination programs had a positive impact not only on the vaccinated groups but also on non vaccinated groups mainly due to a herd immunity effect. [9] In the USA, Hepatitis A vaccine was implemented in 17 states, decreasing the Hepatitis A rate by 88% to a historic low. [10] Vaccination has proven to be safe with no increases in serious adverse events compared with baseline. [11]

### Material & Methods:

#### Study Population

Patients attending Outpatient department or admitted in

P.D.U. medical college, Rajkot, Gujarat, India Dehradun with jaundice or signs and symptoms suggestive of acute viral hepatitis were included in the study.

#### Case Definition and Criteria of Inclusion

Patients with one or more of the following characteristics were included in the study:

1. Acute clinical illness that includes malaise, extreme fatigue, fever, anorexia, vomiting. Combined with right upper quadrant

pain and dark urine.

2. Clinical jaundice and positive bile pigment in urine and elevated total serum bilirubin.

3. History of contact with an acute or known chronic case of viral hepatitis.

### Study Sample

All eligible subjects recorded during the study period extending from From January 2013 to March 2014 were included in the study.

### Sample collection

From January 2013 to March 2014, blood samples were collected from 310 patients suspected of acute viral hepatitis in our institute. Blood samples were collected from all the patients after taking the informed consent. Centrifugation was performed at 5000 rpm to separate the serum. All 310 of the serum samples were tested for anti HAV IgM antibody using commercially available Autobio (anti-HAV (IgM) ELISA, Autobio Diagnostics Co. LTD, China).

### Result:

In our study out of total 310 samples, 65 were positive for HAV each tested for anti HAV IgM antibody to know the recent infection of HAV.

Out of the positive 65 cases 39 were male and 26 were female. 30 (46.15%) were < 5 years of age, 23 (35.38%) were between 6-10 years of age, 5 (7.69%) were between 11-18 years of age and only 7 (10.77%) were between > 18 years of age. There was Negative history of HAV vaccine in all cases.

### Discussion:

Viral hepatitis continues to be a major public health problem in India and other developing countries. Ever since the first epidemics of hepatitis that had occurred in 1955 at Delhi<sup>12</sup>, several outbreaks of hepatitis have continued to occur. In India, available epidemiological data on HAV infection is limited. However, many recent reports have published the changing scenario for sero epidemiological patterns of hepatitis A infection in India<sup>13</sup>. The World Health Organization (WHO) recommends for epidemiological and cost-benefit studies for HAV immunization<sup>14</sup>. Therefore we decided to conduct this study.

In our study HAV infection is more common in male which correlates with previous studies in Hyderabad. [15]

Seroprevalance of HAV in our study shows children < 5 Years are more commonly affected while other study like Mohanavlli *et al* [16] and Dutta *et al* [17] says HAV is more common in child between 5-10 year.

**Table-1: Prevalence of HAV in clinically suspected cases of acute viral Hepatitis**

|                              | <5 year | 6-10 year |
|------------------------------|---------|-----------|
| Mohanavlli <i>et al</i> (16) | 5.9%    | 94.1%     |
| Dutta <i>et al</i> (17)      | 13.4%   | 86.6%     |
| Present study                | 46.15%  | 35.38%    |

The increase in number of infected children in the age group of < 5 years compared to other study could be possibly due to very poor sanitary and hygienic conditions and possibly the occurrence of this epidemiological Shift in this region. At this point, the reasons can only be postulated as the present study was not done at the community level. Probably our HAV vaccination policy needs a review.

## Conclusion:

### Study confirms that,

- HAV most commonly cause acute viral hepatitis in < 5yr, followe by 5-10 yr. In this present study only 7 (10.77%) HAV infection in adults.
- HAV is more common in Male compare to female.

It is concluded that in addition to other preventive measures, if children are to be vaccinated against hepatitis A they should be immunized against HAV by 1 year of age when the maternal antibodies disappear.

## REFERENCE

1. Jacobsen KH, Koopman JS. The effects of socioeconomic development on worldwide hepatitis A virus seroprevalence patterns. *Int J Epidemiol* 2005;34:600-9. || 2. Hadler SC, Webster HM, Erben JJ, Swanson JE, Maynard JE. Hepatitis A in day-care centers. A community-wide assessment. *N Engl J Med* 1980;302:1222-7. || 3. Lednar WM, Lemon SM, Kirkpatrick JW, Redfield RR, Fields ML, Kelley PW. Frequency of illness associated with epidemic hepatitis A virus infection | in adults. *Am J Epidemiol* 1985;122:226-33. || 4. Glikson M, Galun E, Oren R, Tur-Kaspa R, Shouval D. Relapsing hepatitis A. Review of 14 cases and literature survey. *Medicine (Baltimore)* 1992;71:14-23. || 5. Novotny TE. US Department of Health and Human Services: A need for global health leadership in preparedness and health diplomacy. *Am J Public Health* 2006;96:11-3. || 6. Tandon BN. Ninth Triennial International Symposium on Viral Hepatitis and Liver Diseases. Rome 21 st -25 th April 1996. *Trop Gastroenterol* 1996;17:77-83. || 7. Bell BP. Hepatitis A and hepatitis B vaccination of patients with chronic liver disease. *Acta Gastroenterol Belg* 2000;63:359-63. || 8. Centers for Disease Control and Prevention. | Hepatitis A Information for Health Professionals. Available from: <http://www.cdc.gov/hepatitis/HA/index.htm>. [Last accessed on 2012 Jun 05]. || 9. Domínguez A, Oviedo M, Carmona G, Batalla J, Bruguera M, Salleras L, et al. mpact and effectiveness of amass hepatitis A vaccination programme of preadolescents seven years after introduction. *Vaccine* | 2008;26:1737-41. || 10. Wasley A, Samandari T, Bell BP. Incidence of hepatitis A in the United States in the era of vaccination. *JAMA* 2005;294:194-201. || 11. Westblom TU, Gudipati S, DeRousse C, Midkiff BR, Belshe RB. Safety and immunogenicity of an inactivated hepatitis A vaccine: Effect of dose and vaccination schedule. *J Infect Dis* 1994;169:996-1001. || 12. Jindal M, Rana SS, Gupta RK, Das K, Kar P. Serological study of hepatitis A virus infection amongst the students of a Medical College in Delhi & evaluation of the need of vac-cination. *Indian J Med Res* 2002; 115:1-4. || 13. Lakshmi TM, Vaithilingam A, Franklin A, Reddy EP. The prevalence of serological markers of viruses causing acute hepatitis in South Indian population. *International Journal of Biological Medicine Research* 2011; 2(4):925-928. || 14. Hepatitis A vaccines: WHO position paper. *Weekl Epide-miol Rec* 2000; 75:38-42. || 15. Syed R *et al*. Seroepidemiology of hepatitis A virus in Hyderabad, South India. *J Med A lli e d S c i 2 0 1 2 ; 2 ( 2 )* : 58-61. || 16. Dutta AK, Aggarwal A, Kapoor AK, Ray GN, Batra | S.Seroepidemiology of hepatitis A in children in Delhi. *Indian J Pediatr* 2000; 67 : 77-9. || 17. Mohanavalli B, Dhevahi E, Menon T, Malathi S, Thyagarajan SP . Prevalence of antibodies to hepatitis A and hepatitis E virus in urban school childre