

Medial Open Wedge High Tibial Osteotomy for Osteoarthritis of Knee : Long Term Results in 30 Patients of Smimer, Surat, Gujarat



Medical Science

KEYWORDS : Tibia, Knee, Osteoarthritis, Osteotomy

Dr. Tejas patel

Orthopedics Surgeons, Department of Orthopedics, SMIMER, Surat.

Dr. Pratik Patel

Orthopedics Surgeons, Department of Orthopedics, SMIMER, Surat.

Dr. Sweta patel

anaesthesias, department of anaesthesia, GMCS ,surat

ABSTRACT

Introduction : To retrospectively study the long term results of high tibial valgus osteotomy in the management of primary medial compartment osteoarthritis, with special reference to patient satisfaction and functional assessment, we conducted this study in radiologically established 30 knees of patients with primary osteoarthritis of knee. In these patients with medial compartment disease (varus knee), medial open wedge osteotomy was performed using full thickness iliac crest grafts. These were called for follow-up after 2 years and clinico-radiological assesment was done.

Although many surgeons of the west do not favour this osteotomy, we found the prosedure to be quite acceptable to our patients in whom the primary concern of the cost and squatting habits are well taken care of. The authors feel that this osteotomy is still relevant in the third world. Results do deteriorate with time but most patients consider the surgery satisfactory.

Materials and Methods : The authors retrospectively reviewed 30 knees of patients who underwent medial open wedgeosteotomy for primary medial unicompartmental osteoarthritis of knee between 2012 and 2014. Cases of secondary osteoarthritis were excluded from the study. Weight bearing long film X-rayswere obtained from all the symptomatic patientsand were assessed for TFA (tibia-femoral angle).Functional assessment was carried out at the end of 2 years post-operatively.

Results : Results have shown that medial compartment open wedge high tibial osteotomy is very effective in the management of osteoarthritis of knee as far as pain relief, joint stability, quality of life and patient satisfaction is considered.

Conclusion : The authors conclude that medial opening proximal tibial osteotomy is a physiologicallybetter surgery in medial compartment primary osteoarthritis of knee in early stages. The longtermresults deteriorate with time but are stillgratifying enough for the Indian patients as well asfor those of the third world, who cannot affordthe costlier surgeries and implants and also arereluctant to change their squatting habits and job profile.

Introduction : Osteoarthritis is a multifactorial disease but abnormal stress produced by biochemical is one of the major accelerating factors. Minor degree of varus or valgus deformity of knee alters the load on the tibial and femoral condyles. Main symptoms of osteoarthritis are disabling pain, restriction of movements and deformity. Initial symptomatic treatment consists analgesics (NSAIDS), rest, exercise and even local steroid/ anaesthetic injections. Various surgical produres have been described in literature form time to tome like synovectomy, joint debridement, arthrodesis,patellectomy,patelloplasty and meniscectomy. Recent introduction of hemiarthroplasty, arthroplasty and arthroscopic techniques have proved to be excellent in expert hands.

Tibial osteotomies were introduced in 1950's and have been variedly used and modified since then.¹ These open wedge and closed wedge osteotomies have shown variable results. Most of the work cited in the literature has been that of the closed wedge osteotomies for medial compartment disease. These osteotomies change the weight-bearing axis of the knee.² Venous decongestion has also been cited to be one factor responsible for pain relief apart from axial realignment.³

This retrospective study was performed in all patients of primary osteoarthritis of knee who were operated in Surat Municipal Institute of Medical Sciences, Surat between 2012 and 2014. Records of all these patients were studied and 30 consecutive patients were called for follow-up in last quarter of 2014. The results had been previously assessed on pain and patient satisfaction scale at 2 years follow-up.

Although western literature now advocates unicompartmental hemi-arthroplasty and arthroplasty for even unicompartmental osteoarthritis of knee, the results of this study show that in primary osteoarthritis involving medial compartment, medial open wedgeosteotomy still holds the relevance in the third world countries. This is primarily because patients still can not afford the implant and the need for post surgical avoidance of squat-

ting and load carrying habits is not readily acceptable in initial stages of oXOA knee.

Methodology : The authors retrospectively reviewed 30 knees of patients who underwent medial open wedge osteotomy for primary medial unicompartmental osteoarthritis of knee between 2012 and 2014. Cases of secondary osteoarthritis were excluded from the study. Twenty-two patients were female subjects and Eight patients were male subjects. 44% patients were in the age group of 55 to 65 years and 28% were in the age group of 45 to 55 years. 20% of the total were sedentary workers, the disease involved both knees equally. Duration of pain varied from 1 year to 10 years.

Only those patients who did not have ligamentous injuries, dysfunctional spine, hip or ankle, vascular insufficiencies, flexion deformity of more than 20° and varus deformity of more than 25°, were included in the study. Weight bearing long film X-rays were obtained from all the symptomatic patients and were assessed for TFA (tibia-femoral angle).

Surgery was performed under spinal anesthesia and pneumatic tourniquet. The knee was flexed to 90° to avoid popleteal structures. A vertical skin incision was made from just above joint line and carried downwards to antero-medial aspect of tibia. Joint was not opened. Site of osteotomy was selected 1.5 to 2 cm below the joint line and was confirmed by passing a guide wire under image intensifier or X-ray. A fine osteotomy was performed up to 10 mm of the lateral cortex, taking care to preserve it. Medial ligament was sub-periosteally elevated from its inferio- or insertion if greater degree of varus was to be corrected.

Three full thickness cortico-cancellous grafts were obtained from iliac crest by standard technique.

These were placed in the opened osteotomy (Fig. 1) in progressively decreasing height from posterior to anterior side (Fig.2) to avoid flexion deformity of knee. Stability of osteotomy was as-

sessed and staples were used if instability was detected. All patients were applied POP cast from groin to toe for six weeks.

Isometric quadriceps and straight leg raising physiotherapeutic exercises were encouraged from the day of surgery and continued. Union was evaluated at 6 weeks on removal of cast. Knee flexion exercises were started and partial weight bearing with crutches was allowed at 8 weeks. Full weight bearing was allowed at 12 weeks. Functional assessment was carried out 2 years post-operatively. Results were evaluated as in Table 1.

Results : The knee deformity of these 30 symptomatic radiologically proven cases of medial unicompartamental osteoarthritis ranged from 100 varus to 20 valgus. On an average the deformity was 60 varus pre-operatively. Joint space narrowing of medial compartment as compared to the lateral compartment was present in all cases. 40% of patients had restriction of either flexion (06 cases), extension (4 cases) or both (2 cases) preoperatively.

Postoperatively, none of the patients showed deterioration of pain. However, as the follow-up period increased, the number of patients with pain relief in the range of 75 -100% decreased as shown in Table 2, although, no patient had deterioration in pain beyond that of nonoperated knee.

Range of movements were gained in all patients except two in which superficial infection occurred at 2 weeks. Full range of motion of knee was present in 80% patients (24 cases) at 2 years follow-up, but in 60% at late follow-up. Restriction was present postoperatively in 5 cases, mainly in extension, in those early cases in which grafts were not laid in progressively decreasing height from postero-anteriorly. This restriction in 3 cases (of total 5) did not improve even at late follow-up.

There was no case of non-union in the present series. All osteotomies united in 6 to 9 weeks. There was no change in joint space at 2 years in 90% of cases. Marginal improvement gained at 2 years in 24 cases was also lost at later review. It is however difficult to make comment upon, because lateral compartment involvement along with opposite knee involvement makes it difficult to compare joint space.

Tibio-femoral angle (TFA) was improved in all cases except one in which early weight bearing caused collapse of the achieved correction in the peri-operative period itself. Table 3 shows details at variable period of follow-up. No definite statistically significant relation could be made between relief of pain and TFA at either 2 years or at late follow-up of 6 - 8 years.

Final results according to Table 1 at 2 years showed good results in 24 cases (80%), fair in 4 cases (13%) and poor in 2 case (Fig.3). The poor result was in the patients who had superficial infection because of which range of knee motion deteriorated by more than 20°. However, at 6 to 8 years follow-up, the patients showing good results (Fig. 4), decreased marginally to 20 (66%). Fair results at long-term follow-up were present in 4 cases (13%). Few complications were observed in the form of a) superficial infection in one case, which was managed by oral antibiotics and b) deep infection that resulted in knee stiffness, in another 2 cases. Collapse of graft due to early weight bearing resulted in inability to achieve any correction of TBA in one patient. There was no neuro-vascular deficit in any of the cases postoperatively. Inadvertent complete osteotomy occurred in three patients. We applied staples to stabilize the osteotomy in these cases.

Table:1 Evaluation of Results

Good	Fair	Poor
Complete relief of pain	Partial relief of pain	No relief of pain
Normal union of osteotomy	Normal union of osteotomy	Delayed union
Movements either improved or retained at pre-operative level	Movements decreased by $\geq 20^\circ$ of pre-operative level	Movements decreased by $\geq 20^\circ$
Joint stable	Joint stable	Joint stable
Patient fully satisfied	Patient partially satisfied	Not satisfied

Table : 2 Pain relief at various period of follow-up

Pain relief (Using VAS*)	Number of knees (n=30) at 2 years of follow-up	Number of knees (n=30) at > 2 years of follow-up
75-100%	22	15
50-75%	02	09
Less than 50%	06	06

Table : 3 Tibio-femoral angle at various period of follow-up

Pain relief (Using VAS*)	Number of knees (n=30) at 2 years of follow-up	Number of knees (n=30) at > 2 years of follow-up
75-100%	22	15
50-75%	02	09
Less than 50%	06	06

Discussion : Worldwide, osteoarthritis is the most common articular disease of adults 65 years and older. Osteoarthrosis of knee is the commonest of all symptomatic joint arthroses in Indian subcontinent although its incidence is second to that of distal and proximal interphalangeal joints of the hand worldwide.^{4,5} It has a great disability in old age and is frequently symptomatic. Risk factors of OA include age - the strongest risk factor, obesity, genetics, muscle weakness and joint injury.^{6,7,8}

The management of this “disease of old age” includes physical therapy, weight reduction and if needed drug therapy. Once TFA changes from normal, associated with joint space reduction, no amount of drug or physical therapy has been able to revert these changes. Although a plethora of surgical treatment modalities have been cited in literature, all have pitfalls in one or other forms. Hemi-arthroplasty or arthroplasty have been the main stay of treatment of this disorder in the west in recent past.

Patients of this part of the globe have habit of squatting for toilet purposes and also for normal sitting, which is to be totally avoided after any such endeavor. Also, the cost factors involved are prohibitive for most rural Asian population. Manual labour is one of the commonest sources of income in such population and arthroplasty leads to the need of having to change the work profile of such patients.

Conventry technique of high tibial osteotomy gained some significance before arthroplasties became the preferred modality of management in the west. Difficulty of exposure, involvement of lateral popliteal nerve and sacrifice of upper tibio-fibular joint are some of the cited difficulties which made this surgery unpopular, apart from the reason that most long-term studies show that proximal tibial osteotomies deteriorate with time.⁹

Conventional high tibial osteotomy for osteoarthritis of the medial compartment of the knee with closed-wedge or dome osteotomy (DMO) may produce shortening of the patellar tendon and loss of inclination of the proximal tibial plateau or of the offset of the tibial condyle relative to its bony axis. This can make subsequent total knee arthroplasty technically demanding.¹⁰ The reported incidence of revision for even unicompartamental arthroplasty for failed high tibial osteotomy, in a recently reported series, was nine times higher than that for primary unicompartamental arthroplasty.

This retrospective study was carried out to find the relevance of open wedge tibial osteotomy in patients with primary medial compartmental osteoarthritis of knee. Thirty consecutive operated knees of patients in whom this osteotomy was done from 2012 to 2014 were evaluated at 2 years and then later at 6 to 8 years for overall function and pain according to Table 1.

The general observation made by the authors is that deterioration of TFA occurred at longer follow-up period of 6 to 8 years although all the patients in which this occurred did not deteriorate on satisfaction scale. No statistically significant relation could be made between TFA and pain relief at either 2 years or later, in the current series. It is interesting to note that although deterioration of range of pain relief was present in 06 knees over 6 to 8 years, these all did not deteriorate on the satisfaction scale. This might be a reflection of tendency of "acceptability" in the Asian population.

Good result was seen in 66% cases and 16% cases had fair result at even 6 - 8 years follow-up. This observation is similar to that of Majima et al who have reported that in their series that at the final follow-up, the clinical score had deteriorated relative to the 1 year results but still was significantly better than the preoperative score. Their results indicated that greater the surgical valgus correction, the slower the progression of medial joint arthrosis. Lateral joint arthrosis did not progress more quickly after high tibial osteotomy, even if an over-correction was performed.¹¹

Restriction of movement by more than 20° made to patients fare poorly. These cases had infection after suture removal and much improvement of range of motion could not be gained inspite of dedicated physiotherapy. Restriction of extension occurred in those case where placement of grafts had not been proper. This placement should be from postero-anterior direction so that postoperatively osteotomy does not unite in anterior angulation (Fig 2). An advantage of open wedge osteotomy is that leg length is restored and medial ligament and patellar tendon get tightened and their stabilizing function is restored. There are reports in the recent literature, which confirm that patellar 'lowering' occurs more often with open wedge osteotomy (OWO) than with closed wedge osteotomy (CWO) and the latter also produces a high degree of patellar elevation.¹²

Provided an optimal correction is achieved (3 degrees to 6 degrees hypercorrection in valgus osteotomy, 0 degree in varus osteotomy) and provided a horizontal joint line is restored, HTO performed in good indications (Ahlback grade I or II) may provide good results for at least 10 to 15 years. In a recently reported series, it was noted that osteoarthritis had been arrested after a HTO, in 105 cases (69%) whereas it had deteriorated in 47 cases. The main factors associated with further deterioration were insufficient correction and persistence of joint line obliquity.¹³ We did not, however, observe any such co-relation.

We used iliac crest grafts for opening the medial wedge osteotomy and used plaster of Paris cast for immobilization routinely. Stabilization of osteotomy, when required, was done by using staples. Interposition of the three corticocancellous grafts creates sufficient stability for internal fixation to be unnecessary.¹⁴ Early weight bearing caused collapse of the corrected T-F Angle, resulting in a fair result. There has been an upsurge in the techniques of medial opening HTO recently and the use of external fixators and also bone cement as a substitute of bone has been equally gratifying.^{15,16} Many reports also suggest the ease of early weight bearing and total knee arthroplasty after such procedures.

The inability to explain why 6 of our cases, which had a deterioration in the range of pain scale but were still satisfied, is analogous to a recent comparative study of a group of elderly African Americans and white patients with OA of the knee or hip in the United States. The African Americans were found to be more likely than whites to perceive various traditional and complementary care modalities as efficacious and less likely to perceive joint replacement therapy as efficacious.¹⁷ This reflects that patients have different perception of different treatment modalities. Asian patients (Indians) accepted the procedure well and were found satisfied even after 6 to 8 years although the varus had again increased marginally and so had pain.

Leutloff has reported poor results of patients with varus of knee showing radiological signs of osteoarthritis in contrast to those without radiological evidence.¹⁸ This might be a manifestation of irreparable cartilage damage. Others have reported that the outcome with medial open wedge osteotomy is comparable with or better than that of other techniques for osteotomy and that subsequent knee replacement, in cases requiring conversion, is straight forward.¹⁹ Even pattern of scintigraphic uptake is reportedly altered with HTO and that this has paralleled the improvement in Symptoms.²⁰

Conclusions & Recommendations : The authors conclude that medial opening proximal tibial osteotomy is a physiologically better surgery in medial compartment primary osteoarthritis of knee in early stages. The longterm results deteriorate with time but are still gratifying enough for the Indian patients as well as for those of the third world, who can not afford the costlier surgeries and implants and also are reluctant to change their squatting habits and job profile. Its possible role in high degree of varus (more than 20°), secondary OA as well as in lateral compartment disease is out of scope of this study. Complications are minimal and avoidable. Use of external fixators for callosities might be a good alternative but tropical climate deters many a surgeon because of significant risk of pin site infection that can lead to disaster for current and future knee endeavors.

REFERENCE

- Jackson JP, Waught W, Green JP. High tibial osteotomy for osteoarthritis of the knee. *J Bone Joint Surg* 1969; 51B: 88-94. | | 2. Bauer GCH, Insall J, Koshino T. Tibial osteotomy in gonarthrosis. *J Bone Joint Surg* 1969; 51A: | 1545-62. | | 3. Helal B. The pain in primary osteoarthritis of knee. Its causes and treatment by osteotomy. *PostgradMedJ* 1965; 41: 172-81. | | 4. Lawrence RC, Hochberg MC, Kelsey JL et al. Estimates of the prevalence of selected arthritic and musculoskeletal diseases in the United States. *J Rheumatol* 1989; 16: 427-41. | | 5. Oliveria SA, Felson DT, Reed JI et al. Incidence of symptomatic hand, hip and knee osteoarthritis among patients in a health maintenance organization. *Arthritis Rheum* 1995; 38: 1134-41. | | 6. Felson D, Anderson J, Naimark A et al. Obesity and knee osteoarthritis: The Framingham study. *Ann Intern Med* 1988; 109: 18-24 | | 7. Felson D, Zhang Y, Hannan Met al. Risk factors for incident radiographic knee osteoarthritis in the elderly. *Arthritis Rheum* 1997; 40: 728-33. | | 8. Slemenda C, Brandt K, Heilman D et al. Quadriceps weakness and osteoarthritis of the knee. *Ann Intern Med* 1997; 127: 97-104. | | 9. Sangwan SS, Siwach RC, Singh Z, Duhan S. Unicompartmental osteoarthritis of the knee: an innovative osteotomy. *Int Orthop* 2000; 24: 148-50. | | 10. Nakamura E, Mizuta H, Kudo S, Takagi K, Sakamoto K. Open-wedge osteotomy of the proximal tibia hemicallotaxis. *J Bone Joint Surg* 2001; 83B: 1111-5. | | 11. Majima T, Yasuda K, Katsuragi R, Kaneda K. Progression of joint arthrosis 10 to 15 years after high tibial osteotomy. *Clin Orthop* 2000; 381: 177-84. | | 12. Tigani D, Ferrari D, Trentani P, Barbanti-Brodano G, Trentani F. Patellar height after high tibial osteotomy. *Int Orthop* 2001; 24: 331-4. | | 13. Antonescu DN. Is knee osteotomy still indicated in knee osteoarthritis? *Acta Orthop Belg* 2000; 66: 421-32. | | 14. Marti RK, Verhagen RAW, Kerckhoffs GMMJ, Moojen TM. Proximal tibial varus osteotomy. *J Bone Joint Surg* 2001; 83A: 164-70. | | 15. Hnigou P, Ma W. Open wedge tibial osteotomy with acrylic bone cement as bone substitute. *Knee* 2001; 8: 103-10. | | 16. Klinger HM, Lorenz F, Harer T. Open wedge tibial osteotomy by hemicallotaxis for medial compartment osteoarthritis. *Arch Orthop Trauma Surg* 2001; 121: 245-7. | | 17. Ibrahim SA, Siminoff LA, Burant CJ, Kwok CK. Variation in perceptions of treatment and self-care practices in elderly with osteoarthritis: a comparison between African American and white patients. *Arthritis Rheum* 2001; 45: 340-5. | | 18. Leutloff D, Tobian F, Perka C. High tibial osteotomy for valgus and varus deformities of the knee. *Int Orthop* 2001; 25: 93-6. | | 19. Weale AE, Lee AS, MacEachern AG. High tibial osteotomy using a dynamic axial external fixator. *Clin Orthop* 2001; 382: 154-67. | | 20. Bruce W, Macdessi S, Van der Wall H. Reversal of medial compartment osteoarthritic uptake pattern after high tibial osteotomy. *Clin Nucl Med* 2001; 26: 916-8. |