

Study of Surgical Site Infections and antibiotic susceptibility pattern of isolates at a tertiary care hospital in Amritsar



Microbiology

KEYWORDS : Surgical site infections (SSI's), surgeries

Dr.Suchira Chillana

Junior resident , Department of Microbiology, Government Medical College, Amritsar. Punjab

Dr. Bimla Devi

Professor, Department of Microbiology, Government Medical College Amritsar. Punjab

Dr.Pushpa Devi

Professor & Head, Department of Microbiology, Government Medical College Amritsar. Punjab

Dr.Shailpreet Kaur

Assistant Professor, Department of Microbiology, Government Medical college, Amritsar. Punjab

Dr.Kapil Alias Mohit Chilana

Junior Resident, Krishna Institute of Medical Sciences, Secunderabad, Andhra Pradesh

ABSTRACT

Surgical site infections are the third most frequently reported nosocomial infections accounting for 14-16% of all the infections in hospitalised patients. Methodology: Two hundred swabs/pus specimens from the patients developing surgical site infections at Guru Nanak Dev Hospital, GMC, Amritsar were processed in the Department of microbiology during Jan-Nov2015, by standard methods and antibiotic susceptibility testing of all the isolates was done by using Kirby Bauer disc diffusion technique. Results: Of the 153 organisms isolated, the most common was Staphylococcus aureus (47.7%), followed by Pseudomonas aeruginosa (16.99%), Escherichia coli (14.37%), Klebsiella pneumoniae (9.15%), miscellaneous gram negative rods (9.14%) and Streptococcus pyogenes (2.61%). About 30% of the Staphylococcus aureus isolates were found to be methicillin resistant. Conclusion: We should clearly understand and identify this as a problem and devise a system to track, analyze and monitor these.

Introduction:

Surgical site infections (SSI) are the third most frequently reported nosocomial infections accounting for 14-16% of all the infections in hospitalised patients. Among surgical patients SSI are the most common nosocomial infections¹. These remain a complication of surgical procedures resulting in increased morbidity, mortality and cost². The risk of developing a surgical site infection depends upon the balance between factors determining the number of bacteria contaminating the site and the factors determining the resistance of the site against infection^{3,4}.

One of the major problems faced by the surgeons these days is to deal with the post surgical infections, as most of these are being caused by multiple resistant bacteria. Gram positive cocci and Gram negative bacilli are being implicated in most of such cases^{5,6}. The problem of infected surgical sites can only be tackled properly if all these are examined bacteriologically and feedback given to the surgeons well in time, so that they can treat these with appropriate antibiotics⁷. Not only this but, the microbiologist should provide them the guidance regarding proper use of prophylactic antibiotics. But, according to the cure⁷, prevention of surgical site infections by adopting basic principles of asepsis is the key to the solution of this problem⁸.

Material and Methods:

Two hundred wound swabs/pus specimens collected from patients developing surgical site infections during a period from January 2015 to November 2015, were included in the study. Most of our patients were young males (n=80). Rest were females (n=38) and children (n=11). The age range was between 4-65 years and had undergone different kinds of surgery including general surgery (n70), gynecological/obstetric surgery (n48) and orthopedic surgery (n11). Pus swabs/specimens were collected from infected surgical sites by standard technique using commercially available sterile stick swabs. The specimens were immediately transported to the Department of Microbiology, GMC, Amritsar for bacteriological study. All the specimens were inoculated onto blood and MacConkey's agar within two hours

of collection. The agar plates were incubated at 37°C aerobically and were examined for the presence of any growth after 24 hours. Those plates showing no growth were incubated for another 24 hours. The isolates were identified by colonial morphology, Gram's stain and conventional biochemical tests. Antibiotic susceptibility pattern of the isolates was studied using Kirby Bauer method⁹. Mueller Hinton agar (Difco) was used for antibiotic susceptibility testing. Staphylococcus aureus ATCC 25932, Escherichia coli ATCC 25922 and Pseudomonas aeruginosa ATCC 27853 were included as control strains.

Results :

One hundred and fifty three organisms were isolated from the 200 specimens processed. One hundred and five specimens yielded growth of single organism while two isolates were present in rest of the twenty-four cases. The most common pathogen isolate was, Staphylococcus aureus (47.71%), followed by Pseudomonas aeruginosa (16.99%), Escherichia coli (14.37%), Klebsiella pneumoniae (9.15%), Streptococcus pyogenes (2.61%) and miscellaneous gram negative rods (9.14%) including Acinetobacter baumannii, Proteus mirabilis and Citrobacter diversus (Table 1)

Table 1. Pathogens isolated from Surgical site infections

Organism	No.	Percent %
Staphylococcus aureus	73	47.71
Pseudomonas aeruginosa	26	16.99
Escherichia coli	22	14.37
Klebsiella pneumonia	14	9.15
Acinetobacter baumannii	6	3.92
Proteus mirabilis	4	2.61
Citrobacter koseri	4	2.61
Streptococcus pyogenes	4	2.61
Total	153	

In case of Staphylococcus aureus 30% of the isolates were found resistant to methicillin. Antibiotic susceptibility pattern of gram-

positive cocci (Staphylococcus aureus and Streptococcus pyogenes) to other antibiotics are shown in (Table 2)

Table 2. Antibiotic susceptibility pattern(Percent sensitive) of Gram positive cocci

Antibiotic	Staphylococcus aureus %	Streptococcus pyogenes %
Amikacin	86.3	-
Gentamycin	45.2	-
Ciprofloxacin	65.75	-
Cotrimazole	49.31	0
Methicillin	30.1	-
Vancomycin	100	100
Cephalexin	54.79	100

In case of Pseudomonas aeruginosa only 8 isolates (30.7%) were gentamicin sensitive. Quite a few strains were also found resistant to piperacillin-tazobactam, ciprofloxacin and ceftazidime. Antibiotic susceptibility pattern of all the Gram negative rods (GNRs) studied is shown in (Table 3).

Table 3. Antibiotic susceptibility pattern (Percent sensitive) of Gram negative rods (GNRs)

Antibiotic	Pseudomonas aeruginosa	Escherichia coli	Klebsiella pneumoniae	Miscellaneous GNRs
Amikacin	50	54.5	42.8	50
Gentamycin	30.7	36.3	50	57.1
Ciprofloxacin	53.8	40.9	50	28.6
Ceftazidime	50	50	50	71.42
Ceftriaxone	69.2	40.9	64.2	64.2
Sulbactam-ceftriaxone	73	63.6	42.8	71.42
Piperacillin – tazobactam	61.5	72.7	71.4	64.3
Imipenem	84.6	72.7	57.1	57.1

Discussion:

Surgical site infection can be defined as the presence of pus alongwith signs of inflammation in the surgical wound margins. Predisposing underlying conditions for surgical site infections include immunosuppression, irradiation, steroid administration, diabetes mellitus and malnutrition¹⁰. The risk of infection after surgery depends upon the factors including the type and length of surgical procedure; age, underlying conditions and previous history of the patient: skill of the surgeon; diligence with which infection control procedures are applied and the type and timing of pre-operative antibiotic prophylaxis¹¹ Most of the patients included in the study were young males with minimal predisposing factors except that six of these were diabetic. So the factors most probably operative in causing infections in our patients were related to the surgical team or surgical environment. Staphylococcus aureus is considered to be the leading pathogen in such post-surgical wound infections followed by the members of the enterobacteriaceae¹². But in our study the Pseudomonas aeruginosa was the second commonest isolate after Staphylococcus aureus. Otokunefor TV and Datubo-Brown DD also have found similar isolates in most of the patients included in their study¹³.

Most of our isolates were found resistant to the commonly used antibiotics. This is a matter of great concern because treatment of such infections warrants newer and costly antibiotics. The incidence of methicillin resistant Staphylococcus aureus (MRSA) in our study in about 30%. MRSA infections cannot be treated by beta lactarnase resistant penicillins and not even by the cepha-

losporins¹⁴. Treatment of these infections is possible either by the Fluroquinolones (if the isolate is found sensitive) or by the vancomycin only¹⁵. More than 50% were sensitive to ceftriaxone and ciprofloxacin which are thus the minimal choice to treat Pseudomonas aeruginosa infections. Maximum sensitivity of Pseudomonas was seen to Imepenem,sulbactam-ceftriaxone and piperacillin tazobactam.. But an empirical treatment to be really effective against such isolates will have to include either amikacin or one of the carbapenems alone or in combination. Even the Escherichia coli, Klebsiella pneumonia and the other Gram negative isolates in our study are showing fairly high antibiotic resistance. In view of such highly resistant organisms causing wound infections in our hospital. it will become very difficult to treat these cases. So the only hope lies with prevention of such surgical site infections. To achieve this goal we will have to return to the preventive measures including fundamental principles of asepsis. Individual patient risk factors must be identified and modified whenever possible. In addition to the skin asepsis and peri-operative prophylactic antibiotics, care and attention to the theatre environment is also very important¹⁶. Last but not the least, surgical expertise and theatre discipline are the essential components against surgical site infections.

Conclusion

We should clearly understand and identify the SSI as a problem and devise a system to track, analyze and monitor these. Hospital infection control committees should meet regularly and make recommendations at all levels for prevention of such incidents. Otherwise it will be impossible to overcome the serious issues of economic loss and high hospital morbidity and mortality caused by SSI.

References

1. Mangram AJ, Horan TC, Pearson ML, et al. "Guideline for Prevention of Surgical Site Infection 1999" issued by (enters for Disease Control and Prevention through its 1 2—member Hospital Infection Control Practices Advisory Committee. Am. J. Infect. Control, 1999;27:27-30.
2. Bay-Nielsen M. Risk factors for surgical wound infection Ugeskr. Laeger. 1996;158:5749-53
3. Wenzel RP. Prevention and treatment of hospital acquired infection. In: Wyn-gaarden, Smith Ed. Cecil textbook of medicine. 18th ed, Philadelphia, WB Saunders Co., 988, pp.
4. Noer HH Interest and possibilities of post-operative registration of wound infections in Danish orthopaedic department 'A Study Survey. Int. J. Clin.. Monit. Comput., 1990;1:21-26.
5. Bergogne BE, Deere D, Joly ML. Opportunistic nosocomial multiply resistant bacterial nfecti ons—their treatment and prevention. J-Antimicrob. Chemother., 1993;32:Suppl A:39-47.
6. Olson MM, Lee JT. Continuous 10 year wound infection surveillance. Arch. Surg., 1990;125:794-803.
7. Gorback SL, Cartless JG, Nichols L. Epidemiology and prevention of surgical infections. 1st ed, Boston, Little Brown and Co., 1984, pp.
8. Cowan ST. Cowan and Steel's manual for the identification of Medical Bacteria. 2nd Edition, Cambridge University, Press, 1974, pp.
9. Baur AW, Kirby WM, Shris JC, et al. Antibiotic susceptibility testing by a standardized single disk method. Am. J. Clin. Pathol., 1966;45:493-96.
10. Speller OCE. Hospital associated infections. In: Topley and Wilson's Principles of Bacteriology, Virology and Immunology. London, Edward Arnold, 1990;3: 142-67.
11. Nicholas RL. Post-operative infections in the age of drug resistant gram positive bacteria. Am. J. Med., 1998;104(5A):11S-16S.
12. Kotisso B, Aseffa A. Surgical wound infection in a teaching hospital in Ethiopia. East Afr. Med. J., 1998;75:402-405.
13. Otokunefor TV and Datubo-Brown DD. Bacteriology of wound infections in a surgical ward of a teaching hospital. West Afr. J. Med., 1990;9:285-90.
14. Kernodle DS. Failure of cephalosporins to prevent Staphylococcus aureus surgical wound infections. JAMA, 1990;263:961-66.
15. Sta AM, Long MN, Belcher B. Higher overall nosocomial infection rate because of methicillin resistant Staphylococcus aureus. Am. J. Infect. Control, 1993;2 1:70-74.
16. Emmerson M. A microbiologist's view of factors contributing to infection. New Horiz, 1998;6(2 Suppl): S3-10.