

Serum Albumin and Nutritional Status of Patients with Chronic Kidney Disease: A Hospital-Based Study



Medical Science

KEYWORDS :nutritional status, serum albumin, chronic kidney disease

* **FABUSORO Olufemi K.**

Lecturer, Department of Human Nutrition, College of Medicine, University of Ibadan, Ibadan, Nigeria., Registered Dietitian, United Kingdom (Registration no. DT27561), * Corresponding author

ABSTRACT

• *Patients with chronic kidney disease (CKD) often suffer from malnutrition and protein-energy wasting (PEW) resulting in poor nutritional status which is a powerful predictor of mortality. There is a dearth of information on the nutritional status of patients with chronic kidney disease in Ibadan, Nigeria.*

• *This study is a cross sectional study. A total of 32 consenting in-patients were assessed. The mean value of serum albumin and creatinine were 2.88±0.5 g/dL and 11.93±7.78 mg/dL respectively. The BMI of the male and female patients in this study were 22.29± kg/m² and 23.41 kg/m² respectively while Mid-Upper Arm Fat Area (AFA) for the male and the female patients were 7.04±2.12 cm² and 11.78±10.02 cm² respectively. The protein and energy intake for the patients were also low compared to standard values.*

• *It is important to maintain nutritional status in CKD population.*

INTRODUCTION

Malnutrition is recognized to be a serious and common complication of Chronic Kidney Disease (CKD) and is associated with increased morbidity and mortality in children¹ and adults². Approximately 80 percent of deaths due to chronic disease occur in low and middle income countries³. Hospital based data in Nigeria have reported prevalence rates expressed as ratios of hospital admissions of between 1.6 and 8%⁴.

Biochemical analyses are routinely used to assess and monitor nutritional status in patients with CKD. However, none of the currently favoured biochemical nutritional markers have been demonstrated to accurately reflect nutritional status in CKD^{5,6,7}. Nevertheless, serum albumin is still being widely used for research purposes and, in the clinical setting, as a biomarker of nutritional status⁸.

There is paucity of data on the serum albumin, other biochemical markers of nutritional status and anthropometrics of patients with CKD in Ibadan, Nigeria. This study therefore seeks to bridge the gap.

MATERIALS AND METHODS.

Study design: This study was a cross-sectional study.

Study location: The study was carried out at the renal unit of the University College Hospital (UCH), Ibadan, Nigeria.

Study population and sample size:

A total of thirty-two (32) in-patients comprising of 21 males and 11 females were purposively selected and assessed. Chronic Kidney Disease (CKD) is defined as kidney damage or glomerular filtration rate (GFR) <60 mL/min/1.73 m² for 3 months or more, irrespective of cause⁹.

Inclusion criteria:

Male and female adult patients who had been confirmed by nephrologists to have chronic kidney disease with GFR less than 60 ml/min/1.73 m² and those with end stage renal disease that were been treated with haemodialysis for a minimum of 2 months. The patients must have given their consent to participate in the study.

Exclusion criteria:

Any patient with CKD aged <18 years or patient with other serious medical condition such as cancer, HIV, hepatitis and those who refused to give their consent, were excluded from the study.

Data collection procedure:

A standardized semi-structured interviewer administered questionnaire were used to obtain information on the socio-demographic characteristics of the patients. Weight and height were assessed to determine body mass index (BMI) which was categorised as underweight (<18.5kg/m²), normal weight (18.5-24.9kg/m²), overweight (25-29.9kg/m²) and obese (≥30.0kg/m²). Calculations of mid-upper arm fat area (AFA) and arm muscle area (AMA) were based on measurements of mid-upper arm circumference (MAC) and triceps skinfold (TSF). AFA and AMA data were then compared with the data from the National Health and Nutrition Examination Survey (NHANES I and II)¹⁰.

$$\text{Formular 1: } AFA \text{ (cm}^2\text{)} = \frac{MAC \times TSF}{2} - \frac{\pi \times (TSF)^2}{4}$$

$$\text{Formular 2: } AMA \text{ (cm}^2\text{)} = \frac{[MAC - (\pi \times TSF)]^2}{4 \pi}$$

Blood was drawn to about ¾ of the capillary tube. Laboratory analysis was done to determine serum albumin, serum creatinine, urea, Pack cell Volume (PCV), sodium and potassium levels.

Dietary intake was determined by administering the foods served to the patients by the dietetics department, University College Hospital, Ibadan, Nigeria using direct weighing method for 3 days. The study days were 2 week days and one weekend. The protein and energy intake was then calculated using Total Dietary Assessment software.

Data Analysis: Quantitative data was expressed as mean ± standard deviation. Statistical significance was taken as p<0.05.

RESULTS

Table 1: Dietary intake of the patients

	Day	Range	Mean±SD
Body weight (Kg)		34	62.19±8.88
Protein (g/kg body wt)	Day 1	1.70	0.73±0.33
	" 2	1	0.64±0.35
	" 3	1.2	0.70±0.47
Mean protein		1.4	0.69±0.31
Energy (g/kg body wt)	Day 1	39.5	22.00±8.7
	" 2	42.2	20.52±10.38
	" 3	38.0	18.16±11.74

Mean Energy (g/kg body wt)	30.67	20.21±8.41
Mean Energy	1473.86	1275.26±500.4

Table 2: Serum Albumin and other biochemical indices of the CKD patients.

	Range	Mean±SD
Albumin (g/dl)	2.1	2.88±0.50
Urea (mg/dl)	326	151.25±78.02
Creatinine (mg/dl)	28.0	11.93±7.78
Potassium (K ⁺) (mmol/l)	5.7	4.07±1.30
Sodium (Na ⁺) (mmol/l)	32	134.31±7.74
PCV (%)	27	22.66±5.77

Table 3: Anthropometric status of the patients

	Range	Mean±SD
Male		
Age (years)	53	39.57±12.91
BMI (Kg/m ²)	11.19	22.29±4.59
AFA (cm ²)	7.02	7.04±2.12
AMA (cm ²)	40.15	43.02±10.65
Female		
Age (years)	36	36.27±9.95
BMI (Kg/m ²)	20	23.41±6.39
AFA (cm ²)	32.72	11.78±10.02
AMA (cm ²)	25.19	32.70±8.20

Discussion

The kidney Disease Outcome Quality Initiative (K/DOQI)⁹ Clinical Guidelines for nutrition and chronic kidney disease have recommended that BMI of hemodialysis patients be maintained in the upper 50th percentile, which would correspond to a BMI of at least 23.6 for males and 24.3 for females. The mean BMI of the males and females in this study were 22.29± kg/m² and 23.41 kg/m² respectively. This shows that the patients are at increased risk of mortality. In addition, Caravaca *et al.*¹¹ showed that patients with CKD not yet on dialysis within the lowest and the highest quartiles of BMI had higher mortality than the rest of patients, but when patients without comorbidity were studied apart, only those with obesity showed worse survival, suggesting that obesity had a noteworthy impact on mortality in these patients.

At ESRD stage, the effect of overweight (BMI: 25–30 kg/m²) or obesity (BMI: >30 kg/m²) has been repeatedly associated with improved survival whereas, in contrast, a BMI <19 kg/m² was associated with increased mortality^{12,13,14}.

The result also shows that the Mid-Upper Arm Fat Area (AFA) for the male and the female patients were 7.04±2.12 cm² and 11.78±10.02 cm² respectively. When this was compared with their mean age of 39.57±12.91 years and 36.27±9.95 years respectively, it suggests that they were severely depleted when compared with the reference data compiled from National Health and Nutrition Survey (NHANES)¹⁰.

On the contrary, the values of the Mid-Upper Arm Muscle Area (AMA) for the male and the female patients were 43.02±10.65 cm² and 32.70±8.2 cm² respectively which fell between 10th and 15th percentile for the males and between 50th and 75th percentile for the females. This however shows no depletion when the same parameters (i.e Mid-upper arm circumference and triceps skinfold) were used for both the AFA and the AMA.

The protein and energy intake for the patients were 0.69 g/kg/ body wt and 1275.26 kcal respectively. Other stud-

ies have also shown that patients with CKD consume an energy and protein-deficient diet¹⁵. This can be as a result of clinical features such as nausea and anorexia associated with CKD.

The mean value of serum albumin and Creatinine were 2.88±0.5 g/dL and 11.93±7.78 mg/dL respectively. These figures are significantly poor and this can be as a result of patients reporting late to the hospital¹⁶. Present analysis also supported the possibility that animal protein intake was related with serum albumin levels¹⁷. Hypoalbuminemia does not normally occur in otherwise healthy but malnourished individuals until near terminal starvation¹⁸.

Conclusion

In patients with CKD and ESRD, in whom many catabolic signals dominate, it is critical to maintain a dietary protein and energy intake relative to needs. Treatment of concurrent conditions that contribute to catabolism, such as metabolic acidosis, insulin resistance, and systemic inflammation, is of paramount importance for the prevention of protein-energy wasting.

References

1. Furth, S.L., Stablein, D., Fine, R.N., Powe, N.R., Fivush, B.A. (2002). Adverse clinical outcomes associated with short stature at dialysis initiation: a report of the North American Pediatric Renal Transplant Cooperative Study. *Pediatrics*.109, 909–13.
2. Leavey, S.F., Strawderman, R.L., Jones, C.A., Port, F.K., Held, P.J. (1998). Simple nutritional indicators as independent predictors of mortality in hemodialysis patients. *Am J Kidney Dis*. 31, 997–1066.
3. World Health Organization. (2005). Preventing chronic disease: a vital investment. Available from, URL:http://www.who.int/chp/chronic_disease_report/full_report.pdf.
4. Ogun, S.A., Adelowo, O.O., Familoni, O.B., Jaiyesimi, A.E.A., Fakoya, E.A.O. (2000). Pattern and outcome of medical admissions at the Ogun State University Teaching Hospital Sagamu - a three years review. *WJMJ*. 19(4), 304–308.
5. de Mutsert, R., Grootendorst, D.C., Indemans, F., Boeschoten, E.W., Krediet, R.T., Dekker, F.W. (2009). Netherlands Cooperative Study on the Adequacy of Dialysis-II Study Group: Association between serum albumin and mortality in dialysis patients is partly explained by inflammation, and not by malnutrition. *J Ren Nutr* 19, 127–135.
6. Stenvinkel, P., Barany, P., Chung, S.H., Lindholm, B., Heimbürger, O. (2002). A comparative analysis of nutritional parameters as predictors of outcome in male and female ESRD patients. *Nephrol Dial Transplant* 17, 1266–1274.
7. Ikizler, T.A., Wingard, R.L., Harvell, J., Shyr, Y., Hakim, R.M. (1999). Association of morbidity with markers of nutrition and inflammation in chronic hemodialysis patients: A prospective study. *Kidney Int* 55, 1945–1951.
8. Friedman, A.N., Fadem, S.Z. (2010). Reassessment of albumin as a nutritional marker in kidney disease. *J Am Soc Nephrol* 21, 223–230.
9. National Kidney Foundation. K/DOQI. (2002). Clinical practice guidelines for chronic kidney disease: evaluation, classification, and stratification. *Am J Kidney Dis* 392 suppl 1:S1–266.
10. Frisancho, A.R. (1990). Anthropometric Standards for the Assessment of Growth and Nutritional Status. Ann Arbor, The University of Michigan Press.
11. Caravaca, F., Martín, M.V., Barroso, S., et al. (2004). Obesity and mortality in advanced chronic renal failure patients. *Nefrologia*. 24, 453–462.
12. Kalantar-Zadeh, K., Kopple, J.D. (2006). Obesity paradox in patients on maintenance dialysis. *Contrib Nephrol*. 151, 57–69.
13. Kalantar-Zadeh, K., Kuwae, N., Wu, D.Y., et al. (2006). Associations of body fat and its changes over time with quality of life and prospective mortality in hemodialysis patients. *Am J Clin Nutr*. 83:202–210.
14. Kalantar-Zadeh, K., Koopke, J.D., Kilpatrick, R.D., et al. (2005). Association of morbid obesity and weight change over time with cardiovascular survival in hemodialysis populations. *Am J Kidney Dis*. 46:489–500.
15. Norman, L.J., Coleman, J.E., Macdonald, I.A., Tomsett, A.M., Watson,

- A.R. (2000). Nutrition and growth in relation to severity of renal disease in children. *Pediatric Nephrology*. 15, 259–65.
16. Alebiosu, C.O., Ayodele, O.O., Adigun, A., Ina, O.A. (2006). Chronic renal failure at the Olabisi Onabanjo University teaching hospital, Sagamu, Nigeria. *African Health Sciences*. 6(3), 132-138.
17. Watanabe, M., Higashiyama, A., Kokubo, Y., Ono, Y., Okayama, A., Okamura, T; NIPPON DATA80/90 Research Group. (2010). Protein intakes and serum albumin levels in a Japanese general population: NIPPON DATA90. *J Epidemiol*. 20 Suppl 3:S531-6.
18. Rigaud, D., Hassid, J., Meulemans, A., *et al* (2000): A paradoxical increase in resting energy expenditure in malnourished patients near death: The king penguin syndrome. *Am J Clin Nutr* 72:355–360.