


Phytochemicals and Elemental Analysis of *Evolvulus alsinoides* Leaf Extract



Biochemistry

KEYWORDS : *Evolvulus alsinoides*, Phytochemical, Inorganic elements

T.K. Sujayil

Research Scholar, PG and Research Department of Biochemistry, Marudupandiyar College, Tamil Nadu, S. India

T.S. Dhanaraj

PG and Research Department of Biochemistry, Marudupandiyar College, Tamil Nadu, S. India

ABSTRACT

*In the present study to investigate phytochemicals and inorganic elements in *Evolvulus alsinoides*. The results of this study clearly indicate that the preliminary phytochemical analysis of *Evolvulus alsinoides* revealed presence of flavonoids, polyphenol, steroids, tannin, saponins, glycosides, anthroquinones, polyphenol, amino acid, alkaloids, carbohydrate, protein while steroids and Phlobatannins were absent. Quantitative analysis revealed that *Evolvulus alsinoides* rich amount of Total Phenols (324.0 mg/gm), Alkaloids (35.0 mg/gm), Tannin (98.4 mg/gm), Saponin (121.0 mg/gm), Flavonoids 165.0mg/gm) were presented. The vitamin analysis of *Evolvulus alsinoides* showed that the presence of Vitamin E, C and D while vitamin A was absent. The inorganic elements of *Evolvulus alsinoides* showed that the presence of calcium, sodium, potassium, sulphate, phosphate, chloride, Nitrate, magnesium and iron while carbonate was absent. The results of the present study concluded that *Evolvulus alsinoides* may be a good source of phytochemicals, vitamins and minerals. Supplementation of this *Evolvulus alsinoides* leaves may be useful for human health associated emerging diseases such as cardiovascular diseases, diabetes, hypertension and cancer.*

INTRODUCTION

Phytochemicals are a field of increasing attention, both in science and in commerce. As is now generally recognized, many plant compounds and pigments have effects on animals and human beings. There is a great effort now to study and understand at a fundamental level and significant health effects of these compounds. This field is maturing and the health effects of these compounds are now getting the careful study they warrant at both a chemical and a molecular biological level. Identifying bioactive compounds and establishing their health effects are active areas of scientific inquiry. Phytochemicals are biologically active, naturally occurring chemical compounds found in plants, which provide health benefits for humans [1]. They protect plants from disease and damage and contribute to the plant's color, aroma and flavor. In general, the plant chemicals that protect plant cells from environmental hazards such as pollution, stress, drought, UV exposure and pathogenic attack are called as phytochemicals (2,3). Phytochemicals are non-nutritive plant chemicals that have protective or disease preventive properties. Plant produces these chemicals to protect itself but recent research demonstrates that many phytochemicals can protect humans against diseases. There are many phytochemicals fruits and herbs and each works differently (4). The plant kingdom is a rich source of potential drugs. In India, medicinal plants are widely used by all sections of the population, either directly in different indigenous systems of medicine or indirectly in the pharmaceutical preparations of modern medicine. Research on natural resources has been encouraged by the World Health Organization since 1978. Most of the medicinal plants contain a number of chemical constituents such as flavonoids, alkaloids, tannins, saponins, steroids, terpenoids, rotenoids etc. The phytochemical screening of the plants is a preliminary for verification and then these plants may be utilized as new sources of herbal drugs (5).

The mineral elements are separate entities from the other essential nutrients like (proteins, fats, carbohydrates, and vitamins. Animal husbandry had demonstrated the need for minerals in the diet. In this century, biological assay methods clarified the significance and importance of mineral elements for human and animal nutrition and modern analytical techniques led to the detection of trace elements as essential nutrients and this is still an active area of current research. Micronutrient deficiencies are a major public

health problem in many developing countries, with infants and pregnant women especially at risk (6). Infants deserve extra concern because they need adequate micronutrients to maintain normal growth and development (7). The micronutrient deficiencies which are of greatest public health significance are iron deficiency, causing varying degrees of impairment in cognitive performance, lowered work capacity, lowered immunity to infections, pregnancy complications e.g. babies with low birth weight, poor learning capacity and reduced psychomotor skills (6).

Recently, it is clearly known that they have roles in the protection of human health, when their dietary intake is significant. More than 4,000 phytochemicals have been cataloged and are classified by protective function, physical characteristics and chemical characteristics and About 150 phytochemicals have been studied in detail. Since there is no report on the phytoconstituents of methanolic fraction of *Evolvulus alsinoides* leaves extract it was chosen as the subject of this study. Keeping in view, in the preset study to investigate the phytochemical and inorganic elements in *Evolvulus alsinoides* (Tamil: Aannaikatralai).

MATERIALS AND METHODS

Plant materials:

The leaf of *Evolvulus alsinoides* was collected in January 2015 from Tamil University, Thanjavur District, Tamil Nadu, India from a single herb. The leaf was identified and authenticated by Dr. S. John Britto, The Director, the Rabiant Herbarium and centre for molecular systematics, St. Joseph's college Trichy-Tamil Nadu, India. A Voucher specimen has been deposited at the Rabnat Herbarium, St. Josephs College, Thiruchirappalli, Tamil nadu, India.

Preparation of alcoholic extract

The leaf of *Evolvulus alsinoides* was first washed well and dust was removed from the plant. Leaf was washed several times with distilled water to remove the traces of impurities from the plant. The leaf plant was dried at room temperature and coarsely powdered. The powder was extracted with 70% methanol for 24 hours. A semi solid extract was obtained after complete elimination of alcohol under reduced pressure. The extract was stored in refrigerator until used.

Preliminary phytochemicals screening

Chemical tests were carried out on the alcoholic extract using standard procedures to identify the preliminary phytochemical screening following the methodology of Harborne (8), Trease and Evans (9) and Sofowara (10).

Quantitative analysis of phytochemicals

Determination of total phenols (11). Tannin was determined by the method of Van-Burden and Robinson (12). Saponin was determined by the method of Obdoni and Ochuko (13). Flavonoid was determined by the method of Bohm and Kocipai-Abyazan (14).

Qualitative analysis of Vitamins and inorganic elements

Qualitative analysis of Vitamin- A, C and Vitamin -D carried out by the method of Pearson (15). Ash of drug material (500 mg) was prepared and treated with HNO₃ and HCl (3:1 v/v) for 1hour. After the filtration, the filtrate was used to perform the following tests Khandelwal (16).

RESULTS AND DISCUSSION

Phyto is the Greek word for plant. There are many families of phytochemicals and they help the human body in the variety of ways. Phytochemicals may protect human from a host of diseases. Phytochemicals are known as secondary plant metabolites and have biological properties such as antioxidant activity, antimicrobial effect, modulation of detoxification enzymes, stimulation of the immune system, decrease of platelet aggregation and modulation of hormone metabolism and anticancer property. There are more than thousand known and many unknown phytochemicals. It is well-known that plants produce these chemicals to protect themselves, but recent researches demonstrate that many phytochemicals can also protect human against diseases (17).

An assessment of the previous trends and impact of research into the phytochemistry on medicinal plants of the world is quite desirable before considering recent trends. After centuries of empirical use of herbal preparation, the first isolation of active principles alkaloids such as morphine, strychnine, quinine etc. in the early 19th century marked a new era in the use of medicinal plants and the beginning of modern medicinal plants research. Emphasis shifted away from plant derived drugs with the tremendous development of synthetic pharmaceutical chemistry and microbial fermentation after 1945. Plant metabolites were mainly investigated from a phytochemical and chemotaxonomic viewpoint during this period. Over the last decade, however, interest in drugs of plant and probably animal origin has grown steadily (18).

Utilization of medicinal plants has almost doubled in Western Europe during that period. Ecological awareness, the efficacy of a good number of phytopharmaceutical preparations, such as ginkgo, garlic or valerian and increased interest of major pharmaceutical companies in higher medicinal plants as sources for new lead structures has been the main reasons for this renewal of interest. With the development of chemical science and pharmacognosy physicians began to extract chemical products from medicinal plants. A few examples of the products extracted from medicinal plants are - in 1920, quinine was isolated from Cinchona by the French pharmacist, Peletier & Caventou. In the mid-nineteenth century, a German chemist, Hoffmann obtained Aspirin from the bark of the willow. With the active principles in medicinal plants identified and isolated, plant-based prescriptions began to be substituted more and more with pure substances, which were more powerful and easier to prescribe and administer (19).

Phytomedicine almost went into extinction during the first half of the 21st century due to the use of the 'more powerful and potent synthetic drug'. However, because of the numerous side effects of these drugs, the value of medicinal plants is being rediscovered as some of them have proved to be as effective as synthetic medicines with fewer or no side effects and contraindications. It has been proved that although the effects of natural remedies may seem slower, the results are sometimes better on the long run especially in chronic diseases.

In the present study was carried out on the plant sample revealed the presence of medicinally active constituents. The phytochemical characters of the *Evolvulus alsinoides* leaf investigated and summarized in Table-1. The phytochemical screening *Evolvulus alsinoides* leaves showed that the presence of flavonoids, polyphenol, steroids, tannin, saponins, glycosides, anthroquinones, polyphenol, amino acid, alkaloids, carbohydrate, protein while steroids and Phlobatannins were absent.

Table 1: Phytochemical screening of *Evolvulus alsinoides*

S.No	Phytochemical analysis	Results
1	Carbohydrate	++
2	Amino acids	+
3	Polyphenols	+++
4	Tannins	+++
5	Anthroquinone	+++
6	Flavonoids	+++
7	Alkaloids	+++
8	Phlobatannins	-
9	Saponin	+++
10	Steroids	+++
11	phytosterol	-
12	Terpenoids	+++
13	Triterpenoids	++
14	Glycosides	+++

(+) Presence (-) Absence (++) High concentrations

Quantitative analysis

Quantitative analysis revealed that the *Evolvulus alsinoides* contain significant amount of phenols, alkaloids, saponin and terpenoids. Significant amount of Total Phenols (324.0 mg/gm), Alkaloids (35.0 mg/gm), Tannin(98.4 mg/gm), Saponin (121.0 mg/gm), Flavonoids 165.0mg/gm) (Table 2). The above phytoconstituents were tested as per the standard methods. This is because of the pharmacological activity of this plant is used to trace the particular compound.

Table 2: Qualitative analysis of *Evolvulus alsinoides* leaf

S.No	Test	Result (mg/gm)
1	Total Phenols	324.0
2	Alkaloids	35.0
3	Tannin	98.4
4	Saponin	121.0
5	Flavonoids	165.0

Vitamins

Vitamins are organic substances that are essential in tiny amounts for growth and activity of the body. They are obtained naturally from plant and animal foods. Organic in this definition refers to the chemistry and molecules of vitamins. The word organic means that the molecules of the substance contain the element carbon. The term also means

that vitamins can be destroyed and become unable to perform their functions in our bodies. Too much heat, certain kinds of light and even oxygen can destroy some vitamins. The amounts of vitamins ingested from food are measured in micrograms or milligrams.

Vitamin C, or ascorbic acid, is one vitamin humans cannot make; they have to get it from food. Vitamin C helps hold the cells together, heal wounds, and build bones and teeth. The best sources for vitamin C are citrus fruits, strawberries, melons, and leafy green vegetables. Vitamin C also helps to absorb and use Iron. It is important to protect the vitamins in fruits and vegetables from being destroyed; simple ways of doing this include refrigeration, washing them before cutting them, storing them in airtight containers, and avoiding high temperatures and long cooking times (20). The vitamins of the *Evolvulus alsinoides* leaves investigated and summarized in Table-3. The vitamin analysis of *Evolvulus alsinoides* showed that the presence of Vitamin D, E, C while vitamin A was absent.

Table 3: Qualitative analysis of Vitamins

S.No	Test	Result
1.	Vitamins A	-
2.	Vitamins C	+
3.	Vitamins D	+
4.	Vitamins E	+

(+) Presence (-) Absence

Minerals

All human beings require a number of complex organic/inorganic compounds in diet to meet the need for their activities. The important constituents of diet are carbohydrates, fats, proteins, vitamins, minerals and water (21). Every constituent plays an important role and deficiency of any one constituent may lead to abnormal developments in the body. Plants are the rich source of all the elements essential for human beings. There is a relationship between the element content of the plant and its nutritional status. Some elements are essential for growth, for structure formation, reproduction or as components of biologically active molecules while others have some other beneficial effects (22). The elements of the *Evolvulus alsinoides* leaves were investigated and summarized in Table-4. The following elements were found in *Evolvulus alsinoides* leaves. They are calcium, sodium, potassium, sulphate, phosphate, chloride, Nitrate, magnesium and iron while carbonate was absent in *Evolvulus alsinoides* leaves extract.

Table 4: Qualitative analysis of inorganic element in *Evolvulus alsinoides*

S.No.	Inorganic elements	Result
1	Calcium	++
2	Magnesium	+
3	Sodium	++
4	Potassium	++
5	Iron	-
6	Sulphate	+
7	Phosphate	++
8	Chloride	++
9	Carbonate	-
10	Nitrate	+

(+) Presence (-) Absence (++) High concentrations

Minerals are inorganic substances, present in all body tissues and fluids and their presence is necessary for the maintenance of certain physicochemical processes which are essential to life. Minerals are chemical constituents used by the body in many ways. Although they yield no energy, they have important roles to play in many activities in the body. Every form of living matter requires these inorganic elements or minerals for their normal life processes. Minerals may be broadly classified as macro (major) or micro (trace) elements. The third category is the ultra trace elements. The macro-minerals include calcium, phosphorus, sodium and chloride, while the micro-elements include iron, copper, cobalt, potassium, magnesium, iodine, zinc, manganese, molybdenum, fluoride, chromium, selenium and sulfur. The macro-minerals are required in amounts greater than 100 mg/dl and the micro-minerals are required in amounts less than 100 mg/dl. The ultra trace elements include boron, silicon, arsenic and nickel which have been found in animals and are believed to be essential for these animals. Evidence for requirements and essentialness of others like cadmium, lead, tin, lithium and vanadium is weak (23).

CONCLUSION

The results of this study clearly indicate that preliminary phytochemical analysis of *Evolvulus alsinoides* revealed presence of flavonoids, polyphenol, steroids, tannin, saponins, glycosides, anthroquinones, polyphenol, steroids, alkaloids, carbohydrate, protein while amino acid and Phlobatannins were absent. Quantitative analysis revealed that *Evolvulus alsinoides* rich amount of Total Phenols (324.0 mg/gm), Alkaloids (35.0 mg/gm), Tannin (98.4 mg/gm), Saponin (121.0 mg/gm), Flavonoids 165.0mg/gm) were presented. The vitamin analysis of *Evolvulus alsinoides* showed that the presence of Vitamin E, C and D while vitamin A was absent. The inorganic elements of *Evolvulus alsinoides* showed that the presence of calcium, sodium, potassium, sulphate, phosphate, chloride, Nitrate, magnesium and iron while carbonate was absent. The results of the present study concluded that *Evolvulus alsinoides* leaves may be a good source of phytochemicals, vitamins and minerals. Supplementation of this *Evolvulus alsinoides* leaves may be useful for human health associated emerging diseases such as cardiovascular diseases, diabetes, hypertension and cancer.

REFERENCES

- Hasler CM and Blumberg JB (1999) Phytochemicals: Biochemistry and physiology. Introduction. *Journal of Nutrition* 129: 756S-757S.
- Gibson EL, Wardel J, Watts CJ(1998). Fruit and Vegetable Consumption, Nutritional Knowledge and Beliefs in Mothers and Children. *Appetite*; 31: 205-228.
- Mathai K(2000). Nutrition in the Adult Years. In Krause's Food, Nutrition, and Diet Therapy, 10th ed., ed. L.K. Mahan and S. Escott-Stump; 271: 274-275.
- Arora A, Nair MG, Strasburg GM. Structure-activity relationships for antioxidant activities of a series of flavonoids in a liposomal system. *Free Radical Biology and Medicine*. 1998, 24: p1355-1363.
- BNF. Plants: Diet and health. In: Goldberg G (ed.) The Report of the BNF Task Force. Oxford: Blackwell. 2003. P25-23.
- Batra S, Seth S, Husain SA. Mineral elements status in breast cancer. *Breast Cancer Res Treat* 2002, 59:163-170.
- Amirjani MR (2010). Effects of salinity stress on growth, mineral composition, proline content, antioxidant enzymes of soybean. *American Journal of Physiology*, 5: 350-360.
- Harborne J (1973). Phytochemical methods London. Chapman and Hall, Ltd. Iinuma M, Tsuchiya H, Sato M, Yokoyama J, Ohyama M, Ohkawa Y,
- Draft Criteria Trease GE, Evans WC (1989). Pharmacognosy. 11th edn. Brailiar Tiridel can. Macmillan Publishers.U.S. (1984). Environmental

protection Agency document for carbon tetrachloride, criteria and standards Division, office of Drinking Water, Washington, DC.

10. Sofowara, A., (1993). Medicinal plants and Traditional medicine in Africa. *Spectrum Journal Ethnopharmacology*, V.90(2-3): P.249-252
11. Edeoga H.O. Okwu D. E. and Mbaebie B.O. (2005) Phytochemical constituents of some Nigerian medicinal plants. *African Journal of Biotechnology*. 4 (7), pp. 685-688
12. Van-Burden TP, Robinson WC (1981). Formation of complexes between protein and Tannin acid. *J. Agric. Food Chem.* 1: 77.
13. Obdoni BO, Ochuko PO and Scalbert A. (2001). Phytochemical studies and comparative efficacy of the crude extracts of some Homostatic plants in Edo and Delta States of Nigeria. *Global J. Pure Appl. Sci.* 8 p:203-208
14. Bohm BA, Kocipai-Abayazan R (1994). Flavonoids and condensed tannins from leaves of Hawaiian *vaccinium vaticulatum* and *V. calycinum*. *Pacific Sci.* 48: 458-463.
15. Pearson D, The Chemical Analysis of Food, 17th ed. Churchill Livingstone, London. 1976. pp 3-4.
16. Khandelwal, K.R (2006) Practical Pharmacognosy (16th ed.,) Nirali Prakashan, Pune.p98-106.
17. Narasinga Rao(2003). Bioactive phytochemicals in Indian foods and their potential in health promotion and disease prevention. *Asia Pacific Journal of Clinical Nutrition*: 12 (1): 9-22
18. Hamburger M, Hostettmann, K. (1991) Bioactivity in plants: the link between phytochemistry and medicine. *Phytochemistry*. 30; 3864-3874
19. Harvey A (2000). Strategy for discovering drugs from previously unexploited natural products. *Drug Discovery Today*; 5: 294-300.2003.
20. Okwu. D.E. Investigation into the medicinal and nutritional potential of *garcinia Kala* (Bitter kala) Heckel and Demettia tripetala G.Beaker Ph.D. Dissertation of Michael Okpara University of Agriculture Umudike , Nigeria pp. 5-10. 2003.
21. Indrayan, A.K., Sharma, S., Durgapal, D., Kumar, N. and Kumar, M., 2005. Determination of nutritive value and analysis of mineral elements for some medicinally valued plants from Uttaranchal. *Current Sci* 89, 1252- 1255.
22. NewWall, C.A., Anderson, L.A. and Phillipsan, J.D., 1996. *Herbal medicines- A guide for healthcare professionals*. The Pharmaceutical Press, London
23. Murray, C.B., Kagan, C.R., Bawendi, M.G. (2000). Synthesis and characterization of monodisperse nanocrystals and close-packed assemblies. *Annu Rev Mater Sci*, 30:545-610