


EFFECTS OF IMPLEMENTATION OF W.H.O SURGICAL SAFETY CHECK LIST: OUR INSTITUTIONAL ANALYSIS



General Surgery

KEYWORDS: Surgical safety; check list; patient care, implementation

Dr. S.J. Haridarshan

M.S, FMIS, FIAGES, Assistant Professor, Rajarajeswari Medical college and Hospital, Bangalore,India

Dr. Girish.C. S

MS., Post graduate, Rajarajeswari Medical college and Hospital, Bangalore,India

ABSTRACT

BACKGROUND: To err is human. Human errors and flaws in the system and quality of health care persist despite patient safety initiatives and identification of new strategies of improvement. A significant focus in surgery has been to identify strategies to improve patient safety and prevent postoperative complications and adverse events. The World Health Organization (WHO) surgical safety checklist has been shown to decrease mortality and complications and has been adopted worldwide.

METHODS: A retrospective pre-intervention study was done from September 2014 to August 2015, data analysed, necessary training, changes and awareness was implemented over one month and then a prospective study was done post intervention using WHO safety check list from October 2015 – September 2016 both at Rajarajeswari medical college and hospital, Bangalore, India by implementing the safety check list and by corrections of errors present during the pre-intervention phase.

RESULTS: A total of 216 patients were recorded during the pre-intervention phase and a total of 248 patients were included in the post intervention analysis after implementing the WHO safety check list. Our study showed a reduction of intra-operative surgical complications from 5.1% to 2.41% and from 6.48% to 4.44% in post-operative surgical complications after the implementation of safety check list.

There was a significant reduction in anesthesia complications, both during and after surgery post implementation. (Intra-operative-2.78% to 1.61%; Post-operative- 1.4% to 0.8%). We also observed a significant change in the death rates post implementation. Intra operative death reduced from 1.4% to 0.4% and post-operative deaths reduced from 12.04% to 8%.

CONCLUSION: Surgical safety check list improves the quality of patient care and improves perceptions on patient safety. It significantly reduces the morbidity and mortality by reducing the complications and improving the quality of treatment and health care. It is easy to understand and implement the surgical safety check list after training, proper guidance and education. It is a must to implement the surgical safety check list in all institutions to improve the patient safety and to reduce the errors in patient care.

INTRODUCTION:

Health care is a team effort. When human limitations are combined with organizational complexity, human error becomes inevitable. Average error rates of 10% is seen in hospital admissions⁽¹⁾

Numerous changes have since been advocated to improve patient safety like mandating minimum nurse to patient ratios, reducing working hours of trainee/resident doctors, introduction of 'care bundles' that improve patient outcomes, introduction of safety checklists and advances in the science of simulation and teamwork training.

The WHO launched Global Patient Safety Challenge: Safe Surgery Saves Lives (SSSL) in 2006 in response to the global need to improve outcomes in surgery.

A checklist is more than a technical solution; if used properly, it ensures that few things will be reviewed at certain times.

For these reasons, incident reporting appears to be a surrogate marker of safety culture

In our hospital, implementation of surgical safety checklist was started from September 2015.

AIMS AND OBJECTIVES:

The main aim of this study is to assess the effectiveness of surgical safety checklist in our Institution. We also study and analyse the errors in our operation theatre, compare with international standards, train and educate the theatre personnel and implement the necessary corrections to avoid errors and ensure maximum safety for the patients.

METHODS:

Study design

This study was conducted in the department of general surgery at Rajarajeswari medical college and hospital, Bangalore, India. All the elective surgical cases were included and all the emergency cases were excluded in our study.

A Retrospective study of pre-intervention from September 2014 to August 2015 was done to analyse the errors in surgical safety and then, a Prospective study of post intervention from October 2015 – September 2016 was done by implementing the safety check list and by corrections of errors present during the pre-intervention phase.

A local investigator was chosen who was trained in the identification and reporting of process, measures, errors and complications. This person worked on the study full-time and did not have clinical responsibilities at the study site. Data analysis and implementation of the check list was done by the principal investigator, in this case the author and his assistant, the co-author.

Ten essential objectives for safe surgery were identified:

- (i) correct site of surgery,
- (ii) provision of safe anaesthesia,
- (iii) management of airway problems,
- (iv) management of haemorrhage,
- (v) avoiding known allergies,
- (vi) minimizing the risk of surgical site infection,
- (vii) preventing the retention of swabs and instruments,
- (viii) accurate identification of specimens,
- (ix) effective communication within the surgical team and
- (x) routine surveillance of surgical outcomes.

Intervention

After collecting baseline data retrospectively, the local investigator was given information to identify errors and deficiencies and was then asked to implement the 19-item WHO safe-surgery checklist. The checklist consists of an oral confirmation by surgical teams at the time of sign in, time out and sign out.

Departments of Surgery and Anaesthesiology, the operation theatre personnel and the design team created an education program focused on the checklist. The education program consisted of oral, online education module and an educational video showing the checklist process, elements involved, theatre and perioperative service area. A weekly frequently asked question communications to surgical team members and theatre. The local investigator

introduced the checklist to operating-room staff, using lectures, written materials, or direct guidance. The checklist was introduced over a period of 1 month. Personnel training regarding the benefits of the checklist and the checklist process was conducted for a period of one month.



FIGURE 1

Data Collection

Standardized data sheets completed by the local investigators and the clinical teams involved in surgical care were analysed. Perioperative data included the demographic characteristics of patients, procedural data and the type of anaesthesia used as per the WHO surgical safety check list.

Patients were followed prospectively until discharge or for 30 days, whichever came first, for death and complications. Outcomes were identified through chart monitoring and communication with clinical staff. Completed data forms were directed to primary investigator.

WHO SURGICAL SAFETY CHECKLIST

FIGURE 2

Knowledge of the safety checklist was analyzed based on a questionnaire used before and after implementation for the surgeons, anesthesiologists and the theatre personnel. The scoring was done on a scale of ten and the following results were drawn as per Table 1.

	Before implementation	After implementation
Doctor (consultants)	6	9
Doctor (residents)	4	7
Anaesthetologists	5	8
Nursing staff	2	5
Technical staff	1	4

TABLE 1 QUESTIONNAIRE

INTRODUCTION	SETTING	OPERATION	POST PROCEDURE
Patient medically fit for procedure?	Crash kit available?	Patient identity, procedure and consent checked?	Inadvertent events documented?
Procedure details and type of anesthesia reviewed?	Oxygen source and suction checked?	Is the site and side marked and identified?	Patient and the doctors' satisfaction assessed?

Post procedure plans reviewed?	Anticipated duration recorded?	Patient monitoring as per protocol?	
		Adverse events discussed prior to surgery?	
		Antibiotic prophylaxis?	
		DVT prophylaxis?	

TABLE 2

The questionnaire as shown in Table 2 was based on a Likert type scale and a yes/no question. Post implementation also included the above questionnaire along with open ended questions like the overall satisfaction of using the check list, ease and benefits of using the check list and if it would be beneficial in implementing the check list for patient's safety.

RESULTS

A total of 216 patients were included in the pre-intervention study and a total of 248 patients were included in the post-intervention study. The intra-operative and the post-operative complications both due to surgery and the anaesthesia were assessed and recorded (Table 3).

INTRAOPERATIVE		POSTOPERATIVE	
SURGICAL	ANESTHETIC	SURGICAL	ANESTHETIC
Haemorrhage	Decreased saturation	Surgical site infections	Inadequate analgesia
Long duration of surgery	Inadequate fluid management / shock	Unplanned return to OT	Pneumonia
Wrong surgery	Cardiac arrest	Burst abdomen	Spinal headache
iatrogenic damage	Multiple pricks/intubations	Multiple transfusions	
	Inefficient induction/relaxation	Prolonged hospital stay	
		DVT	

TABLE 3

The effectiveness of the safety check list was studied and analyzed by comparing the intra operative and post operative complications before and after its implementation.

INTRAOPERATIVE SURGICAL COMPLICATIONS		INTRAOPERATIVE ANESTHETIC COMPLICATIONS		POSTOPERATIVE SURGICAL COMPLICATIONS		POSTOPERATIVE ANESTHETIC COMPLICATIONS	
Before implementation	After implementation	Before implementation	After implementation	Before implementation	After implementation	Before implementation	After implementation
11(5.1%)	6(2.41%)	6(2.78%)	4(1.61%)	14(6.48%)	11(4.44%)	3(1.4%)	2(0.8%)

TABLE 4

Our study showed a reduction of intra-operative surgical complications from 5.1% to 2.41% and from 6.48% to 4.44% in post-operative surgical complications after the implementation of safety check list (Table 4).

There was a significant reduction in anesthesia complications, both during and after surgery post implementation. (Intra-operative- 2.78% to 1.61%; Post-operative- 1.4% to 0.8%) as shown in Table 4.

Intraoperative death		Postoperative death	
Before implementation	After implementation	Before implementation	After implementation
3 (1.4%)	1 (0.4%)	26 (12.04%)	20 (8%)

TABLE 5

We also observed a significant change in the death rates post implementation. Intra operative death reduced from 1.4% to 0.4% and post-operative deaths reduced from 12.04% to 8% (Table 5).




FIGURE 3

The data we analyzed before and implementation of the safety check list was comparable with the studies done by B Haynes^[2] et al and Haugen et al^[3]. Complications before implementation in our study was 15.76% as compared to Haynes et al^[2] which was 21% and 2.93% in Haugen et al^[3] study. Post implementation, our study showed 9.26% in the complication rate while their studies showed 12% and 2.77% respectively (Figure 3).

Death rate in our study was 13.44% and 1.20% before and after implementation, while the study done by B Haynes et al^[2] had a comparison of 1.10% and 0.30% before and after implementation. Study done by Haugen et al^[3] had no mortality rate (Figure 3).

DISCUSSION

The implementation of WHO surgical safety checklist showed significant and positive changes, with overall improvement in the awareness of patient safety, communication, and quality of care. Improvements were noted in the perception of value and participation in the time-out process, surgical team communication, and the establishment and clarity of patient care needs.

Use of this check list will help to bring about effective modifications in the process of execution as an ongoing quality improvement in our institution.

The checklist has garnered significant worldwide enthusiasm, with programs implemented in 26 countries and more than 3,000 hospitals worldwide within 3 years of its introduction.^[4]

It has been suggested that the simplicity of the checklist facilitates a rapid and effective implementation process^[5]. The surgical safety checklist has been shown to improve postoperative morbidity and mortality^[6]. Poor communication between surgeons and surgical team members has been previously reported, and team training may be required for effective checklist use^[7].

Humans are fallible, and this checklist enhances consistency in surgical team performance at critical times, fostering good communication, teamwork, and a culture of patient safety.^[8]

There was improvement in patient safety following implementation of surgical safety checklist in our study suggested by a reduction in percentage of complications by 5.3% and percentage of deaths by 4.03%.

CONCLUSION

By this study, it is eminently evident that surgical safety checklist can be implemented in all the institutions.

Implementation of surgical safety checklist is a prerequisite in surgical health care team to reduce the errors associated and for improved patient safety.

Knowledge about surgical safety checklist in surgical team has scope for improvement.

REFERENCES:

1. N. Sevdalis, L. Hull1 and D. J. Birnbach. Improving patient safety in the operating theatre and perioperative care: obstacles, interventions, and priorities for accelerating progress. *British Journal of Anaesthesia*; 2012; 109:3-16
2. B. Haynes, G. Weiser, R. Berry et al. A surgical safety checklist to reduce morbidity and mortality in a global population. *N Engl J Med*. 2009; 360(3):491-499
3. A.S. Haugen, E. Softeland, G.E. Eide et al. Impact of WHO's surgical safety checklist on a safety culture in the operating theatre. *British J Anaesthesia*. 2013; 110(5): 807-815
4. World Health Organization. Safe Surgery Saves Lives. 2013 <http://www.who.int/patientsafety/safesurgery/en/index.html>. Accessed December 20, 2012.
5. Haynes AB, Weiser TG, Berry WR, et al. Safe Surgery Saves Lives Study Group. A surgical safety checklist to reduce morbidity and mortality in a global population. *N Engl J Med*. 2009 Jan 29; 360(5):491-499. Epub 2009 Jan 14.
6. Bliss LA, Ross-Richardson CB, Sanzari LJ, et al. Thirty-day outcomes support implementation of a surgical safety checklist. *J Am Coll Surg*. 2012 Dec; 215(6):766-776. Epub 2012 Aug 28.
7. Fourcade A, Blache JL, Grenier C, Bourgain JL, Minvielle E. Barriers to staff adoption of a surgical safety checklist. *BMJ Qual Saf*. 2012 Mar; 21(3):191-197. Epub 2011 Nov 7.
8. Changes in safety attitude and relationship to decreased postoperative morbidity and mortality following implementation of a checklist-based surgical safety intervention. Haynes AB, Weiser TG, Berry WR, Lipsitz SR, Breizat AH, Dellinger EP, Dziekan G, Herbosa T, Kibatala PL, Lapitan MC, Merry AF, Reznick RK, Taylor B, Vats A, Gawande AA, Safe Surgery Saves Lives Study Group. *BMJ Qual Saf*. 2011 Jan; 20(1):102-7.