



## COMPARATIVE STUDY OF OXIDATION OF dl-ALANINE BY DIFFERENT Cr(Chromium) (VI) OXIDANTS IN PARTIAL NON-AQUEOUS MEDIUM

### Chemistry

**Shilpa Rathor**

Research Scholar, Pacific academy of Higher Education and Research University, Udaipur

**B. K. Dangarh**

Asstt. Prof. Of Chemistry, Govt. P. G. College, Neemuch, M.P.

### ABSTRACT

A comparative study of oxidation of dl-Alanine by different Cr (VI) oxidants [PCC, PDC, and QDC] has been studied in DMF-H<sub>2</sub>O medium in the presence of perchloric acid spectrophotometrically. The reactions are found first order with respect to all three oxidants, [H<sup>+</sup>], and [dl-Alanine]. Michaelis-Menten type kinetics was observed. The reaction rate increased with increasing volume percentage of DMF in reaction mixture. The reaction was studied at different temperature [298-323K] & the activation parameters were computed. The oxidation product was identified as Cr (III) and acetaldehyde. The oxidation rate order was found with respect to different oxidants are: QDC > PDC > PCC.

### KEYWORDS:

Oxidation, dl-Alanine, HClO<sub>4</sub>, Oxidants.

### 1. INTRODUCTION

Chromium is one of the essential elements for proper metabolism of insulin, protein, fat and carbohydrate. But Cr (VI) is easily taken up by cells and undergoes reduction to give the trivalent form through other intermediate oxidation states such as Cr (V) and Cr (IV). Different Cr (VI) oxidants are used for the oxidation of amino acids.

Hiran et al. reported the oxidation of glycine<sup>[1]</sup> and tyrosine<sup>[2]</sup> by PBC in aqueous acetic acid in presence of perchloric acid. Kinetics of oxidation of Alanine<sup>[3]</sup>, L-Cysteine<sup>[4]</sup>, Methionine<sup>[5]</sup>, by PCC. Karim and Mohant<sup>[6-10]</sup> studied the oxidation of several amino acids by QDC in DMF in presence aqueous sulfuric acid. Dangarh & Thakur reported the kinetic and mechanistic study of oxidative transformation of some alpha-amino acid<sup>[11-13]</sup> by PDC in aquo-acetic acid medium. There seems to be no reports on the comparative study of oxidation of dl-Alanine by different Cr (VI) oxidants.

### 2. MATERIALS AND METHODS

Oxidants Pyridiniumchlorochromate<sup>[14]</sup>, Pyridiniumdichromate<sup>[15]</sup> Quinoliniumdichromate<sup>[16]</sup> were prepared by the method describe in the literature. The purity of the oxidants was checked by spectral analysis. dl-alanine (A.R. grade) was used as supplied and purity was checked by its melting point. All other chemicals used in this investigation were of analytical reagent grade. Double distilled water was used throughout in the investigation.

The rate measurements were carried out at 313K in 2M HClO<sub>4</sub> under the condition [alanine] >> [oxidants], in the solvent system of 30-70 % (v/v) DMF-H<sub>2</sub>O. The progress of the reaction was followed by measuring the absorbance of oxidants at 354 nm in one cm cell placed in the thermostatted compartment of Systronics VISISCAN -167 spectrophotometer.

The kinetics run were followed for more than 60-70% completion of the reaction and good first order kinetics were observed.

### 3. RESULTS AND DISCUSSION

#### 3.1 Stoichiometry and product analysis

To determine the stoichiometry of a reaction a known slight excess of oxidant was added to a known amount of alanine, by keeping all other condition of reaction is constant and after some hours the residual of oxidant was determined spectrophotometrically at 354nm. The product of oxidation was corresponding aldehyde i.e. acetaldehyde and was identified by its 2, 4-dinitro phenyl hydrazine derivative.

The stoichiometry of the reaction was found to correspond to the equation:



[Oxidant]                      [dl-Alanine]                      [Acetaldehyde]

#### 3.2 Effect of Oxidants

When substrate was in excess, the rate of which oxidants disappears followed the first -order kinetics. The first -order rate constants are independent of the initial concentration of the oxidants when varied in the range (0.5 – 4.0) X 10<sup>-3</sup> mol/dm<sup>3</sup> at 313K. Among three Cr (VI) oxidants dl-Alanine oxidized faster with QDC.

#### 3.3 Effect of Substrate

The effect of alanine concentration on the reaction rate studied at constant [Oxidants], [HClO<sub>4</sub>], temperature and DMF% [table 1]. The reaction rate increases with increasing of alanine concentration. A plot of log k<sub>obs</sub> against log [subs] gives a straight line [fig-1] with respect to three oxidants. This revealed that the rate of oxidation is first order with respect to substrate. It has been found that the plot of (1/k) versus (1/[subs]) is straight line with small intercept, indicates that Michaelis-Menten type kinetics is observed. The reaction proceeds through the formation of a complex between the oxidant and dl-Alanine.

The variation of the rate of oxidation of alanine with oxidants can be expresses as

$$d[\text{oxidants}]/dt = k[\text{dl-alanine}][\text{oxidants}]/K_m + [\text{dl-alanine}]$$

#### 3.4 Effect of H<sup>+</sup> ion

In order to study the effect of [H<sup>+</sup>] on the rate of reaction, kinetic runs were carried out keeping the concentrations of all other reactants are constant and varying the [H<sup>+</sup>] with HClO<sub>4</sub>. From [table 1], it is clear that the rate of the reaction increases with increase in [H<sup>+</sup>]. When the logarithms of k<sub>obs</sub> values were plotted against logarithms of the corresponding [H<sup>+</sup>], linear plots with a positive slope was obtained and indicate that first order reaction with respect to the hydrogen ion concentration. Under the present experimental conditions (1-3 mol dm<sup>-3</sup> [H<sup>+</sup>]), the protonated chromium (VI) is presumed to be the reactive species since the rate increases with increase in [H<sup>+</sup>].

#### 3.5 Effect of solvent composition

The influence of variation of dielectric constant of the medium was carried out by the changing DMF (% v/v) in the reaction medium, keeping other conditions remaining constants and the reaction rate was affected considerably. The rate of reaction increased with an increasing volume percentage of DMF [table 1], suggesting that a low dielectric medium favors the oxidation. A plot of log k against 1/D is linear with a positive slope for alanine under study. This indicates an ion-dipole type of interaction in the rate -determining step.

#### 3.6 Effect of Temperature

The reaction was studied at different temperatures (298-323 K), keeping other experimental conditions constant. The rate constant of the reaction was found to increase with increasing temperature [table 2]. From the Arrhenius plot of log k<sub>obs</sub> versus 1/T [fig-2], activation energy and other thermodynamic parameters was calculated [table 3]. The entropy of activation is negative as expected for bimolecular reaction. The negative value also suggests the formation of a cyclic intermediate from non-cyclic reactants in the rate determining step. The complex formation is proved by the plot of 1/k<sub>obs</sub> against 1/[dl-alanine]. It has been pointed out that if entropy of activation is negative

and small the reaction will be slow.

#### 4. CONCLUSION

At room temperature the reaction between alanine and oxidants are very slow in the low perchloric acid concentration in DMF-water medium. The oxidation of alanine by oxidants are first order with respect to [oxidants], [dl-Alanine] and [HClO<sub>4</sub>]. The Zwitter ionic form of alanine is oxidized to give aldehyde. The reaction was studied at different temperatures. In the temperature range of 298-323 K, Arrhenius equation is valid. The negative value of entropy indicates that the complex C is more ordered than reactant. The description of the mechanism is consistent with all experimental evidence. Among three Cr (VI) oxidants dl-Alanine oxidized faster with QDC. The reactivity of oxidation of dl-Alanine towards three oxidants was found to be QDC>PDC>PCC.

**TABLE NO. 1 Effect of [Substrate], [H<sup>+</sup>] and Solvent [Oxidants]=1 x 10<sup>-3</sup>M T=313 K**

| [Subs] x 10 <sup>2</sup> M | [HClO <sub>4</sub> ] x M | DMF % v/v | k <sub>obs</sub> X 10 <sup>5</sup> Sec <sup>-1</sup> |       |       |
|----------------------------|--------------------------|-----------|--|-------|-------|
|                            |                          |           | PCC  | PDC   | QDC   |
| 1                          | 2                        | 30        | 16.04  | 20.07 | 20.26 |
| 1.5                        | 2                        | 30        | 20.65  | 24.98 | 23.41 |
| 2                          | 2                        | 30        | 25.10  | 27.71 | 25.48 |
| 2.5                        | 2                        | 30        | 29.55  | 30.24 | 27.63 |
| 3                          | 2                        | 30        | 33.93  | 34.54 | 29.44 |
| 1                          | 1                        | 30        | 7.29   | 4.64  | 6.83  |
| 1                          | 1.5                      | 30        | 12.05  | 10.82 | 15.27 |
| 1                          | 2                        | 30        | 16.04  | 20.07 | 20.26 |
| 1                          | 2.5                      | 30        | 22.14  | 40.91 | 35.15 |
| 1                          | 3                        | 30        | 30.97  | 52.89 | 46.78 |
| 1                          | 2                        | 10        | 8.13   | 9.51  | 15.35 |
| 1                          | 2                        | 20        | 11.89  | 12.09 | 17.77 |
| 1                          | 2                        | 30        | 16.04  | 20.07 | 20.26 |
| 1                          | 2                        | 40        | 26.36  | 36.65 | 27.67 |
| 1                          | 2                        | 50        | 34.00  | 49.32 | 43.10 |


**TABLE NO. 2 [dl-Alanine] = 1 X 10<sup>-2</sup> M [HClO<sub>4</sub>] = 2M [Oxidants] = 1 X 10<sup>-3</sup>M [DMF]=30% v/v**

| Temperature In K | k <sub>obs</sub> X 10 <sup>5</sup> Sec <sup>-1</sup> |       |       |
|------------------|--|-------|-------|
|                  | PCC  | PDC   | QDC   |
| 298              | 5.29   | 9.78  | 7.94  |
| 303              | 7.98   | 10.01 | 9.09  |
| 308              | 13.93  | 16.46 | 13.08 |
| 313              | 16.04  | 20.07 | 20.26 |
| 318              | 26.63  | 33.66 | 27.21 |
| 323              | 43.06  | 49.43 | 52.50 |


**TABLE NO. 3 THERMODYNAMIC PARAMETERS [Substrate] = 1 X 10<sup>-2</sup> M [HClO<sub>4</sub>] = 2M [Oxidant] = 1 X 10<sup>-3</sup>M [DMF]=30% v/v T=313K**

| Alanine with | log A | ΔE <sub>a</sub> <sup>#</sup> kJ mol <sup>-1</sup> | ΔS <sup>#</sup> J mol <sup>-1</sup> K <sup>-1</sup> | ΔH <sup>#</sup> kJ mol <sup>-1</sup> | ΔG <sup>#</sup> kJ mol <sup>-1</sup> |
|--------------|-------|---|---|--------------------------------------|--------------------------------------|
| PCC          | 8.70  | 64.70   | -82.29  | 62.09                                | 87.85                                |
| PDC          | 7.04  | 54.17   | -114.06   | 51.56                                | 87.27                                |
| QDC          | 7.95  | 59.64   | -96.51  | 57.04                                | 87.24                                |

**Fig-1 Variation of rate with substrate concentration by different oxidants**



**Fig-2 Variation of rate with Temperature by different oxidants**



#### REFERENCES:

- Hiran B.L., Nalwaya N., and Jain A.; J. Ind. Chem. Soc., 79, 587, (2002).
- Hiran B.L., Nalwaya N., and Jain A.; Kinetics and Catalysis, 45(3), 371, (2002).
- Hiran B.L., Meena M. L. and kunthwal J.; Der Pharma Chemica, 2(5),470, (2010).
- Adari K. K., Nowduri A. and Vani P; Transition Metal Chemistry, 31,745, (2006).
- Sharma V., Sharma P.K. and Banerji K.K.; J. Indian Chem. Soc., 74(8), 607, (1997).
- Karim E., Mahanti M.K.; Oxidation Communication, 14, 157, (1991).
- Karim E., Mahanti M.K.; Oxidation Communication, 15, 211, (1992).
- Karim E., Mahanti M.K.; Oxidation Communication, 16, 1471, (1992).
- Karim E., Mahanti M.K.; Oxidation Communication, 19, 582, (1996).
- Karim E., Mahanti M.K.; Oxidation Communication, 21, 559, (1998).
- Dangarh B. K. and Thakur S. S.; Ultra Chemistry, 12(1), 9, (2016).
- Dangarh B. K. and Thakur S. S.; IJSR, 5(6), 370, (2016).
- Dangarh B. K. and Thakur S. S.; J. Sci. Res. Phar. Chem. Bio. Sci., 1 (2), 41, (2016).
- Core E.J. and Suggs W.T.; Tetrahedron Lett., 31, 2647, (1975).
- Corey E.J. and Shmidt G.; Tetrahedron Lett., 399, (1979).
- K. Balasubramanian and V. Prathiba, Ind. J. Chem., 25B, 326 (1986).