



STUDY OF OSTEOPOROSIS AND ITS RELATIVE FACTORS IN FACULTY OF SPORT AND PHYSICAL EDUCATION AND FACULTY OF MEDICINE

Orthopaedics

Al-Sadek, T*	Belhoul European Hospital – UAE, Dubai, Department of Orthopedics and Traumatology. *Corresponding Author
Al-Sadek, A	Medical University of Sofia – Bulgaria.
AL-Sadek, L	University Hospital “Prof. Dr. St. Kirkovich” - Stara Zagora, Bulgaria, Department of Dermatology & Venerology.
Khawatmi, N	Medical University of Sofia – Bulgaria.

ABSTRACT

Background: Osteoporosis is a metabolic bone disease. This complication will reduce bone mineral density (BMD) and reduce bone strength. Lack of sufficient physical activity and reception of insufficient levels of sunshine are the factors that contribute to this disorder. The present research seeks to study osteoporosis and its relative factors among faculty of sport and physical education in Palestine.

Materials and Method: This is a cohort research conducted on two groups including 100 students of the Faculty of sport and physical education and Faculty of Medicine. The status of all participants was tracked for 5 variables including vitamin D, calcium, Femoral neck BMD, Total lumbar spine BMD, and Total hip BMD for a period of six months (at the beginning and the end of the period) and the resulting raw data was analyzed using IBM SPSS Statistics v.21. The level of significance in this research was set to p-value < 0.05.

Results: This was a cohort research conducted on 200 participants in 2 groups consisting of the faculty of sport and physical education and Faculty of Medicine. The participants aged from 18 to 24 years old with an average age of 20.08 ± 1.06 years. The results obtained through studying the five variables indicated that after a six month follow up period, only vitamin D and calcium levels among the faculty of sport and physical education had increased significantly compared to the first assessment (P-value < 0.05). Other variables failed to exhibit a significant change within this period (P-value > 0.05). Regardless of statistical significance, the general trend of change among faculty of sport and physical education indicated increase, but a decreasing trend was observed among Faculty of Medicine.

Conclusion: Exposure to sunshine and physical activities play a major and significant role in increasing vitamin D and calcium levels. Otherwise, vitamin D deficiency and unfavorable consequences such as osteoporosis are inevitable.

KEYWORDS

Osteoporosis, Metabolic Diseases, Fracture, Bone Mass.

Introduction:

Osteoporosis and Rickets are important bone metabolic diseases. Rickets is caused by the lack of vitamin D and mineral materials. According to WHO (world health organization), osteoporosis is a skeletal disorder whose most important property is reduced bone strength. As a result, the patient with this complication will be more prone to bone fractures. WHO considers osteoporosis as the third most important global issue following heart diseases and cancers and it is titled silent epidemics (1). This complication is one of the most common metabolic diseases during the old age ranging from mild cases with no fracture to acute cases and various fractures. As the researchers believe, this disease is one of the risk factors contributing to death toll and disease. This disease plays a major role in the life quality of the elderly due to its high costs (2-4). Higher rates of osteoporosis and fractures associated with it are observed among women than men (1). In other words, 1 out of every 4 women suffers from this complication, while this ration among men is 1 to 8 (5). A frequency rate of 30% has been reported for this disease in Europe. A study conducted in Austria showed that as many as 470000 Austrian women older than 50 are suffering from osteoporosis (6). The results of many studies show that one woman out of every five older than 50 years old suffers from osteoporosis. The same research claims that most women older than 50 have experienced wrist, pelvic, and lumbar fractures (7). As for Iranian women, a frequency rate of 32.4% has been reported for lumbar osteoporosis while this frequency of femoral osteoporosis has been reported to be 5.9% (5). The results of the same research show that as many as 75% of all fractures are caused by osteoporosis (8). Neonates, the elderly, those exposed to insufficient sunshine, those who have problems absorbing fat, the black people, the obese and those who have undergone stomach bypass operation are in the danger of lack of Vitamin D (10). Fracture is one of the most important complications of osteoporosis. Reduced physical performance, pain, less mobility, depression and social seclusion are the most important consequences of fractures caused by osteoporosis. Medical and non-medical interventions are used to treat and prevent this complication. The most obvious feature of osteoporosis is the reduced level of minerals and bone matrix (11). The results of various researches points to the fact that exercising, taking sufficient levels of calcium and vitamin D and exposure to sunshine help slow down

reduction of bone mass (12). Proper and regular physical activities play a major role in preventing this disease (13). Further to improving bone health, proper physical activities help create balance and coordination in the body, improve muscle power and general body health (14). Most researches on osteoporosis have been focused on the elderly particularly on ladies in Postmenopausal period. Few researches have studied the prevalence of its relative factors among university students. What's more, the present research has also studied the role of the daily schedule of various faculty's in bone health. The present research seeks to study the relative factors of osteoporosis among various ranks of Palestinian students. As a matter of fact, the present research seeks to study and compare the effect of various schedules in the Faculty of sport and physical education and Faculty of Medicine. (in terms of morning exercise, physical activity and exposure to sunshine) on osteoporosis (assessed by measuring bone density) and its relative factors such as vitamin D and calcium levels.

Materials and Method:

This is a prospective, cohort research conducted on the students Faculty of sport and physical education (Al Aqsa university) and Faculty of Medicine Al Azhar University – Gaza, Palestine in 2014-2016. Those who had no previous history of taking Vitamin D or pathological fractures took part in the research. If the participants took medicines without doctor's prescription, they would be eliminated due to low levels of vitamin D. Based on previous studies and with due consideration of the inclusion and exclusion criteria, the participants were divided into two groups each consisting of 100 students. The participants took part in the research voluntarily and with due consideration of medical ethics principles after obtaining the written consent. 10% of the first three-years students in the Faculty of sport and physical education and Faculty of Medicine were evaluated prospectively. The participants were studied in terms of vitamin D and calcium levels and bone mineral density in the lumbar, femoral and hip areas. These laboratory assessments were conducted once at the beginning of the research. Without any intervention, these students were once again assessed in terms of vitamin D, calcium, and bone mineral density levels. IBM SPSS Statistics v.23 was used to analyze the data and the normality of data was studied for all variables. Proper statistical tests were utilized based on the normality status. Paired

Samples t-test and Wilcoxon Test were used to analyze the data. The level of statistical significance in this research (P-value) was set to less than 0.05.

Results:

The participants aged 18 to 24 with an average of 20.08 ± 1.06 years. The weight and height of the participants were also measured. It turned out that the participants weighed from 51 to 90 kg with an average of 70.13 ± 7.90 kg and their average height was 175.85 ± 5.48 cm. A review of age, weight, and height of participants in each group showed that those in the Faculty of Medicine exhibited higher means in all

three variables than those in the Faculty of sport and physical education. Wilcoxon non-parametric test was used to compare levels of Vitamin D in the first and second assessment for participants. The results showed that in spite of reduced vitamin D levels in the second phase of assessment among students of Faculty of Medicine Al Azhar university, this difference was not significant (P-value > 0.05). On the other hand, the results of this statistical test showed that the average level of vitamin D among students of the Faculty of sport and physical education (Al Aqsa university) in the second assessment had increased significantly compared to the first assessment (P-value < 0.05) (table 1).

Table 1: Ranking and comparison of vitamin D levels in the first and second assessment among the two groups

Wilcoxon Test								
	Variables		Mean	Std. Deviation	N-Mean Rank	P-Mean Rank	Z	P-Value
Faculty of Medicine students		F-D3	24.041	14.174	49.23	51.88	-0.120	0.904
		S-D3	23.675	7.553				
Faculty of sport and physical education students	Variables	F-D3	14.354	8.620	26.14	52.33	-8.053	0.001
		S-D3	19.595	8.290				

F: First stage measurement S: Second stage measurement

The vitamin D levels of the participants in the first and second phase of assessment was also studied. A review of data pointed to an unexpected fact indicating that a large number of students in both groups were in the deficient (Vit-D < 10) and insufficient (vit-D < 30) levels, but the improving trend of vitamin D levels was more obvious among cadet school students. Next, the calcium levels measured in two phases among Faculty of sport and physical education students and Faculty of Medicine students were analyzed and compared with one another. The

statistical analysis by Wilcoxon non-parametric test confirmed the fact that the calcium levels measured in the two phases exhibit no significant change among medical school students and this level has stayed relatively constant and fixed (P-value > 0.05). However, observations point to the fact that the calcium levels measured in the second phase were significantly more than what was observed in the first phase (P-value < 0.05) (table 2).

Table 2: Ranking and comparison of calcium levels in the first and second phases of assessment among the two groups

Wilcoxon Test								
	Variables		Mean	Std. Deviation	N-Mean Rank	P-Mean Rank	Z	P-Value
Faculty of Medicine students		F-Calcium	9.58600	0.482841	47.55	47.42	-1.447	0.148
		S-Calcium	9.51600	0.605817				
Faculty of sport and physical education students	Variables	F-Calcium	9.43700	0.446910	15.63	49.42	-8.242	0.001
		S-Calcium	9.93800	0.421584				

F: First stage measurement S: Second stage measurement

An analysis of the difference between the measured values of Total lumbar spine BMD in the both defined stages for both groups was one of the main goals of this research which we will discuss in details here. In this analytical process where the statistical parametric test of Paired

Samples T-test was used for both groups, it turned out that after 6 months no statistically significant difference was observed between the two groups in terms of lumbar spine BMD levels (P-value > 0.05) (table 3).

Table 3: Total lumbar spine BMD comparison in the double stages of measurement in both groups

Paired Samples Statistics							
	Variables		Mean	Std. Deviation	t	d	P-Value
Faculty of medicine students		F-Total lumbar spine BMD	0.98625	0.082785	1.727	99	0.087
		S-Total lumbar spine BMD	0.97602	0.079998			
Faculty of sport and physical education students	Variables	F-Total lumbar spine BMD	0.98694	0.091588	-1.366	99	0.175
		S-Total lumbar spine BMD	0.99493	0.083123			

F: First stage measurement S: Second stage measurement

Femoral neck BMD is another parameter measured in this research analyzed following two stages of measurement in each group. The results of parametric statistical test of Paired Samples t-test among medical school students failed to show a statistically significant difference between the values recorded in both stages (P-value > 0.05).

As a consequence of Wilcoxon non-parametric statistical test failed to show a statistically significant difference between the first measurement data and the first stage among Faculty of sport and physical education students (P-value > 0.05) (table 4).

Table 4: Femoral neck BMD comparison in the double stages of measurement in both groups

Paired Samples Statistics							
	Variables		Mean	Std. Deviation	t	d	P-Value
Faculty of Medicine students		F-Femoral neck BMD	0.96553	0.127235	1.438	99	0.154
		S-Femoral neck BMD	0.96164	0.129567			
Faculty of sport and physical education students	Variables	F-Femoral neck BMD	1.01585	0.132028	47.66	41.26	-1.948
		S-Femoral neck BMD	1.01055	0.130832			

F: First stage measurement S: Second stage measurement

Another goal of the research was to study, investigate and compare Total hip BMD changes within the period of research (6 months) for each group. The results of the statistical analysis of non-parametric Wilcoxon test showed that in spite of rising changes in the second stage

of measurement, no statistically significant difference was observed between the two stages of Total hip BMD measurement among the two groups (P-value > 0.05) (table 5).

Table 5: Total hip BMD comparison in the double stages of measurement in the two groups

Wilcoxon Test								
	Variables		Mean	Std. Deviation	N-Mean Rank	P-Mean Rank	Z	P-Value
Faculty of Medicine students		F- Total hip BMD	1.00620	0.114051	47.66	41.26	-1.783	0.075
		S- Total hip BMD	1.01317	0.099195				

Faculty of sport and physical education students	Variables	F- Total hip BMD	1.08741	0.123115	53.42	39.53	-1.860	0.063
		S- Total hip BMD	1.08256	0.144709				
F: First stage measurement S: Second stage measurement								

Discussion:

Various researches have pointed to factors such as calcium shortage, lack of vitamin D, lack of mobility and proper physical activities, insufficient exposure to sunshine, genetic and inherited factors, Menopause and previous history of certain diseases are the factors that intensify osteoporosis. Vitamin D deficiency (VDD) creates imbalance and unnatural conditions in levels of calcium, phosphor, and bone metabolism. VDD contributes to reduced absorption of calcium and phosphor from your diet and this will increase levels of parathyroid hormone (PTH). Higher osteoclastic activities will create weak local bone centers as a result of parathyroid hormone which will culminate in reduced bone mineral density (BMD) and osteoporosis (15). Measuring bone mineral density (BMD) is one of the most important methods to diagnose osteoporosis. Considering the great importance of this issue in the health and life quality of people, the researchers in the present study have studied osteoporosis and its relative factors among various ranks of the students of Palestine. Various groups are working on osteoporosis and the factors that affect it and they keep publishing their results. A review of the related literature and an analysis of the results achieved by those researchers and comparing them with the present research points to a general alignment and rational agreement between data. In a 20-year cohort research ending in 2014, Lindqvist PG et al studied the effects of receiving sunshine on the death toll regardless of the disease (all-cause mortality). They arrived at the conclusion that failing to expose to sunshine is a risk factor for all-cause mortality. Although the first glance at this research fails to show a correlation between this research and its goals with our research, the general view and conclusion of our research can justify its correlation with the above-said study (16). Based upon the reports released in a review research by Holick in 2007(17) and according to the epidemiologic research by Eobinson et al (18), the global rate of Vitamin D insufficiency is 50%. A glance at the present research shows that the rate of insufficient vitamin D in both groups at the beginning and at the end of the research was more than 50% which shows the insufficient rate of vitamin D has a great prevalence. At the end of the research, the rate of vitamin D insufficiency was 15% less than the global level which indicates the positive effect of physical activity and exposure to sunshine on vitamin D and calcium levels. Many authors such as Fu X et al (19), Langsetmo et al (20), Moayyeri (21), Zernicke et al (22) have reported the positive effect of physical activity on BMD through mechanical loading mechanism. A clear example of these researches are athletes who have a higher BMD as a result of greater mechanical loading mechanism (23). In a research by Alghadir et al (2015) in order to assess the role of physical activities in bone mineral density (BMD) and bone metabolism indicators among 350 young volunteers, it was shown that body mass index (BMI), physical activity, low calcium consumption and lifestyle play a major role in bone minerals density and pre-awareness of osteoporosis among adults (24). Taking into consideration the intangible and insignificant change of BMD in our research, we may conclude that changes in this variable were not significant possibly due to the limited time of the research and absence of any intervention.

Conclusion:

Physical activity and exposure to sunshine plays a major and central role in the increasing trend of vitamin D and calcium level in the group of Faculty of sport and physical education students and absence of these factors among Faculty of Medicine students is responsible for reduced levels of vitamin D and calcium. We may also predict that reduced levels of vitamin D and calcium caused by lack of physical activity, unhealthy diet, insufficient exposure to sunshine, etc. will reduce BMD level over a long period and may culminate in Osteopenia and Osteoporosis.

References:

1. WHO Study Group WTRS.
2. Chrischilles E, Shireman T, Wallace R. Costs and health effects of osteoporotic fractures. *Bone*. 1994;15(4):377-86.
3. Darvishi M, Saeedi AA, Amini F. Evaluation of Vitamin D Levels in Patients with Tuberculosis. *The Journal of Clinical and Analytical Medicine*. 2017;9(Special Issue 2).
4. Morris R, Masud T. Measuring quality of life in osteoporosis. *Age and ageing*. 2001;30(5):371-3.
5. Khan A. Advances in osteoporosis therapy. 2003 update of practical guidelines. *Canadian family physician*. 2003;49(4):441-7.
6. Jahlka B, Dörner T, Terkula R, Quittan M, Bröll H, Erlacher L. Health-related quality of life in patients with osteopenia or osteoporosis with and without fractures in a geriatric

rehabilitation department. *WMW Wiener Medizinische Wochenschrift*. 2009;159(9):235-40.

7. Ström O, Borgström F, Kanis JA, Compston J, Cooper C, McCloskey EV, et al. Osteoporosis: burden, health care provision and opportunities in the EU. *Archives of osteoporosis*. 2011;6(1):59-155.
8. Keramat A, Patwardhan B, Larjani B, Chopra A, Mithal A, Chakravarty D, et al. The assessment of osteoporosis risk factors in Iranian women compared with Indian women. *BMC musculoskeletal disorders*. 2008;9(1):28.
9. Wagner CL, Greer FR. Prevention of rickets and vitamin D deficiency in infants, children, and adolescents. *Pediatrics*. 2008;122(5):1142-52.
10. Bischoff-Ferrari HA, Dawson-Hughes B, Staehelin HB, Orav JE, Stuck AE, Theiler R, et al. Fall prevention with supplemental and active forms of vitamin D: a meta-analysis of randomised controlled trials. *BMJ (Clinical research ed)*. 2009;339:b3692.
11. Gallagher JC. Effect of early menopause on bone mineral density and fractures. *Menopause*. 2007;14(3):567-71.
12. Kelley GA, Kelley KS. Exercise and bone mineral density at the femoral neck in postmenopausal women: a meta-analysis of controlled clinical trials with individual patient data. *American journal of obstetrics and gynecology*. 2006;194(3):760-7.
13. Ernst E. Can exercise prevent postmenopausal osteoporosis? *British journal of sports medicine*. 1994;28(1):5.
14. Henderson NK, White CP, Eisman JA. The roles of exercise and fall risk reduction in the prevention of osteoporosis. *Endocrinology and metabolism clinics of North America*. 1998;27(2):369-87.
15. Nair R, Maseeh A. Vitamin D: The "sunshine" vitamin. *Journal of Pharmacology & Pharmacotherapeutics*. 2012;3(2):118-26.
16. Lindqvist PG, Epstein E, Landin-Olsson M, Ingvar C, Nielsen K, Stenbeck M, et al. Avoidance of sun exposure is a risk factor for all-cause mortality: results from the Melanoma in Southern Sweden cohort. *Journal of internal medicine*. 2014;276(1):77-86.
17. MF. H. Vitamin D deficiency. *The New England journal of medicine* 2007;357(3):266-81 Epub 2007/07/20. 2007.
18. Robinson PJ, Bell RJ, Lanzafame A, Kirby C, Weekes A, Piterman L, et al. The prevalence of vitamin D deficiency and relationship with fracture risk in older women presenting in Australian general practice. *Australasian journal on ageing*. 2013;32(3):177-83.
19. Fu X, Ma X, Lu H, He W, Wang Z, Zhu S. Associations of fat mass and fat distribution with bone mineral density in pre- and postmenopausal Chinese women. *Osteoporosis international : a journal established as result of cooperation between the European Foundation for Osteoporosis and the National Osteoporosis Foundation of the USA*. 2011;22(1):113-9.
20. Langsetmo L, Hitchcock CL, Kingwell EJ, Davison KS, Berger C, Formso S, et al. Physical activity, body mass index and bone mineral density-associations in a prospective population-based cohort of women and men: the Canadian Multicentre Osteoporosis Study (CaMos). *Bone*. 2012;50(1):401-8.
21. Moayyeri A. The association between physical activity and osteoporotic fractures: a review of the evidence and implications for future research. *Annals of epidemiology*. 2008;18(11):827-35.
22. Zernicke R, MacKay C, Lorincz C. Mechanisms of bone remodeling during weight-bearing exercise. *Applied physiology, nutrition, and metabolism = Physiologie appliquee, nutrition et metabolisme*. 2006;31(6):655-60.
23. Kohrt WM, Bloomfield SA, Little KD, Nelson ME, Yingling VR. American College of Sports Medicine Position Stand: physical activity and bone health. *Medicine and science in sports and exercise*. 2004;36(11):1985-96.
24. Alghadir AH, Gabr SA, Al-Eisa E. Physical activity and lifestyle effects on bone mineral density among young adults: sociodemographic and biochemical analysis. *Journal of Physical Therapy Science*. 2015;27(7):2261-70.