



KNOWLEDGE AND PRACTICES AMONG NURSES ON POST EXPOSURE PROPHYLAXIS OF HIV IN MEDICAL COLLEGE, JHANSI (UP)

Community Medicine

Vimal Arya

Assistant Professor, Department of Community Medicine, MLBMC, Jhansi, U.P. -
Corresponding Author

B.N.Singh

Assistant Professor, Department of Community Medicine, WCMS, Jhajjar, Haryana -
Corresponding Author

ABSTRACT

Background: Hospital staffs, especially the nurses who have direct contact with patients and body substances are at higher risk of occupational hazards.

Objectives: To assess the knowledge and practice regarding post exposure prophylaxis of HIV among the nurses and to find out association between Knowledge and selected demographic variables.

Methodology: The study was conducted among the nurses working in medical-surgical units of MLBMC. 80 Nurses working in medical and surgical units were selected by stratified random sampling technique. Data were collected by using semi structured questionnaire. Chi-square was used to find out association.

Result: Majority of the respondents fall in the group of 20-25 years, mean age=24.28, 81.2 % were staff nurses. Regarding the knowledge about PEP, 87.5 % knew the meaning of post exposure prophylaxis of HIV. 81.2 % had the knowledge regarding the first aid for an occupational exposure.

Conclusion: Knowledge regarding PEP was found to be satisfactory in majority of respondents but practice was found to be less.

KEYWORDS:

Knowledge, Practices, HIV, Post Exposure Prophylaxis (PEP), Nurse.

Introduction- AIDS is far more than a health crisis. It is a threat to development itself." [1]

The global AIDS epidemic continues to grow and there is concerning evidence that some countries are seeing resurgence in new HIV infection rates which were previously stable or declining [2]. PEP is a medical response given to prevent the transmission of pathogens after potential exposure. PEP for HIV refers to a set of comprehensive services to prevent HIV infection in exposed individuals. These services include, first aid care, counselling and risk assessment, HIV testing based on informed consent, and depending on risk assessment, the provision of short term (28 days) antiretroviral (ARV) drugs, with follow up and support [3]. After the evaluation, health care workers should provide counselling on risk-reduction behaviour to the exposed person regardless of how the individual was exposed, and of whether or not antiretroviral (ARV) drugs will be recommended for PEP, as such, counselling can reduce the risk of future exposures [4]. With increasing number of HIV positive clients in eastern region of Uttar Pradesh and nurses being at high risk to occupational transmission of HIV the investigator felt need to assess the knowledge and practice regarding PEP of HIV among the nurses.

Material and Methods-

Research design: Descriptive study design. **Research setting:** The study was conducted among the nurses working in medical-surgical units of MLBMC, Jhansi (UP) **Sample:** Nurses working in medical unit I, II, III and surgical unit I, II, III, Who fulfil the set selection criteria, were included in this study. **Sample size:** 80 nurses were included in the study. **Sampling technique:** Population proportionate Stratified sampling technique was used to select the study subject. Two strata were made, one medical and one surgical unit. From both the strata using population proportionate method 40 nurses were selected from Medical units and 40 were selected from the surgical units randomly. **Research instrument:** Semi structured questionnaire was used to find out the demographic profile along with knowledge, and practices of nurses regarding post exposure prophylaxis of HIV. Pre-testing of the tool was done among 10% of total subjects. A detailed list of all the nurses was prepared before starting the data collection. Ethical clearance from concerned authorities was obtained. Informed verbal consent from each subject was obtained prior to interview.

Statistical Analysis- Data was entered and analyzed on Epi-info. Descriptive statistics i.e. percentage, mean and standard deviation was used to report findings of the study. Inferential statistics i.e. Chi Square test, was used to find the association between the knowledge and selected demographic variables.

Results –

Table 1- Socio-demographic details of nurses (n=80)

Factors	Category	Frequency (%)
Age (yrs)	<20	4(5)
	20-25	35(44)
	25-30	20(25)
	>30	21(26)
Hospital units	Medicine I	20 (25)
	Medicine II	10 (12.5)
	Medicine III	10 (12.5)
	Surgery I	20 (25)
	Surgery II	10 (12.5)
	Surgery III	10 (12.5)
Experience	≤1 year	10 (12.5)
	1-5 years	40 (50)
	>5 years	30 (37.5)
Designation	Staff nurse	65 (81.25)
	Senior staff nurse	10 (12.5)
	Matron	5 (8.75)

Table 1 shows total 80 nurses including 40 from medical units and 40 from surgical units were included in the study. Majority of respondents (44%) belonged to the age group of 20-25 years with the mean age=27.28, SD=3.4, range=18-35 yrs. Majority (81.25) of respondents were staff nurses followed by senior staff nurse (19.5%). Regarding experience 50% of respondents had an experience of 1-5 years followed by >5 years (37.5%).

Table 2- Training on HIV and PEP (n=80)

Characteristics	Category	Frequency (%)
Training on HIV	Yes	30 (37.5)
	No	50 (62.5)
Duration	2 days	12 (15)
	4 days	18 (22.5)
Training on PEP	Yes	15 (18.75)
	No	65 (81.25)
Duration	1 hour	15 (18.75)

Table 2- shows that only 37.5% of respondents had received in-service training on HIV/ AIDS and among those who received training on HIV/AIDS, 60% of respondents had taken the training of 4 days followed by 40% of 2 days. Regarding in- service training on post exposure prophylaxis (PEP) of HIV, only 18.75 % of them had received in- service training on PEP of HIV and the duration of the course was 1 hour.

Table 3- Knowledge regarding PEP and drugs used in it (n=80)

Characteristics	Category	Frequency (%)
Definition of PEP*	Yes	70 (87.5)
	No	10 (12.5)
Necessity*	Yes	70 (87.5)
	No	10 (12.5)
First aid*	Yes	65 (81.25)
	No	15 (18.75)
Duration	Yes	15 (18.75)
	No	65 (81.25)
ART	Yes	50 (62.5)
	No	30 (37.5)
Combination drug therapy	Yes	30 (37.5)
	No	50 (62.5)

*- statistically significant association (p<0.05)

Table 3- shows that 87.5% of respondents had the knowledge about the meaning of post exposure prophylaxis of HIV and same percentage of respondents, knew the necessity of post exposure prophylaxis (PEP) for nurses after occupational exposure with HIV source. Similarly 81.25% of respondents had the knowledge about the first aid for an occupational exposure and 18.75% of them could know the duration of PEP of HIV is 28 days. Result also reveals that 62.5% of respondents had the knowledge about the anti retroviral drug which was first used as PEP of HIV (Zidovudine (ZDV)), 37.5% of respondents had the knowledge about the combination of two drug therapy.

Table 4- Practices receiving PEP after exposure (n=80)

Characteristics	Category	Frequency (%)
Did you ever seek help from PEP of HIV services	Yes	20 (25)
	No	60 (75)
Did you try to find out HIV status of source	Yes	24 (30)
	No	56 (70)
Did you receive pre-test and post test counselling for that exposure	Yes	5 (8.75)
	No	75 (93.75)
Did you check your HIV status for that exposure	Yes	5 (8.75)
	No	75 (93.75)
Did you ever received ART	Yes	5 (8.75)
	No	75 (93.75)

Table 4 shows that only 25% of respondents had sought help from PEP of HIV services, only 30% of them tried to find out the HIV status of the source, only 6.25% of them received pre test and post test counselling for that exposure, 31% of them checked their HIV status for that exposure.

Discussion- In the present study majority of respondents (44%) belonged to the age group of 20-25 years with the mean age=27.28, SD=3.4, range=18-35 yrs. Majority (81.25) of respondents were staff nurses followed by senior staff nurse (19.5%). Regarding experience 50% of respondents had an experience of 1-5 years followed by >5 years (37.5%). Only 37.5% of respondents had received in-service training on HIV/ AIDS and among those who received training on HIV/AIDS, 60% of respondents had taken the training of 4 days followed by 40% of 2 days. The finding of the current study is contradictory to the finding of the study conducted by Lamichanne J, Aryal B. and Sharma Dhakal K. in Medical Colleges of Chitwan District, revealed that cent percent respondents had not received any training on post exposure prophylaxis of HIV. Seventy-eight percent of respondents answered that self learning was their sources of information regarding PEP followed by co-workers (48%), working experience (11%) & mass media (2%) [5]. The finding of current study is higher to the finding of a study of Baheti, Tullu & Lahiri which showed that only 40.8% nurses had knowledge about the fact that the exposed site must be immediately washed with soap and water [6]. Bairy et al showed that 98% nurses had knowledge that the first-aid procedure after needle prick injury is to wash the site with soap and water and in the present study only 5.0% of them had the knowledge about the best time for initiation of Post exposure prophylaxis and 30.0% of them could know the duration of PEP of HIV is 28 days [7]. Similarly this finding is also supported by a study of Avachat, Phalke & Dhumale which showed that, 47% of respondents' knew when post

exposure prophylaxis should start [8,9] The finding of current study is different than the finding study of Owolabi et al which showed that only 30.9% of respondents' could correctly identify the drugs used and duration of PEP [10].

Conclusions- Based on the findings of the study, it is concluded that nurses working on medical and surgical units of MLBMC had fair level of knowledge about PEP of HIV. Knowledge regarding PEP was found to be satisfactory in majority of respondents but practice was found to be less. And there is significant association between the knowledge with demographic variables but in few characteristics only. Hence educational intervention should be carried out periodically to enhance their practice.

Acknowledgement- We would like to express our sincere gratitude to the Head of Department of Medicine and Surgery for providing us the opportunity and facilities to conduct this research.

Conflict of interest- none declared

References-

- Annan K. Quotes for fighting AIDS [internet]. 2009 [cited 2017 Mar 17]. Available from :URL:http://aids-quotes.htm
- Global AIDS epidemics [online]. 2009 [cited 2017 Feb 20]. Available from: URL:http://data.unaids.org/pub/FactSheet/2009/20091124-FS-global-en.pdf.
- Post exposure prophylaxis for HIV infection: clinical protocol for the WHO European region [online]. 2007 [cited 2017 Feb 18]; 469-88. Available from: URL: http://www.euro.who.int/document/SHA/e90840-chapter-13.pdf.
- Lee R. Occupational transmission of blood borne diseases to healthcare workers in developing countries: meeting the challenges. *Journal of Hospital Infection* [serial online] 2009 [cited 2017 Apr 4]; 72:285-91. Available from: URL: http://www.elsevierhealth.com/journals/jhin
- Lamichanne Jharna, Aryal Bijay, Dhakal Kalplana Sharma. Knowledge of Nurses on post exposure prophylaxis of HIV in Medical Colleges of Chitwan District, Nepal. *Int J Pharm Biolog Arch.* 2012; 3(6): 1394-9.
- Baheti AD, Tullu MS, Lahiri KR. Awareness of Health Care Workers Regarding Prophylaxis for Prevention of Transmission of Blood-Borne Viral Infections in Occupational Exposures. *AlAmeen J Med Sci.* 2010; 3
- Bairy KL, Ganaraja B, Indira B, Thiyagar N, Choo CM, See CK. Awareness of post-exposure prophylaxis guidelines against occupational exposure to HIV in Hospital Sungai Petani. *Med J Malaysia.* 2005;60(1)
- Alenyo R, Fualal J, Jombwe JJ. Knowledge, attitude and practices of staffs towards post-exposure prophylaxis for HIV infection at Mulago Hospital in Uganda. *East and Central African Journal of Surgery* [serial online] 2009 Jul-Aug [cited 2017 Feb 18]; 14(2) Available from: URL:http://www.bioline.org.br/pdf?js09042
- S Avachat, DB Phalke, GB Dhumale. Awareness regarding pre- and postexposure prophylaxis for human immunodeficiency virus/acquired immunodeficiency syndrome among nursing students. *Indian Journal of community medicine online.* 2007. Available from: http://www.ijcm.org.in/article.asp?issn=0970
- Owolabi RS, Alabi P, Ajayi S, Daniel O, Ogundiran A, Akande TM, et al. Knowledge and practice of post-exposure prophylaxis (PEP) against HIV infection among health care providers in a tertiary hospital in Nigeria. *J Int Assoc Physicians AIDS*