



Elevation of Free Prostate - Specific Antigen (fPSA) in Cancer Patients Adenosine Deaminase (ADA) Binds and Stimulates Plasminogen (pg) activation on 1-LN Human Prostate Cancer cells.

Urology

M.A. Majid Adil	Department of Urology, Princess Esra Hospital & Research Centre, Deccan College of Medical Sciences, Hyderabad, Telangana State, India
Shaikh Mahmood	Department of Physiology, Owaisi Hospital & Research Centre, Deccan College of Medical Sciences, Hyderabad, Telangana State, India
Mohd. Abbas Hyder	Department of Biochemistry, Owaisi Hospital & Research Centre, Deccan College of Medical Sciences, Hyderabad, Telangana State, India
Shahana Sarwar	Department of Biochemistry, Princess Esra Hospital & Research Centre, Deccan College of Medical Sciences, Hyderabad, Telangana State, India

ABSTRACT

The Prostate Specific Antigen (PSA) is the best and most commonly used tumor marker for prostate in clinical practice these days^(1,2). It is a glycoprotein has molecular weight of 33KD. PSA is very routinely used in the diagnosis and follow up of prostate disease, specially prostate cancer^(3,4). Prostate Specific Antigen (PSA) is formed in two different molecular types in serum; they are free PSA (fPSA) and complex PSA (cPSA), the serum of which form the total PSA (tPSA). fPSA is mostly used in the detection of Prostate cancer⁽⁵⁾. Free PSA 28 kda is eliminated by glomerular filtration. The data revealed that the patients with end stage of renal dysfunction prolonged dialysis with increase percent of fPSA^(6,7,8,9).

Introduction: Adenosine Deaminase (ADA) is expressed in intracellular by all the cells. It is also associated with the cell surface multifunctional. Glycoprotein CD26/ dipeptidyl peptidase IV. CD26 is expressed on the surface of human prostate cancer. 1-LN cells acting as a receptor for plasminogen (pg). Since ADA and pg bind CD26 of distinct nearby sites. A possible interaction is found between these two proteins on the surface of 1-LN cells. Human ADA binds to CD26 on the surface 1-LN cells. Thus PSA, ADA may be factors regulating events in prostate cancer cells that occur when pg binds to the cell surface and in activated in prostate cancer (10-13).

Method: A total number of 201 patients with age group of (Median age 58 years, interquartile age range 47-68) with Chronic Kidney Disease (CKD). Their Glomerular Filtration Rate (GFR) was 24 ml/min/1.73m². (interquartile range is 18-34) by determined by iohexol clearance method. A separate biochemical and quantitative estimation was done for serum Adenosine Deaminase (ADA) and Plasminogen (pg) was done by fluoroimmunoassay method and are found elevated as serum Adenosine Deaminase (ADA) > 10IU/L and plasminogen (pg) > 0.2 IU/L in prostate cancer patients (14-17). Control included 350 subjects, attended a positive cancer screening test with no diagnostic prostate cancer.

Results: Median t PSA levels and percent of f PSA were significantly higher (p<0.001) in patients with renal dysfunction, 0.45µg/L and 48.2%. Compared to control, 0.28µg/L and 28.8% respectively. Effect of the interaction between Plasminogen (pg) and Adenosine Deaminase (ADA) on plasminogen activation or Adenosine Deaminase (ADA) enzymatic activity that plasminogen (pg) binds to CD26 on 1-LN cells (18-20). The present study shows the effect of increasing concentration of ADA on pg activation by 1-LN cells monolayer are single pg concentration 0.2 µM and also on pg activation by t-PSA.

Conclusion: The present study of percent of free Prostate Specific Antigen (fPSA) Adenosine Deaminase (ADA) on plasminogen (pg) are significantly influenced by moderately impaired renal function in patients with Chronic Kidney Disease (CKD) and in prostate cancer.

KEYWORDS

Prostate Specific Antigen (PSA), Adenosine Deaminase (ADA), Plasminogen (pg), CD26, 1-LN Cells, Chronic Kidney Disease (CKD), Glomerular Filtration Rate (GFR) and prostatic cancer.

Introduction:

The estimation of serum prostate specific antigen (PSA), Adenosine Deaminase (ADA) and Plasminogen (pg) are the important clinical methods for the detection and monitoring prostate cancer. PSA is found in two different molecular forms in the blood, free prostate specific antigen (fPSA), and complexed prostate specific antigen (cPSA). Free PSA (fPSA) with a molecular weight of ~28 kDa and Complexed PSA (cPSA) ~90 kDa. Complexed PSA (cPSA) alpha-1-antichymotrypsin, alpha-1-antitrypsin and alpha-2-macroglobulin. The serum of fPSA and cPSA correspond to the conventional immunodetected total PSA (TPSA). Adenosine Deaminase (ADA) is expressed intracellular by all cells. It is also associated with the cell surface multifunctional glycoprotein CD₂₆/dipeptidyl peptidase IV. CD₂₆ is expressed on the surface of human prostate cancer 1-LN cells acting as a receptor for plasminogen (pg). Since ADA and pg bind CD₂₆ as distinct nearby sites. A possible interaction is found between these two proteins on the surface of 1-LN cells. Thus prostate specific antigen (PSA) Adenosine Deaminase (ADA) and Plasminogen (pg) are the factors regulating events in prostate cancer cells that occurs when Plasminogen (pg) binds to the cell surface and are activated in prostate cancer.

Methods:

Men with chronic kidney disease (CKD) undergone medical examination at the department of urology Princess Esra Hospital a teaching hospital to Deccan College of Medical Sciences, Hyderabad, Telangana State, India. Blood samples for PSA, ADA and plasminogen analysis were collected at the same time as the routine determination of

glomerular filtration rates (GFRs). The blood samples were allowed to clot for 15 mins at room temperature subsequently centrifuged at 3500 rpm for 10 mins and then immediately stored at -70°C for pending analysis. Measurement of serum creatinine, t-PSA, f-PSA, ADA and plasminogen were carried in all blood samples. Fluoroimmunoassay technique was used to measure t-PSA and f-PSA, from which percent of f-PSA was calculated. A quantitative estimation of creatinine was determined by enzymatic method using the enzymes creatinine amidohydrolase and creatinine amidohydrolase. The reference range was 0.5-1.5md/dl. The enzymatic activity of ADA was determined by phosphate buffer pH 7.4 containing 100 mm adenosine. ADA activity was expressed as a 265 nm/min. Binding of ADA to pg attached to CD26 and inhibition of its binding to pg by pg structures modules. Increasing concentration of pg was added to CD26 with a single concentration of ADA. It is suggested that ADA facilitates pg activation. It is only the stimulation of pg activation by ADA. When pg is bound to 1-LN cells increase the concentration of ADA which was added to 1-LN cells.

Results:

The characterisation study of patients and controls are present in table 1. In the current study evaluated PSA levels and percent of f-PSA is 201 men with chronic kidney disease (CKD). The majority of patients had moderated to severe renal insufficiency 96 % has GFR < 59 ml/min/1.73m². The median GFR was 23ml/min/1.73m².

There was no significant difference in t-PSA in the study patients compared with controls (median 1.08 ug/L versus 0.95 ug/L ; P=0.15).

Both median f-PSA and PSA, how were significantly higher ($p=0.01$) in the study patients (0.45 ug/L and 47.8 % respectively) than in the controls (0.29 ug/L and 29.9 % respectively). Among the study patients, 69(68 %) has percent f-PSA > 40 % and 39(39 %) has percent f-PSA>50 %.

Hence there is association between GFR and percent f PSA. Regression analysis showed significant association of lower GFR with higher percent Fpsa ($p=0.036$). Both pg2 and ADA bind to CD26 and ADA binds directly to pg2, the capacity of pg2 to influence the binding of ADA to CD26, increasing concentration of pg enhance the binding of ADA to the complex CD26-pg2. Stimulation of pg activation by ADA when pg is bound to 1-LN cell or immobilized CD26 .An increase concentration of ADA were added to 1-LN cells pg2 (0.2u) at 37°C.

Discussion:

Total Prostate specific antigen (tPSA), Percent free prostate-specific antigen (fPSA) are frequently measured to discriminate of prostate cancer from benign prostate disorder. It has been confirmed a higher percent fPSA indicating a less risk of cancer. By using a cut off of < 25 %. Prostate cancer detection in men with t-PSA in the range 4-10 ug/L. Diminished renal elimination of fPSA may, therefore effect percent fPSA and an accuracy as a diagnostic marker for prostate cancer. This study shows that men with chronic kidney disease and impaired renal function have significantly higher serum levels of fPSA (median 0.45 ug/L versus 0.29ug/L) and significantly higher percent fPSA (median 45 % versus 30 %) compared to control. As many as 69% of the study patients present with a percent of fPSA > 40 %. In addition to the above study ADA binding influences the function of CD₂₆ - 1-LN cells.

The binding region for pg and ADA is CD₂₆ are in close proximally. The study suggests a novel role for the complex CD₂₆-ADA. In this centre the levels of CD26 and ADA have been evaluation in prostate cancer. Prostate cancer may be higher levels of the protease plasmin, resulting from higher levels of CD₂₆ and ADA which facilitates pg localization and activation on the surface of the cells.

TABLE 1: Characteristics of the study participants

Characteristic	Study Patients (n=201)	Controls (n =350)	p-value
Age	58 (47,68)	58(55-64)	-
GFR*(ML/min/1.73m ²)	24 (16,33)	-	-
Creatinine (ug/dl)	1.8 (0.5-1.5)	-	-
Total PSA (ug/L)	1.08(0.28,1.87)	0.95(0.01,1.55)	0.15
Free PSA (ug/L)	0.45(0.28,0.76)	0.29(0.20,0.43)	<0.001
Percent free PSA	47.2(37.5,55.1)	29.9(22.4,38.8)	<0.001
Renal Diseases, No.of patients (%)			
Diabetic Nephropathy	25(25%)	-	-
Renal Transplants	20(20%)	-	-
Glomerulonephritis	22(22%)	-	-
RenalDisease (non-specific)	15(15%)	-	-
Nephrosclerosis	16(16%)	-	-
Miscellaneous renal disease	2(2%)	-	-

*GFR measured by iohexol clearance. P-values were obtained by linear regression, with adjustment for age.

Fig.1




Fig.2




Fig.3




Table 2. Adenosine Deaminase,(ADA) CD₂₆ (1-LNCCells) and Plasminogen(Pg) In patients and controls of prostate cancer.

	PATIENTS	CONTROL	P-VALUES
Adenosine Deaminase (ADA),	40±10	8±3	1.7
CD ₂₆ (1-LN Cells)	0.8±0.2	0.2±0.1	<0.001
Plasminogen (pg),	1.4±0.3	0.2±0.1	<0.001

References:

- Stenman UH, Leinonen J, Alfthan H, et al. A complex between prostate-specific antigen and alpha 1-antichymotrypsin is the major form of prostate-specific antigen in serum of patients with prostatic cancer: assay of the complex improves clinical sensitivity for cancer. *Cancer Res.* 1991;51:222-226.
- Christensson A, Laurell CB, Lilja H. Enzymatic activity of Prostate-specific antigen and its reactions with extracellular serine proteinase inhibitors. *Eur J Biochem.* 1990;194:755-763.]
- Lilja H, Christensson A, Dahlen U, et al. Prostate-specific antigen in serum occurs predominantly in complex with alpha 1-antichymotrypsin. *Clm Chem.* 1991;37:1618-1625.]
- Christensson A, Bjork T, Nilsson O, et al. Serum prostate specific antigen complexed to alpha 1-antichymotrypsin as an indicator of prostate cancer. *J Urol.* 1993;150:100-105.]
- Roddam AW, Duffy MJ, Hamdy FC, et al. Use of prostate-specific antigen (PSA) isoforms for the detection of prostate cancer in men with a PSA level of 2-10 ng/ml: systematic review and meta-analysis. *Eur Urol.* 2005;48:386-399. discussion 398-389.
- Partin AW, Piantadosi S, Subong EN, et al. Clearance rate of serum-free and total PSA following radical retroper prostatectomy. *Prostate Suppl.* 1996;7:35-39.
- Bjork T, Ljungberg B, Piironen T, et al. Rapid exponential elimination of free prostate-specific antigen contrasts the slow, capacity-limited elimination of PSA complexed to alpha 1-antichymotrypsin from serum. *Urology.* 1998;51:57-62.
- Lilja H, Haese A, Bjork T, et al. Significance and metabolism of complexed and noncomplexed prostate specific antigen forms, and human glandular kallikrein 2 in clinically localized prostate cancer before and after radical prostatectomy. *J Urol.* 1999;162:2029-2034. discussion 2034-2025.
- Martin BJ, Cheli C, Davis R, et al. cPSA and fPSA elimination in African-American men. *Prostate Cancer Prostatic Dis.* 2003;6:163-168.
- Gregorakis AK, Malourovvas D, Stefanakis S, et al. Free/Total PSA (F/T ratio) kinetics in patients with clinically localized prostate cancer undergoing radical prostatectomy. *Clin Chim Acta.* 2005;357:196-201.
- Kilic S, Yalcinkaya S, Guntekin E, et al. Determination of the site of metabolism of total, free, and complexed prostate-specific antigen. *Urology.* 1998;52:470-473.
- Birkenmeier G, Struck F, Gebhardt R. Clearance mechanism of prostate specific antigen and its complexes with alpha2-macroglobulin and alpha1-antichymotrypsin. *J Urol.* 1999;162:897-901.
- Bruun L, Ekberg H, Bjork T, et al. Rapid elimination by glomerular filtration of free prostate specific antigen and human kallikrein 2 after renal transplantation. *J Urol.* 2004;171:1432-1435.
- Fischer K, Hamza A, Wicht A, et al. Shift of the f/t-PSA quotient in relation to renal insufficiency: consequences for the early detection of prostate carcinoma in patients with terminal renal failure. *Anticancer Res.* 2007;27:1945-1948.
- Kamali K, Zargar MA. Effects of renal transplantation on serum-free and total PSA levels. *Transplant Proc.* 2007;39:1027-1028.
- Sasagawa I, Kubota Y, Hayami S, et al. Serum levels of total and free prostate specific antigen in men on hemodialysis. *J Urol.* 1998;160:83-85.
- Douville P, Tiberi M. Effect of terminal renal failure on the ratio of free to total prostate-specific antigen. *Tumour Biol.* 1998;19:113-117.
- Cristalli, G., Costanzi, S., Lambertucci, C., Lupidi, G., Vittori, S., Volpini, R., and Camaioni, E. (2001) *Med. Res. Rev.* 21, 105-128.
- Hershfield, M. S., and Mitchell, B. S. (2001) in *The Metabolic and Molecular Basis of Inherited Diseases* (Scriver, C. R., Beaudet, A. L., Sly, W. S., and Valle, D., eds) 8th Ed., pp 2585-2625, McGraw-Hill Inc., New York
- Nishihara, H., Ishikawa, S., Shinkai, K., and Akedo, H. (1973) *Biochem. Biophys. Res. Commun.* 302, 429-442