



A STUDY OF PATIENTS OF TOTAL THYROIDECTOMY GIVEN CALCIUM BY ASSESSING THEIR PARATHYROID SCORE.

Surgery

Dr. Kundan Gedam	Professor in Dept. of General surgery, MGM Medical college, Kamothe, Navi Mumbai.
Dr. Manajitsinh Jadhav	3 rd year Junior resident, Dept. of General surgery, MGM Medical college, Kamothe, Navi Mumbai.
Dr. Parnasha Chemburkar	2 nd year Junior resident, Dept. of General surgery, MGM Medical college, Kamothe, Navi Mumbai
Dr. Meghna Kinjalk	1 st year Junior resident, Dept. of General surgery, MGM Medical college, Kamothe, Navi Mumbai
Dr. Atul K. Gowardhan	Assistant Prof. in General Surgery, MGM Medical College, Kamothe, Navi Mumbai.

ABSTRACT

In this study, we scored the dis-colouration of the parathyroid glands using a new parathyroid scoring system and evaluated the correlation between the parathyroid score and duration of required calcium supplementation after total thyroidectomy.

Materials and Methods: A total of 316 patients undergoing total thyroidectomy between November 2016 and April 2017 were enrolled in this retrospective study. Parathyroid scoring was performed by one experienced surgeon..

Results: Parathyroid scores were positively correlated with ionized PTH concentrations at 2 hours ($r = 0.053$, $p < 0.001$), 2 weeks ($r = 0.056$, $p < 0.001$), 3 months ($r = 0.032$, $p < 0.001$), 6 months ($r = 0.072$, $p < 0.001$), and 1 year ($r = 0.071$, $p < 0.001$) after thyroidectomy. Parathyroid scores were significantly and inversely associated with the duration of required calcium supplementation ($p = 0.001$).

Conclusions: Parathyroid scores at the end of surgery might be helpful for predicting the degree of postoperative hypocalcemia after total thyroidectomy.

KEYWORDS

thyroidectomy, parathyroid

Introduction:

Postoperative hypoparathyroidism is the most common complication after total thyroidectomy, owing to unintentional injury or decreased blood flow to the parathyroid glands. Because the thyroid glands and parathyroid glands play very important roles in maintaining homeostasis in the body, they should remain intact. However, when total thyroidectomy is necessary because of thyroid cancer, it is important to completely remove the malignant portion while avoiding damage to the remaining portion. Hypoparathyroidism is a major cause of hypocalcemia, lengthens hospital stays, and necessitates repetitive laboratory tests and continued administration of both calcium and vitamin D [1, 2, 3].

Although the diverse causes of hypocalcemia include excessive discharge of calcium into urine due to surgical stress, hungry bone syndrome due to hyperthyroidism or hypercalcitoninemia, and vitamin D deficiency, the most important causes include physical damage to the four parathyroid glands, which are adjacent to the thyroid glands, during a total thyroidectomy; disturbance of the blood supply to the parathyroid glands resulting from damage to the blood vessels; and removal of the parathyroid glands by unintentional resection [4,5]. Therefore, an anatomical understanding of the thyroid glands and parathyroid glands in addition to meticulous surgical techniques are required to prevent these causes of hypoparathyroidism; although specialized surgeons make fewer mistakes with regard to damage to the glands, they do still occur [6,7].

After a total thyroidectomy, hypocalcemia reportedly occurs in approximately 10–50% of patients, and approximately 0.5–2% of these patients have permanent hypocalcemia that persists for at least one year [1, 8–12]. Side effects of total thyroidectomy-induced hypocalcemia include Chvostek's sign, Trousseau's sign, numbness, and paresthesia of the fingertips; these cause discomfort for patients and reluctance to be discharged early. Long-term progressive changes can result in abnormal skeletal microstructures [13,14].

Therefore, the purpose of the present study was to categorize the condition of the parathyroid gland using a simple scoring system immediately after a total thyroidectomy and determine the correlation between post-operative intact parathyroid hormone (iPTH) levels and duration of required calcium supplementation.

Materials and methods:

A total of 316 patients undergoing total thyroidectomy with central node dissection (CND) between March 2011 and April 2017 were studied in the Surgery department of MGM medical college & Hospital, Navi Mumbai, a tertiary care teaching hospital in western Maharashtra, India. An ipsilateral CND (pretracheal and ipsilateral paratracheal node dissection) was performed routinely for prophylactic or therapeutic purposes, and a bilateral CND was performed for patients with bilateral thyroid cancer. The subjects were divided into three groups based on the duration of calcium replacement: group I did not require calcium and vitamin D supplements or required them for ≤ 2 weeks; group II required supplementation for ≤ 6 months; and group III required supplementation for ≥ 6 months (Table 1). In the patients who recovered from hypoparathyroidism, calcium and vitamin D replacement was stopped.

Results:

Group I consisted of 217 patients (68%) who did not experience transient hypoparathyroidism and 43 patients (13.5%) who required calcium and vitamin D supplementation for ≤ 2 weeks. In group III, 18 patients (5.75%) who had permanent hypoparathyroidism for at ≥ 6 months, 6 patients (1.9%) recovered from hypoparathyroidism after 1 year; of the remaining 12 patients (3.8%), 2 patients (0.6%) recovered from hypoparathyroidism in the third year after surgery.

As shown in Table 1, 3 groups were compared clinicopathological information, but only postoperative parathyroid scores were significantly different ($p < 0.001$) between the 3 groups (Table 1).

Within each group, iPTH levels decreased with increasing duration of calcium supplementation, at 2 hours, 2 weeks, 6 months, and 1 year after surgery (Table 2).

Discussion:

The scoring method (parathyroid score) using color that we developed to determine the status of the parathyroid glands after total thyroidectomy was able to represent the condition of the parathyroid well, as indicated by the association between the duration of calcium supplementation and iPTH values. Furthermore, the scores were correlated with iPTH levels. Therefore, the scores could be a good

indicator for postoperative hypocalcemia. Hypocalcemia remains one of the most frequent complications after a total thyroidectomy, in addition to postoperative hematoma and hoarseness. Hypocalcemia occurs due to unintentional damage or decreased blood flow to the parathyroid glands [4] and can cause complaints of a tingly sensation and muscle cramping. Therefore, an important part of post-thyroidectomy management involves the continuous monitoring of calcium levels and supplementation of calcium immediately after hypoparathyroidism is identified. However, it is important to avoid conventional blood tests in patients with no possibility of hypocalcemia.

In a study conducted in 2010, a comparison of discolored parathyroid glands with normal-colored parathyroid glands and autotransplanted glands indicated that discolored parathyroid glands have a transient deterioration in function [16].

In addition, the four parathyroid glands do not function evenly. Even with similar colors, the amount of parathyroid hormones secreted by individual parathyroid glands could be considerably different, and large, dominant parathyroid glands could secrete relatively large amounts of hormones. Therefore, when a dominant parathyroid has been severely damaged or excised, the risk of postoperative transient hypoparathyroidism is high, even with high parathyroid scores for the remaining glands.

Methods to treat unavoidable damage to the parathyroid glands include auto-transplantation and calcium and vitamin D supplementation. Although auto-transplantation of the parathyroid glands is reportedly effective for long-term recovery of parathyroid gland function [17,18], the post-recovery parathyroid gland hormone levels were much lower with auto-transplantation of the parathyroid glands when there was no preserved parathyroid gland compared with auto-transplantation of the parathyroid glands when there was at least one preserved parathyroid gland [19].

In our study, there were 20 cases detected where autotransplantation was performed. Among those 20 patients, 14 patients were in group 1, 6 in group 2, and no patient was observed in group 3 (data not shown). Six of the 20 patients with autotransplantation needed more than 2 weeks of medication, but after 6 months, all patients reached normocalcemia with normal intact parathyroid hormone level. Compared to this finding, 18 out of 50 patients without autotransplantation who needed more than 2 weeks of medication, medication for more than six month was needed (data not shown). Although it might not be statistically significant due to small sample size, autotransplantation helps in maintaining normal calcium level.

In conclusion, the need for and duration of calcium supplementation after total thyroidectomy were significantly related with the parathyroid score. Therefore, the parathyroid score, which is a simple method, might be effective for predicting and managing postoperative hypocalcemia, saving time and cost. These findings are important for surgeons.

Tables:

TABLE 1: Analysis of risk factors affecting the duration of calcium requirement.

	Oral medication			p value
	Group I	Group II	Group III	
Number of patients (%)	260(82.3)	38(12.0)	18(5.75)	
Age, mean	19-77(45.50)	22-65(46.34)	30-65(45.89)	0.911
Sex				0.303
Female (%)	204(78.5)	33(86.8)	16(88.9)	
Male (%)	56(21.1)	5(13.2)	2(11.1)	
Largest primary tumor diameter in cm	0.899(0.1-3) [*]	0.942(0.4-2.5) [*]	1.058(0.2-4.0) [*]	0.486
Extrathyroidal extension (%)	147(56.8)	19(51.4)	9(52.9)	0.799
Multifocality (%)	73(28.1)	15(39.5)	5(27.8)	0.516
Central nodal status (mean)				
dissected lymph nodes	0-18(5.62) [*]	1-20(6.37) [*]	1-15(7.00) [*]	0.259
metastatic lymph nodes	0-13(1.27) [*]	0-5(0.97) [*]	0-9(1.50) [*]	0.632
Thyroiditis				0.175
Yes	89(34.2)	10(27.0)	9(52.9)	
No	171(65.8)	27(73.0)	8(52.9)	
MRND				0.692
Yes	28(10.8)	5(13.2)	1(5.6)	
No	232(89.2)	33(86.8)	17(94.4)	
Parathyroid score (mean)	7.32	6.45	4.89	<0.001

Table 2: iPTH and parathyroid score according to Ca+ medication.

	Oral medication			p value
	Group I	Group II	Group III	
Pre-OP iPTH (mean)	46.35	40.96	40.27	0.075
iPTH: 2 hours (mean)	16.80	5.03	4.21	<0.001
iPTH: 2 weeks (mean)	32.47	11.02	7.89	<0.001
iPTH: 6 months (mean)	38.59	28.08	7.91	<0.001
iPTH: 1 year (mean)	32.55	24.77	12.56	<0.001

References:

- Walker Harris V, Jan de Beur. Postoperative hypoparathyroidism: Medical and surgical therapeutic options. *Thyroid*. 2009; 19: 967-973.
- Toniato A, Boschin IM, Piotta A, Pelizzo M, Sartori P. Thyroidectomy and parathyroid hormone: tracing hypocalcemia-prone patients. *Am J Surg*. 2008; 196: 285-288.
- Abboud B, Sargi Z, Akkam M, Sleilaty F. Risk factors for postthyroidectomy hypocalcemia. *J Am Coll Surg*. 2002; 195: 456-61.
- Bilezikian JP, Khan A, Potts JT Jr, Brandi ML, Clarke BL, Shoback D et al. Hypoparathyroidism in the adult: epidemiology, diagnosis, pathophysiology, target-organ involvement, treatment, and challenges for future research. *J Bone Miner Res*. 2011; 26: 2317-2337.
- Prime MP, de Diego JJ, Hardisson D, Madero R, Gavilan J. Factors related to nerve injury and hypocalcemia in thyroid gland surgery. *Otolaryngol Head and Neck*. 2001; 124: 111-114.
- McCullough M, Weber C, Leong C, Sharma J. Safety, efficacy, and cost savings of single parathyroid hormone measurement for risk stratification after thyroidectomy. *Am Surg*. 2013; 79: 768-774.
- Chow TL, Choi CY, Chiu AN. Postoperative PTH monitoring of hypocalcemia expedites discharge after thyroidectomy. *Am J Otolaryngol*. 2014; 35: 736-740.
- Asari R, Passeler C, Kaczirek K, Scheuba C, Niederle B. Hypoparathyroidism after total thyroidectomy: a prospective study. *Arch Surg*. 2008; 143: 132-138.
- McHenry CR, Speroff T, Wentworth D, Murphy T. Risk factors for postthyroidectomy hypocalcemia. *Surgery*. 1994; 116: 641-648.
- Pattou F, Combemale F, Fabre S, Carnaille B, Decoux M, Wemeau JL et al. Hypocalcemia following thyroid surgery: incidence and prediction of outcome. *World J Surg*. 1998; 22: 718-724.
- Erbil Y, Barbaros U, Temel B, Turkoglu U, Issever H, Bozboru A et al. The impact of age, vitamin D(3) level, and incidental parathyroidectomy on postoperative hypocalcemia after total or near total thyroidectomy. *Am J Surg*. 2009; 197: 439-446.
- Sywak MS, Palazzo FF, Yeh M, Wilkinson M, Snook K, Sidhu SB et al. Parathyroid hormone assay predicts hypocalcemia after total thyroidectomy. *ANZ J Surg* 2007; 77: 667-670.
- Rubin MR, Dempster DW, Zhou H, Shane E, Nickolas T, Sliney J Jr et al. Human chorionic gonadotropin measurements in parathyroid carcinoma. *Eur J Endocrinol*. 2008; 159: 469-474.
- Rubin MR, Bilezikian JP. Hypoparathyroidism: Clinical features, skeletal microstructure and parathyroid hormone replacement. *Arq Bras Endocrinol Metabol*. 2010; 54: 220-226.
- Payne RB, Little AJ, Williams RB, Milner JR. Interpretation of serum calcium in patients with abnormal serum proteins. *Br Med J*. 1973; 4: 643-646.
- Promberger R, Ott J, Kober F, Mikola B, Karik M, Freissmuth M et al. Intra- and postoperative parathyroid hormone-kinetics do not advocate for autotransplantation of discolored parathyroid glands during thyroidectomy. *Thyroid*. 2010; 20: 1371-1375.
- Lo CY. Parathyroid autotransplantation during thyroidectomy. *ANZ J Surg* 2002; 72: 902-907.
- Cavallaro G, Iorio O, Centanni M, Porta N, Iossa A, Gargano L et al. Parathyroid Reimplantation in Forearm subcutaneous Tissue During Thyroidectomy: A Simple and Effective Way to Avoid Hypoparathyroidism. *World J Surg*. 2015; 39: 1936-1942.
- Kihara M, Miyauchi A, Kontani K, Yamauchi A, Yokomise H. Recovery of parathyroid function after total thyroidectomy. *Head Neck* 2003; 25: 799-807.
- Gupta S, Chaudhary P, Durga CK, Naskar D. Validation of intra-operative parathyroid hormone and its decline as early predictors of hypoparathyroidism after total thyroidectomy: A prospective cohort study. *Int J Surg*. 2015; 18: 150-153.