

EFFECT OF LOW DOSE MIDAZOLAM WITH LOW DOSE DEXMEDETOMIDINE ON BRADYCARDIA AND SEDATION AFTER SPINAL BLOCK: A PROSPECTIVE CASE CONTROL STUDY

Anesthesiology

Dr. Kashyap Jha Manju	Professor and HOD Department of Anaesthesiology and Critical Care, Pravara Institute of Medical Sciences, Loni.
Dr. Prashant Rai*	MD Scholar Department of Anaesthesiology and Critical Care, Pravara Institute of Medical Sciences, Loni. *Corresponding Author
Dr. Bhumika Rangparia	MD Scholar Department of Anaesthesiology and Critical Care, Pravara Institute of Medical Sciences, Loni.

ABSTRACT

Introduction: Sedation makes it convenient for the patient, the anaesthesiologist and the surgeon during surgeries under regional anaesthesia. Midazolam in doses of 0.05mg/Kg produces good sedation and excellent amnesia but has no specific analgesic properties and causes depression of upper airway muscle tone in the elderly, resulting in a higher incidence of airway obstruction. Midazolam is commonly used drug for sedation during regional anaesthesia, which produces good sedation and excellent amnesia but causes upper airway obstruction and respiratory depression. Dexmedetomidine has been used as a sedative for various surgical and medical procedures, but loading dose of 1µg/kg has been associated with higher incidences of bradycardia, whereas lower doses produce inadequate sedation.

Aim: To evaluate the effect of low dose midazolam on bradycardia and sedation during low dose dexmedetomidine infusion after spinal block.

Methodology: In this prospective case control study, 80 patients undergoing surgeries under spinal anaesthesia were studied under two groups, Group D (n=40) who received 1mcg/kg dexmedetomidine as loading dose and Group DM (n=40) who received inj. midazolam 0.025mg/kg along with 0.5mcg/kg dexmedetomidine as loading dose. Both the groups were maintained by continuous infusion of dexmedetomidine at 0.04mcg/kg/hr. All patients were monitored for heart rate, mean arterial pressure, respiratory rate, oxygen saturation, Observer's Assessment of Alertness/Sedation Scale (OAA/S) and complications.

Results: Mean heart rate was significantly lower in group D compared to group DM after 5 and 10 min of loading ($P < 0.0001$). Incidence of bradycardia requiring Atropine was significantly higher ($P = 0.0011$) in group D (22/40) than group DM (7/40). OAA/S score was significantly lower in group DM at 5 minutes ($P=0.0049$) and 10 minutes ($P=0.0004$). Other hemodynamic variables were comparable in both groups.

Conclusion: Low dose Midazolam with low dose Dexmedetomidine offered better results for bradycardia and sedation compared to routine dose of Dexmedetomidine alone.

KEYWORDS

Dexmedetomidine, Midazolam, bradycardia, sedation, Spinal block

Introduction:

Regional anaesthesia has many advantages like maintaining spontaneous breathing, relaxing necessary muscles for surgery, providing postoperative analgesia, avoiding the risks of intubation and pulmonary aspiration [1]. Staying awake, early family contact, and early food intake are major advantages from patients point of view [2]. However, the fear of surgery complemented with strange surroundings of the operation theatre, sophisticated machines and masked faces of strange personals makes the patient uncomfortable and fearful [3, 4]. Continuous supine position for a prolonged duration and inability to move the body during regional anaesthesia further increases the discomfort causing many patients to panic. [5]. Sedation has been shown to increase patient satisfaction and acceptance of regional anaesthesia [6, 7]. Sedation makes it convenient for the patient, the anaesthesiologist and the surgeon during surgeries under regional anaesthesia.

The goals of sedation during regional anesthesia include rapid achievement of adequate sedation, its maintenance at a constant level during the surgical procedure and awakening the patient quickly at the end. Many techniques and drug regimens have been tried for providing adequate sedation during regional anaesthesia [8, 9, 10]. Continuous infusions have been proven to produce lesser side effects, faster recovery, easy controllability over the desired depth of sedation and, should the regional block prove to be ineffective, easy conversion to general anesthesia [11, 12].

Midazolam has a fast onset and short recovery time, because of which it is one of the most widely used sedatives in spinal anesthesia. Midazolam in doses of 0.05mg/Kg produces good sedation and excellent amnesia but has no specific analgesic properties and causes depression of upper airway muscle tone in the elderly, resulting in a higher incidence of airway obstruction [13, 14, 15].

Dexmedetomidine, Alpha 2 -adrenergic agonist has both analgesic and sedative properties, when used as an adjuvant to regional anesthesia [16, 17]. They potentiate the effect of local anesthetics and prolong the duration of both motor, sensory spinal blockade and postoperative

analgesia [18, 19, 20]. Other claimed advantages include minimal respiratory depression with cardio-protection, neuroprotection and renoprotection [21]. Hence dexmedetomidine has been used as a sedative for various purposes, including ICU sedation, awake fiberoptic intubation, various surgical and medical procedures. In general practice dexmedetomidine needs to be given as an initial loading dose of up to 1µg/kg followed by a continuous infusion. But during the initial loading period or too rapid administration can lead to bradycardia due to sympatholytic effects [22-27], and using dose lesser than this cause inadequate sedation. Study by Yun-Sic Bang et al [28]. suggests adding low dose Midazolam and reducing Dexmedetomidine loading dose reduce chances of bradycardia while maintaining adequate sedation. However, no other study has been conducted on this subject. The present study is to evaluate the effect of low dose midazolam with low dose dexmedetomidine on bradycardia and sedation during dexmedetomidine infusion after spinal block.

Aims and objectives:

Aim of this study is to evaluate effects of low dose midazolam on bradycardia and sedation during dexmedetomidine infusion.

Methodology:

This double blinded, prospective case control study was conducted after obtaining the institutional ethical committee clearance & written informed consent from the patient. 80 patients of either gender between 18 to 60 years of age, belonging to ASA grade I & II, undergoing surgeries under spinal anaesthesia at P.R.H. Loni, willing for study and sedation after spinal block were included for the study. Patients with known hypersensitivity to study drugs, on β -blockers or anti-coagulants, with psychiatric disorder, third degree heart block, hypotension (mean arterial pressure < 70 mmHg), bradycardia (baseline heart rate < 60 beats/min), pregnant or breast feeding female were not included in the study. All patients were pre-medicated tab. Ranitidine 150mg orally on the night before surgery and kept nil orally from midnight. On the day of surgery intravenous access secured with 18-gauge venous cannula and preloading with 10 ml/ kg of lactated Ringer's solution was done.

After noting the base line readings, patients were administered spinal

block under aseptic precautions using 27G Quincke's needle, in left lateral position at L₃-L₄ level with hyperbaric Inj. Bupivacaine 13mg + 25 micrograms Inj. Fentanyl. Patient developing bradycardia (HR <50 bpm) or hypotension (MAP <60 mm Hg or fall in MAP more than 20%) after spinal block were managed and omitted from the study. After the confirmation of successful spinal anesthesia with adequate level of blockade, sedation was administered. Patients were studied under two groups depending on the type of sedation received. Group DM (n = 40)- received Inj. Midazolam 0.025mg/kg (diluted up to 4cc in normal saline) intravenously and Inj. Dexmedetomidine 0.5µg/kg intravenously over a period of 10 min as loading dose followed by continuous infusion of 0.4µg/kg/hr. Group D (n = 40)- received 4cc of normal saline intravenously and Inj. Dexmedetomidine 1µg/kg intravenously over a period of 10 min as loading dose followed by continuous infusion of 0.4µg/kg/hr. Study drugs were prepared by anesthesiologist not involved in data recording, thus patient and investigator were blinded from the study drugs. All patients received oxygen at 5L/min via face mask and intravenous Ringers lactate solution for maintenance. Observer's Assessment of Alertness/Sedation Scale (OAA/S) was used for assessment of sedation level.

Table 1: Observer's assessment of alertness/sedation (OAA/S) scale

Observation	Score level
Responds readily to name spoken in normal tone	5
Lethargic response to name spoken in normal tone	4
Responds only after name is called loudly and/or repeatedly	3
Responds only after mild prodding or shaking	2
Does not respond to mild prodding or shaking	1

Heart rate (HR), Mean arterial pressure (MAP), respiratory rate (RR), oxygen saturation (SpO₂) and the Observer's Assessment of Alertness/Sedation Scale (OAA/S) were recorded at baseline (T1), after spinal block (T2), before dexmedetomidine administration (T3), after dexmedetomidine administration every 5mins till first 20 min (T5, T10, T15, T20) and every 20 min till 80 min (T40, T60, T80). Inadequate sedation (OAA/S > 3) after 15 minutes of starting infusion were managed by rescue midazolam (0.5mg intravenously) and repeated whenever required to maintain OAA/S < 3. Side effects at any point during the surgery were documented and managed. In case of bradycardia (HR <50 bpm) Inj. Atropine Sulphate 0.5mg was given intravenously, for hypotension (MAP <60 mm Hg or fall in MAP more than 20%) Inj. Ephedrine HCl 5 mg in incremental doses were given, for respiratory depression (rate <8 or SpO₂ <90%) the depth and rate of respiration was increased with painful stimuli, airway insertion and stopping the infusion. Nausea/Vomiting by Inj. Ondansetron 4mg intravenously. The infusion was discontinued after completion of surgery and patients were shifted to post-anaesthesia care unit (PACU) till OAA/S = 5 was achieved. MAP, HR, SpO₂, OAA/S were noted on arrival and at discharge from PACU. Time taken for OAA/S score to reach 5 after stopping the study drug was noted. Patients were observed for any complications postoperatively for 24 hrs.

Statistical analysis:

All data were entered into a proforma in excel sheet (MS Office 2016) and statistical analysis was done using SPSS software (Statistical Package for Social Science [SPSS] version 19.0 for Windows, SPSS, Inc). Chi square test, students unpaired t test was used for inter group comparisons and repeated measures ANOVA for within the group comparisons. P value < 0.05 was considered statistically significant.

Observations and Results:

Effect of spinal anaesthesia was adequate in all the patients and none required administration of general anaesthesia, none of the patients were excluded from the study after spinal block and thus all 80 patients were considered for evaluation process (40 in each group).

Table 2: Demographic profile


	Group D	Group DM	Statistical test	P Value
Age (year)	35.88 ± 11.5	38.4 ± 8.97	Unpaired t test	0.277
Gender (M/F)	23/17	19/21	Chi square test	0.502
Weight (Kg)	66.78 ± 10.34	63.35 ± 10.98	Unpaired t test	0.1549
Height (cm)	163.13 ± 9.51	160.45 ± 9.18	Unpaired t test	0.2046
ASA grading (I/II)	21/19	18/22	Chi square test	0.655

Both groups were comparable in terms of age, gender, weight, height and ASA grading (p>0.05).


Table 3: Mean HR (Intraoperative)

TIME INTERVAL	GROUP D		GROUP DM		P
	MEAN	S.D.	MEAN	S.D.	VALUE
T1	78.23	7.96	80.05	7.14	0.2838
T2	81.50	8.31	83.10	7.20	0.3601
T3	79.60	8.71	81.03	7.03	0.4232
T5	59.25	8.09	67.10	6.98	<0.0001
T10	63.93	10.42	65.30	5.66	<0.0001
T15	66.35	7.86	64.35	5.52	0.1919
T20	64.78	6.73	64.83	4.33	0.9686
T40	65.25	5.79	66.35	4.55	0.3479
T60	66.28	5.80	67.40	4.72	0.3444
T80	67.33	5.97	68.25	4.49	0.4395

Graph 1: Intraoperative mean HR



Graph 2: Patients requiring Atropine




Significant fall in mean heart rate was observed in both groups (P < 0.05) after 5 min of dexmedetomidine loading dose. Mean heart rate was significantly lower in group D compared to group DM at 5 and 10 min of loading (P < 0.0001). 22 patients (55%) in group D required Atropine intraoperatively as compared to 7 patients (17.50%) in group DM and the difference was statistically significant (P = 0.0011).

Table 4: Intraoperative mean MAP

TIME INTERVAL	GROUP D		GROUP DM		P VALUE
	MEAN	S.D.	MEAN	S.D.	
T1	91.28	11.60	92.40	10.00	0.6435
T2	85.40	11.82	85.05	9.47	0.8842
T3	82.30	11.77	83.15	9.53	0.7236
T5	80.45	12.21	80.90	9.64	0.8553
T10	79.30	11.81	79.68	9.75	0.9105
T15	78.65	11.15	79.65	8.99	0.6599
T20	78.00	10.87	79.03	8.75	0.6436
T40	78.95	10.92	80.48	9.08	0.4991
T60	79.68	10.87	81.40	9.09	0.4438
T80	81.05	11.34	82.68	9.17	0.4830

Graph 3: Intraoperative mean MAP



Significant fall in mean MAP was observed in both groups ($P < 0.05$) after spinal block and 5 min of dexmedetomidine loading dose. Fall in both groups were comparable at all-time intervals.

Table 5: Intraoperative mean RR

TIME INTERVAL	GROUP D		GROUP DM		P VALUE
	MEAN	S.D.	MEAN	S.D.	
T1	16.03	2.81	15.73	2.18	0.5958
T2	17.00	2.32	16.63	2.22	0.3525
T3	16.58	2.36	15.98	2.15	0.2045
T5	16.25	2.17	15.73	2.22	0.2326
T10	16.75	2.37	16.40	2.46	0.3662
T15	16.98	2.56	16.03	2.63	0.1052
T20	16.28	2.01	15.38	2.72	0.0968
T40	15.68	2.29	15.15	2.67	0.3478
T60	15.08	2.48	14.50	2.80	0.3038
T80	15.40	2.57	14.93	3.17	0.3890

Graph 4: Intraoperative mean RR





Table 6: Intraoperative mean SpO2

	GROUP D		GROUP DM		P VALUE
	MEAN	S.D.	MEAN	S.D.	
T1	98.88	0.79	98.85	0.77	0.8864
T2	98.98	0.80	98.73	0.85	0.1787
T3	99.03	0.80	99.13	0.88	0.5309
T5	98.88	0.88	99.23	0.77	0.0621
T10	99.08	0.83	99.08	0.92	0.999
T15	99.23	0.77	99.18	0.87	0.2719
T20	99.08	0.83	98.88	0.79	0.2728
T40	98.75	0.71	99.05	0.85	0.0892
T60	99.15	0.74	99.30	0.76	0.3718
T80	99.08	0.80	99.03	0.86	0.7884

Graph 5: Intraoperative mean SpO2




Mean Respiratory Rate and mean SpO2 were comparable in both the groups at all-time intervals and no respiratory depression was observed in either group.

Table 7: Mean OAA/S score (Intraoperative)

TIME INTERVAL	GROUP D		GROUP DM		P VALUE
	MEAN	S.D.	MEAN	S.D.	
T1	5.00	0.00	5.00	0.00	--
T2	5.00	0.00	5.00	0.00	--
T3	5.00	0.00	5.00	0.00	--
T5	4.43	0.71	3.93	0.83	0.0049
T10	3.48	0.75	2.83	0.81	0.0004
T15	2.58	0.64	2.48	0.78	0.5328

T20	2.40	0.50	2.38	0.67	0.8497
T40	2.25	0.49	2.28	0.55	0.8318
T60	2.33	0.53	2.35	0.48	0.8253
T80	2.48	0.55	2.53	0.60	0.6993

Graph 6: Mean OAA/S



Significant fall in mean OAA score was observed in both groups ($P < 0.05$) after 5 min of dexmedetomidine loading dose. Mean OAA/S score was significantly lower in group DM at 5 minutes ($P=0.0049$) and 10 minutes ($P=0.0004$). However, it remained comparable in both groups during rest of the intraoperative period. 2 patients in group DM and 3 in group D required midazolam supplementation to maintain adequate sedation, however the difference was insignificant ($P = 0.6442$).

Graph 7: Patients requiring rescue midazolam





Table 8: Post-operative monitoring

	GROUP D		GROUP DM		P
	MEAN	SD	MEAN	SD	VALUE
HEART RATE					
At Arrival	70.43	6.19	71.38	5.18	0.4591
At Discharge	70.88	6.33	72.43	5.27	0.2374
MEAN ARTERIAL PRESSURE					
At Arrival	83.20	11.29	83.73	9.22	0.8203
At Discharge	84.08	11.28	84.18	9.22	0.9665
RESPIRATORY RATE					
At Arrival	15.55	2.36	15.23	2.31	0.5362
At Discharge	16.00	2.39	15.50	2.36	0.3494
SPO2					
At Arrival	99.05	0.78	99.15	0.86	0.5891
At Discharge	99.13	0.82	99.18	0.81	0.7852
OAA/S SCORE					
At Arrival	3.25	0.74	3.10	0.78	0.3804
At Discharge	5.00	0.00	5.00	0.00	--
RECOVERY TIME					
Time taken for OAA/S to reach 5	39.05	19.07	34.13	17.21	0.2291

Graph 8: Mean recovery time



All hemodynamic variables were comparable in both groups during post-operative period. Mean recovery time after stopping dexmedetomidine infusion was 39.05 ± 19.07 minutes in group D and 34.13 ± 17.21 minutes in group DM with no significant difference ($P = 0.229$).

No other complication was reported in any patient during intraoperative and postoperative period.

Discussion:

In our study both the groups were comparable for age, sex, height, weight, duration of surgery.

Dexmedetomidine provides excellent sedation without respiratory depression and also has added advantage of prolonged duration of motor & sensory blockade and postoperative analgesia after spinal block [18, 19, 20].

Multiple studies have reported higher incidence of bradycardia (more than 40%) during initial loading with dexmedetomidine due to sympathetic blockade [22 to 27]. It has been suggested that dexmedetomidine has a dose-dependent effect on hemodynamic profile which may be altered by the loading dose [29, 30]. In our study we used dose of dexmedetomidine to reduce the incidence of bradycardia.

We observed, mean heart rate was significantly lower in group D compared to group DM at 5 and 10 min of loading ($P < 0.0001$). Significantly lower incidence ($P = 0.0011$) of bradycardia was observed in group DM with 7 patients (17.50%) compared to 22 patients (55%) in group D required atropine intraoperatively. Similar results were observed by Yun-Sic-Bang et al. who found patients requiring atropine as 15/33 in group receiving $1\mu\text{g/kg}$ vs. 5/32 in group receiving $0.5\mu\text{g/kg}$ dexmedetomidine, $P = 0.009$ [28]. Mizrak et al. also observed no significant bradycardia with loading dose of $0.5\mu\text{g/kg}$ before general anaesthesia [31].

The incidence of bradycardia was significantly lower in group DM compared to the group D probably because the sympatholytic action and vagal mimetic effect of dexmedetomidine is reduced due to the low dose.

Loading with dexmedetomidine in dose of $1\mu\text{g/kg}$ provides faster and better sedation compared to $0.5\mu\text{g/kg}$ [29, 30]. There have been reports of the synergistic enhancement of their sedative effects when midazolam and dexmedetomidine are used in combination [32]. In our study we used low dose midazolam 0.025mg/kg as adjunct to low dose $0.5\mu\text{g/kg}$ dexmedetomidine for loading in group DM.

Using OAA/S scale to assess sedation, we observed mean OAA/S score was significantly lower in group DM at 5 minutes ($P=0.0049$) and 10 minutes ($P=0.0004$). Thus faster and better onset in group receiving midazolam and dexmedetomidine. OAA/S was comparable in both groups during rest of the intraoperative period and number of patients requiring rescue midazolam in both the groups were comparable. Similar results were observed by Yun-Sic-Bang et al. who found significantly lower BIs and OAA/S at 5 and 10 min after loading in group receiving $0.5\mu\text{g/kg}$ dexmedetomidine.

Dexmedetomidine being a α_2 -adrenergic agonist causes an α_2 -adrenoceptor induced vasoconstrictive response in the peripheral vasculature which increases the blood pressure initially. Then due to both centrally and peripherally mediated sympatholytic action, hypotension follows [33]. This decrease in the sympathetic outflow and circulating catecholamine levels [25-27] as well as the vagal mimetic effect [34] of dexmedetomidine cause a decrease in the HR and BP.

In our study we observed Significant fall in mean MAP in both groups ($P < 0.05$) after spinal block and 5 min of dexmedetomidine loading dose. Fall in both groups were comparable at all-time intervals.

Respiratory depression was not observed in any of the group, probably due to use of low dose of midazolam (0.025mg/kg).

Hemodynamic variables and mean recovery time were comparable among both the groups postoperatively. No other side effects were observed in any of the group.

These findings suggest that the earlier sedation effect after the start of dexmedetomidine was achieved as a result of the administration of low dose midazolam until later sedation was achieved by dexmedetomidine. Also, both dexmedetomidine and midazolam complemented each other intraoperatively, enabling an optimal level of sedation to be achieved and maintained. More time was required to achieve optimal sedation with dexmedetomidine alone.

Conclusion:

Midazolam 0.025 mg/kg with a halved loading dose ($0.5\mu\text{g/kg}$) of dexmedetomidine was superior in terms of producing sedation and lower incidence of bradycardia than with dexmedetomidine alone ($1\mu\text{g/kg}$), without causing respiratory depression or hemodynamic instability.

References:

- Hwoe G, Seung HB, Seong WB, Hae KK, Sang WS, Kyung HK. Optimal dose of dexmedetomidine for sedation during spinal anesthesia, Korean J Anesthesiol 2013 May 64(5): 426-431
- De Andres J, Valia JC, Gil A, Bolinches R. Predictors of patient satisfaction with regional anaesthesia. Reg Anesth 1995;20:498-505.
- Aeshnourne K, Albaladejo P, Smail N, Heriche C, Sitbon P, Guenon JP, et al. Information and anaesthesia: What does the patient desire? Ann Fr Anesth Reanim 2000;19:577-81.
- Badner NH, Nielson WR, Munk S, Kwiatkowska C, Gelb AW. Preoperative anxiety: Detection and contributing factors. Can J Anaesth 1990;37:444-7.
- Schneider TW, Minto CF. Predictors of onset and offset of drug effect. Eur J Anaesthesiol 2001;23:26-31.
- Wu CL, Naqibuddin M, Fleisher LA. Measurement of patient satisfaction as an outcome of regional anesthesia and analgesia: a systematic review. Reg Anesth Pain Med 2001;26:196-208.
- Practice guidelines for sedation and analgesia by non-anesthesiologists. American Society of Anesthesiologists Task Force on Sedation and Analgesia by Non-Anesthesiologists. Anesthesiology 2002;96:1004-17.
- Höhener D, Blumenthal S, Borgat A. Sedation and regional anaesthesia in the adult patient. Br J Anaesth 2008;100:8-16.
- Helgeson LE. Sedation during regional anaesthesia: Inhalational versus intravenous. Curr Opin Anaesthesiol 2005;18:534-9.
- Nemethy M, Paroli L, Williams-Russo PC, Blanck TJ. Assessing sedation with regional anaesthesia: Inter-rater agreement on a modified Wilson Sedation Scale. Anesth Analg 2002;94:723-8.
- Smith I, Monk TG, White PF, Ding Y. Propofol infusion during regional anesthesia: sedative, amnestic, and anxiolytic properties. Anesth Analg 1994;79:313-19. 136 Topics in Spinal Anaesthesia
- Casati A, Fanelli G, Casaletti E, et al. Clinical assessment of target-controlled infusion of propofol during monitored anaesthesia care. Can J Anaesth 1999;46:235-39
- De Andrés J, Bolinches R. Comparative study of propofol and midazolam for sedation in regional anesthesia. Rev Esp Anesthesiol Reanim 1993;40:354-59.
- Leiter JC, Knuth SL, Krol RC, Bartlett D Jr. The effect of diazepam on genioglossal muscle activity in normal human subjects. Am Rev Respir Dis 1985;132:216-19.
- Montravers P, Dureuil B, Desmonts JM. Effects of i.v. midazolam on upper airway resistance. Br J Anaesth 1992;68:27-31.
- Gabriel JS, Gordin V. Alpha 2 agonists in regional anaesthesia and analgesia. Curr Opin Anaesthesiol 2001;14:751-3. [PUBMED]
- Mauro VA, Brandão ST. Clonidine and dexmedetomidine through epidural route for post-operative analgesia and sedation in a cholecystectomy. Rev Bras Anesthesiol 2004;4:1-10.
- Streibel S, Gutzler JA, Schneider MC, Aeschbach A, Kindler CH. Small-dose intrathecal clonidine and isobaric bupivacaine for orthopedic surgery: A dose-response study. Anesth Analg 2004;99:1231-8. [PUBMED]
- Dobrydnyov I, Axelsson K, Thorn SE, Matthiesen P, Klockhoff H, Holmstrom B, et al. Clonidine combined with small-dose bupivacaine during spinal anesthesia for inguinal herniorrhaphy: A randomized double-blinded study. Anesth Analg 2003;96:1496-503.
- Sethi BS, Samuel M, Sreevastava D. Efficacy of analgesic effects of low dose intrathecal clonidine as adjunct to bupivacaine. Indian J Anaesth 2007;51:415.
- Panzer O, Moitra V, Sladen RN. Pharmacology of sedative-analgesic agents: Dexmedetomidine, remifentanyl, ketamine, volatile anesthetics, and the role of peripheral mu antagonists. Crit Care Clin 2009;25:451-69.
- Grant SA, Breslin DS, MacLeod DB, Gleason D and Martin G. Dexmedetomidine infusion for sedation during fiberoptic intubation: a report of three cases. J Clin Anesth 2004; 16: 124- 126.
- Riker RR, Shehabi Y, Bokesch PM, Ceraso D, Wisemandle W, Koura F, Whitten P, Margolis BD, Byrne DW, Ely EW, Rocha MG; SEDCOM (Safety and Efficacy of Dexmedetomidine Compared With Midazolam) Study Group. Dexmedetomidine vs midazolam for sedation of critically ill patients: a randomized trial. JAMA 2009; 301: 489-499.
- Hong JY, Kim WO, Yoon Y, Choi Y, Kim SH and Kil HK. Effects of intravenous dexmedetomidine on low-dose bupivacaine spinal anaesthesia in elderly patients. Acta Anaesthesiol Scand 2012; 56: 382-387.
- Ebert TJ, Hall JE, Barney JA, Uhrich TD and Colino MD. The effects of increasing plasma concentrations of dexmedetomidine in humans. Anesthesiology 2000; 93: 382-394.
- Talke P, Chen R, Thomas B, Aggarwal A, Gottlieb A, Thorborg P, Heard S, Cheung A, Son SL and Kallio A. The hemodynamic and adrenergic effects of perioperative dexmedetomidine infusion after vascular surgery. Anesth Analg 2000; 90: 834-839.
- Talke P, Richardson CA, Scheinin M and Fisher DM. Postoperative pharmacokinetics and sympatholytic effects of dexmedetomidine. Anesth Analg 1997; 85: 1136-1142.
- Yun-Sic Bang, Eunsun So, Seongsu Kim, Duk-Hee Chun. Effects of low dose midazolam on bradycardia and sedation during dexmedetomidine infusion. Int J Clin Exp Med 2016;9(6):11838-11844.
- Lee S, Kim BH, Lim K, Stalker D, Wisemandle W, Shin SG, et al. Pharmacokinetics and pharmacodynamics of intravenous dexmedetomidine in healthy Korean subjects. J Clin Pharm Ther. 2012;37:698-703. [PubMed]
- Hong GH. Edited by the Korean Society of Anesthesiologists, editors. Anesthesiology and Pain Medicine. 2nd ed. Seoul: Ryo Moon Gak; 2010. Monitoring and Anesthesia Record; pp. 142-143.
- Mizrak A, Ganidagli S, Cengiz MT, Oner U and Saricicek V. The effects of DEX premedication on volatile induction of mask anesthesia (VIMA) and sevoflurane

- requirements. *J Clin Monit Comput* 2013; 27: 329-334.
32. Bol CJ, Vogelaar JP, Tang JP and Mandema JW. Quantification of pharmacodynamic interactions between dexmedetomidine and midazolam in the rat. *J Pharmacol Exp Ther* 2000; 294: 347-355.
33. Bloor BC, Ward DS, Belleville JP and Maze M. Effects of intravenous dexmedetomidine in humans. II. Hemodynamic changes. *Anesthesiology* 1992; 77: 1134-1142.
34. De Jonge A, Timmermans PB and van Zwieten PA. Participation of cardiac presynaptic alpha 2-adrenoceptors in the bradycardiac effects of clonidine and analogues. *Naunyn Schmiedeberg Arch Pharmacol* 1981; 317: 8-12.