



ROLE OF DEXAMETHASONE IN BRACHIAL PLEXUS BLOCK

Anaesthesiology

Dr. Chandni Mehra	Post Graduate, Deptt of Anaesthesiology & Critical Care Nims Medical College & Hospital, Jaipur.
Dr. Zara Wani*	Post Graduate, Deptt Of Anaesthesiology & Critical Care Nims Medical College & Hospital, Jaipur. *Corresponding Author
Dr. Meenaxi Sharma	Professor, Deptt of Anaesthesiology & Critical Care Nims Medical College & Hospital, Jaipur.

ABSTRACT

Objective: To evaluate the effect of dexamethasone added to (lignocaine) on the onset and duration of axillary brachial plexus block.

Material and Methods: A total of 100 patients, who were scheduled for elective hand and forearm surgery under axillary brachial plexus block, were randomly allocated to group A in which patients received 40 ml 1.5% lidocaine with 2 ml of isotonic saline (0.9%) and group B in which patients received 40 ml 1.5% lidocaine with 2 ml of dexamethasone (8 mg). Nerve stimulator with insulated needle for multiple stimulations technique was used to locate the brachial plexus nerves. After the injection onset of action and duration of sensory blockade of brachial plexus were recorded at 5 minutes and 15 minutes interval.

Results: Group A showed the onset of action of 21.64 ± 2.30 min and in group B it was 15.42 ± 1.44 min ($p < 0.001$). Duration of nerve block was 115.08 ± 10.92 min in group A and 265.42 ± 16.56 min in group B ($p < 0.001$).

Conclusion: The addition of dexamethasone to 1.5% lignocaine solution in axillary brachial plexus block prolongs the duration of sensory blockade significantly.

KEYWORDS

Brachial plexus block, Dexamethasone, Lignocaine

INTRODUCTION:

The brachial plexus is formed by the ventral rami of C5-C6-C7-C8-T1, occasionally with small contributions by C4 and T2. There are multiple approaches to blockade of the brachial plexus, beginning proximally with the interscalene block and continuing distally with the supraclavicular, infraclavicular, and axillary blocks¹. Advantages of using brachial plexus over general anaesthesia are that there are less chances of decrease in blood pressure, undesirable decreases in cardiac output, central nervous system depression, respiratory depression, loss of protective airway reflexes (such as coughing), need for tracheal intubation and mechanical ventilation, and residual anaesthetic effects. Although brachial plexus block is not without risk, it is usually less invasive and affects fewer organ systems than general anaesthesia². Brachial plexus block is one of the most commonly used peripheral nerve blocks in clinical practice. It can be used as the sole anaesthetic technique or in combination with general anaesthesia for intra-operative and postoperative analgesia. Continuous catheterization of the brachial plexus is one of the best methods of postoperative analgesia. This block was performed by William Steward Halsted first time in 1889. He directly exposed the brachial plexus in the neck to perform the block and used cocaine^{3,4}. After that Herschel⁵, Kulenkampf⁶, Winnie and Collins⁷ described the percutaneous, supraclavicular and perivascular approach to the brachial plexus block respectively. The drugs commonly used for brachial plexus block are 2% lignocaine with adrenaline, 0.5% bupivacaine and by adding opioids to the local anaesthetic. Neuraxial blockade in elderly population needs less sedation, favouring early mobilization and excellent analgesia postoperatively⁸. Regional anaesthesia for forearm surgery can be provided by brachial plexus block through axillary approach, where surrounding nerves are injected with local anaesthetic agents⁹. Increasing the duration of local anaesthetic action is often desirable as it prolongs surgical anaesthesia and analgesia. Different additives have been used to prolong regional blockade¹⁰. Vasoconstrictors can be used to reduce vascular absorption of the local anaesthetic. Synergistic interaction can also occur when drugs affect different critical points along a common pathway¹¹. Different additives have been used to prolong regional blockade. It has been recently demonstrated that the nerve blocks are the ideal techniques for day case surgery¹². Addition of dexamethasone to the local anaesthetic increases the duration of the block^{13,14}. The purpose of this study was to evaluate the effect of dexamethasone added to (lignocaine) on the onset and duration of axillary brachial plexus block.

MATERIAL AND METHODS:

This randomized controlled trial was conducted on patients, who were

scheduled for elective hand and forearm surgery under axillary brachial plexus block, of either sex, aged 20 to 80 years having ASA grade I and II, who were willing to undergo the trial. Patients with a history of peptic ulcer disease, diabetes mellitus, hepatic or renal failure, pregnant women, morbidly obese with BMI more than 35, having coagulopathy or taking any premedications like opioids, benzodiazepines, and clonidine or having allergy to amide type local anaesthetics were excluded. One hundred patients were included in the study and randomly divided into two equal groups using random numbers table. The study was conducted after approval from the Hospital Ethical Committee and all data was collected after the informed consent of patients. Group A received 1.5% lignocaine plain only and in group B 8 mg dexamethasone was added in 1.5% lignocaine. Neither epinephrine nor bicarbonate was added to mixtures. All local anaesthetic solutions and adjuvant drugs were prepared by an anaesthesiologist not involved in the performance of block, patient care, or data collection to control the bias. On arrival to the operating room, standard monitoring was established (pulse oximetry, electrocardiography, and non-invasive arterial blood pressure monitoring) and oxygen was delivered via a Venturi facemask at a rate of 3 L/min. After insertion of an 18 gauge IV catheter in a peripheral vein in the contra-lateral arm and administration of 1 to 2 mg IV midazolam, axillary block was performed with the patient in the supine position and the upper arm abducted 90° and the elbow flexed at 110°. A nerve stimulator with a 24-gauge 7 cm sportte needle was used for precise localization of each nerve. The stimulation frequency was set at 3 Hz, the duration of stimulation at 0.1 ms, and the intensity of the stimulating current was initially set to deliver 3 mA and then gradually decreased. The position of the needle was considered to be acceptable when an output current Sensory blockade of each nerve was rated by the patient on a verbal analogue scale from 100% (normal sensations) to 0% (no sensation). The onset time of the sensory blockade was defined as the time between the end of last injection and the total abolition of the pinprick response. The duration of sensory block was considered as the time interval between the administration of the local anaesthetic and the first postoperative pain. The patients and the anaesthesiologist who evaluated the onset and duration of sensory blockades were blinded as to the mixture used. Data was entered in SPSS version 10.0 for statistical analysis. For quantitative variables, mean and standard deviation (SD) were calculated for description and independent samples' t-test was applied for comparison. For qualitative variables, frequency and percentage were calculated for description and chi-square test was applied for comparison. A p-value < 0.05 was considered as significant.

RESULTS:

The mean age of patients in group A was 41.18 ± 13.34 years while in group B it was 40.42 ± 12.50 years. In group A, male to female ratio was 3.5:1 and in Group B, male to female ratio was 7.3:1. In Group A, 29 (58%) patients were of ASA I while in group B, 25 (50%) patients were of ASA I. Both the groups were comparable with respect to age ($p = 0.769$), gender ($p = 0.813$) and ASA status ($p = 0.422$). (Table-1)

Table-1: Comparison of gender and ASA status between both the groups.

Variables	Group A (n=50)	44 (88%)	p-value
Gender:			0.183
male	39 (78%)	44 (88%)	
female	11 (22%)	6 (12%)	
ASA			0.422
ASA I	29 (58%)	25 (50%)	
ASA II	21 (42%)	25 (50%)	

The onset of action and the duration of nerve block showed a significant difference in the two groups. Group A showed the onset of action of 21.64 ± 2.30 min and in group B it was 15.42 ± 1.44 min ($p < 0.001$). Average duration of nerve block in group A was 115.08 ± 10.92 min while in group B it was 265.42 ± 16.56 min ($p < 0.001$). Table-2 shows the frequency of duration or sensory block between two groups.

Table-2: Comparison of duration of action between the groups.

Duration of Action	Group A (n=50)	Group B (n = 50)
< 130	50 (100%)	0 (0%)
131- 172	0 (0%)	0 (0%)
173 – 214	0 (0%)	0 (0%)
215 - 256	0 (0%)	18 (36%)
>256	0 (0%)	32 (64%)

In group A, all (100%) patients had the duration between 89-130 minutes, while in group B majority (64%) had duration > 256 minutes. This showed that group B had a more prolonged duration of brachial plexus sensory blockade as compared to group A.

DISCUSSION:

The addition of corticosteroid to local anaesthetic prolonged duration of blockade of the peripheral nerves. In this regard different studies were conducted to increase the duration of sensory and motor blockade and decreasing the onset of analgesia. In a prospective study conducted on sixty adult patients undergoing various orthopaedic surgeries on forearm and around the elbow under supraclavicular brachial plexus block by Maruf et al¹⁵ have concluded that addition of dexamethasone as an adjuvant to 2% lignocaine or 0.5% bupivacaine results in significantly early onset and markedly prolonged duration of analgesia without any unwanted effects. Parrington and colleagues¹⁶ in their study added 8 mg of dexamethasone to 30 ml 1.5% mepivacaine during supraclavicular brachial plexus blockade. The dexamethasone group showed a longer duration of analgesia: 332 (225- 448 min) vs. 228 (207-263 min) min in the control group, whereas the onset time of sensory and motor blockade were similar in both groups. In our study, in group A onset of action was 21.64 minutes and duration of axillary brachial plexus block was 115.08 minutes and in group B onset was earlier i.e., 15.42 minutes and duration of sensory blocked was prolonged i.e., 265.42 minutes. Our study results are comparable to the study by Movafegh et al¹⁷ who, in a randomized double-blind, study evaluated the effects of dexamethasone added to lignocaine on the onset and duration of axillary brachial plexus block in sixty patients scheduled for elective hand and forearm surgery. However, in their study the onset times of sensory and motor block were similar in the two groups but the duration of sensory (242 ± 76 versus 98 ± 33 min) blockade was significantly longer in the dexamethasone than in the control group ($p < 0.01$). Another study suggested that methylprednisolone can increase the duration of sensory and motor block when added to mepivacaine and bupivacaine¹⁸. Hamid et al¹⁹ found that butorphanol, a morphinan-type synthetic opioid analgesic exhibiting partial agonist at the μ opioid receptors and agonist activity at κ opioid receptors when added to 1% lignocaine can increase the duration of sensory blockade significantly. The mechanism of the analgesia induced by corticosteroids is not fully understood. This effect is suspected to be mediated by their anti inflammatory or immunosuppressive effects^{20,21}. The use of corticosteroids as an adjuvant to local anaesthetic for peripheral nerve blocks rarely has

been described, and its mechanism of action is not clearly understood. Corticosteroids cause skin vasoconstriction on topical application. The vasoconstriction effects of topical steroids are mediated by occupancy of classical glucocorticoid receptors rather than by nonspecific pharmacological mechanisms^{22,23}. According to the traditional theory of steroid action, steroids bind to intracellular receptors and modulate nuclear transcription. In our study, dexamethasone produced a relatively rapid effect which cannot be explained by the above mechanism²⁴. Therefore, vasoconstriction, the presumed mechanism of action for epinephrine's adjunctive effect on local anaesthetics, is probably not responsible for block prolongation by dexamethasone. Corticosteroids may have a local effect on the nerve; the dexamethasone effect may be related to this action²⁵. One possibility is that prolongation of local anaesthetic block occurs because of systemic effects of dexamethasone. Some authors believe that analgesic properties of corticosteroids are the result of their systemic effects^{26,27}. To elucidate the mechanism of action of dexamethasone and to answer the question of whether these results were attributable to a local or systemic effect, well designed further studies are warranted. Adding a steroid to local anaesthetic solution may not be indicated for all patients. For example, diabetic patients may experience hyperglycemia and patients with a continuing infectious process may be detrimentally affected by the anti inflammatory effects of steroids. The use of dexamethasone to increase the duration of action of local anaesthetics is not an indication of this drug. This study led us to hypothesize that it may be useful in situations in which epinephrine must be used with caution (e.g., hypertension, ischemic heart disease).

CONCLUSION:

We conclude that, the addition of dexamethasone 8 ml to 1.5% lignocaine (38 ml) solution in axillary brachial plexus block prolongs the duration of sensory and motor blockade. However, further well designed studies are needed to evaluate the optimal dose and mechanism of action of dexamethasone to be used to prolong the brachial plexus block duration.

REFERENCES:

- Fisher L, Gordon M, Wolfe SW, Hotchkiss RN, Pederson. Green's Operative Hand Surg 6th ed., Philadelphia, Elsevier/Churchill Livingstone, 2011: 25-40.
- Boedeker BH, Rung GW, Zajchuk R, Bellamy RF, Grande CM. Regional anesthesia. Textbook of Military Medicine, Part IV: Surg Combat Casualty Care. 1: Anesthesia and Perioperative Care of the Combat Casualty. Washington, DC: Borden Institute. 1995; 251-86.
- Halsted WS. Practical comments on the use and abuse of cocaine suggested by its invariably successful employment in more than a thousand minor surgical operations. New York Med J, 1885; 42: 294-5.
- Crile GW. "Anesthesia of nerve roots with cocaine, Cleveland Med J 1897 2: 355.
- Hirschel G. Anesthesia of the brachial plexus for operations on the upper extremity. Munchener Medizinische Wochenschrift German 1911; 58: 1555-6.
- Kulenkampff D. On anesthesia of the brachial plexus. Chirurgie ; German, 1911; 38: 1337-40.
- Winnie AP, Collins VJ. The subclavian perivascular technique of brachial plexus anesthesia. Anesthesiology 1964; 25: 35-63.
- Pöpping DM, Elia N, Marret E, Wenk M, Tramèr MR. Clonidine as an adjuvant to local anesthetics for peripheral nerve and plexus blocks: A meta-analysis of randomized trials. Anesthesiology 2009; 111: 406.
- Aasboe V, Raeder JC, Grogaard B. Betamethasone reduces postoperative pain and nausea after ambulatory surgery. Anesth Analg 1998; 87: 913-7.
- Tsui BC, Wagner A, Finucane B. Regional anaesthesia in the elderly: a clinical guide. Drugs Aging 2004; 21: 895-9.
- Handoll HHG, Koscielniak-Nielsen ZJ. Single, double or multiple injection technique for axillary brachial plexus block for hand, wrist or forearm surgery. Cochrane Database Syst Rev 2006; 25: Cd003842.
- Capdevila X, Dadure C. Perioperative management for one day hospital admission: regional anesthesia is better than general anesthesia. Acta anaesthesiol Belg 2004; 55: 33-6.
- Adnan T, Elif AA, Ayshe K. Clonidine as an adjuvant for lidocaine in axillary brachial plexus block in patients with chronic renal failure. Acta Anaesthesiol Scand 2005; 49: 563-8.
- Tandoc MN, Fan L, Kolesnikov S, Kruglov A. Adjuvant dexamethasone with bupivacaine prolongs the duration of interscalene block: a prospective randomized trial. J Anesth 2011 Oct; 25(5): 704-9. Epub 2011 Jun 17.
- Islam SM, Hossain MHMD, Maruf AA. Effect of addition of dexamethasone to local anesthetics in supraclavicular brachial plexus block. JAFMC Bangladesh 2011; 7(1).
- Parrington SJ, O'Donnell D, Chan VW, Brown-Shreves D, Subramanyam R, Qu M, et al. Dexamethasone added to mepivacaine prolongs the duration of analgesia after supraclavicular brachial plexus blockade. Reg Anesth Pain Med 2010; 35: 422-6.
- Movafegh A, Razavian M, Hajimaohamadi F, Mevsamie A. Dexamethasone added to lidocaine prolongs Axillary brachial plexus blockade. Anesth Analg 2006 102; 263-7.
- Stan T, Goodman E, Cardida B, Curtis RH. Adding methylprednisolone to local anesthetic increases the duration of axillary block. Reg Anesth Pain Med 2004; 29: 380-1.
- Mir IH, Hamid A. Addition of Butorphanol to Lidocaine prolongs duration of the Axillary Brachial Plexus Block. The Internet Journal of Anesthesiology 2008; 16(1): 92.
- McCormack K. The spinal actions of nonsteroidal anti-inflammatory drugs and the dissociation between their anti-inflammatory and analgesic effects. Drugs 1994; 47: 28-45.
- Ahlgren SC, Wang JF, Levine JD. C-fiber mechanical stimulus-response functions are different in inflammatory versus neuropathic hyperalgesia in the rat. Neuroscience 1997; 76: 285-90.
- Marks R, Barlow JW, Funder JW. Steroid-induced vasoconstriction: glucocorticoid

- antagonist studies. *J Clin Endo Meta* 1982; 54: 1075-7.
23. Seidenari S, Di Nardo A, Mantovani L, Giannetti A. Parallel intraindividual evaluation of the vasoconstrictory action and the antiallergic activity of topical corticosteroids. *Exp Dermatol* 1997; 6: 75-80.
 24. Taguchi H, Shingu K, Okuda H, Matsumoto H. Analgesia for pelvic and perineal cancer pain by intrathecal steroid injection. *Acta Anaesthesiol Scand* 2002; 46: 190-3.
 25. Devor MD, Gorvin-Lippmann R, Raber P. Corticosteroids suppress ectopic neural discharge originating in experimental neuromas. *Pain* 1985; 22:127-37.
 26. Aasboe V, Raeder JC, Groegaard B. Betamethasone reduces postoperative pain and nausea after ambulatory surgery. *Anesth Analg* 1998; 87: 913-7.
 27. Baxendale BR, Vater M, Lavery KM. Dexamethasone reduces pain and swelling following extraction of third molar teeth. *Anaesthesia* 1993; 48: 961-4.