



AMELOBLASTIC CARCINOMA -SECONDARY TYPE, A RARE CASE REPORT AND DISTINCTION FROM MALIGNANT AMELOBLASTOMA

Surgery

Dr Rajesh B Dhirawani	Prof and HOD, Department of Oral and Maxillofacial Surgery, Hitkarini Dental College and Hospital
Dr Sumit Asrani	Senior Lecturer, Department of Oral and Maxillofacial Surgery, Hitkarini Dental College and Hospital
Dr Anshalika Agrawal	PG Resident (I) Department of Oral and Maxillofacial Surgery, Hitkarini Dental College and Hospital
Dr Sauvik Singha*	PG Resident (II) Department of Oral and Maxillofacial Surgery, Hitkarini Dental College and Hospital *Corresponding Author
Dr Kshitij Aggarwal	PG Resident (III) Department of Oral and Maxillofacial Surgery, Hitkarini Dental College and Hospital

ABSTRACT

Ameloblastic Carcinoma (AC) primarily is an aggressive odontogenic tumour, which histologically consist of features of both ameloblastoma and carcinoma. This requires more aggressive surgical approach than its benign counterpart. Malignant Ameloblastoma (MA) microscopically appears benign but shows metastasis. The authors reported an extremely rare secondary type of Mandibular Ameloblastic Carcinoma of a 50 year old Indian male with a complaint of swelling, trismus, significant bone resorption with tooth mobility. This article discussed about the clinical features, radiographic features, management, and histological characteristics of AC and its distinction from MA.

KEYWORDS

INTRODUCTION-

Ameloblastic Carcinoma (AC) is a rare malignant odontogenic tumour that has features of ameloblastoma intermingled with features of carcinoma, regardless of whether it has metastasized¹ i.e. ameloblastoma in which there is histologically malignant transformation with less differentiated metastatic growth². Malignant Ameloblastoma (MA) is the lesions that metastasizes and are well differentiated benign lesions AC in both primary and metastatic sites³. In contrast to ameloblastoma, presents more aggressive clinical features, such as rapid growth, perforation of cortex, painful swelling and trismus⁴. AC mostly occurs in posterior mandible region with only 1/3rd cases in maxilla region, and mostly occurs in around 30 yrs of age¹⁰. Radiographic findings include a poorly defined radiolucency, sometimes with focal radiopacities due to necrosis with dystrophic calcifications^{7, 8}. (Ac)ording to latest WHO 2005 update, (AC) are classified in to Primary type i.e. histologic features of ameloblastoma with cytologic atypia and Secondary type, i.e. malignant transformation of pre-existing benign ameloblastoma¹¹. Most cases of (AC) are denovo and less than 1% of ameloblastomas undergo malignant transformation. Hence secondary type is less frequent than primary¹²⁻¹⁵. In this article, a case of (AC) secondary type is presented and its clinical examination, radiographic examination, histopathology and management are discussed. Meanwhile the published articles about the (AC) are reviewed and its distinction from MA is pointed.

CLINICAL EXAMINATION –

A 50-year-old Indian male patient referred to our department with the complaint of rapidly growing painless swelling of the right side of lower jaw crossing midline with difficulty in chewing, multiple mobile teeth over affected region. He had the past history of a similar lesion over the right lower jaw, which was treated by Hemimandibulectomy with Disarticulation of right TMJ 12 years back. The histopathology then was suggestive of ameloblastoma. On extraoral examination, a firm diffuse non-tender swelling over right side of f(Ac)e extended superiorly from the right infraorbital region to the lower border of mandible inferiorly. Medially, it extended up to one-third of both upper and lower lips with an elevated angle of mouth and laterally 1 inch anterior to the angle of mandible with more medio-lateral expansion measuring about 10 cm × 8 cm × 6 cm [FIGURE 1A]. Neither paresthesia was associated with the swelling nor was any regional lymph node palpable. Intraoral examination revealed a painless firm proliferative growth crossing midline up to the right anterior border of ramus. The swelling was negative on aspiration [FIGURE 1B].

RADIOGRAPHIC EXAMINATION-

3 Dimensional computed tomography showed hypodense image of approximately 5 cm × 7 cm size present over left side mandibular and submandibular region, an area of resorption over left mandibular premolar region with att(Ac)hed 2.4 mm titanium reconstruction plate. [FIGURE 3]

INCISIONAL BIOPSY-

The patient was then planned for an incisional biopsy under local anesthesia. Histopathological examination with H & E stain was conclusive of diagnosis of AC.

SURGERY-

He was planned for resection of tumor under general anesthesia. The submandibular incision was given and the tumor appeared firm in consistency, surrounding the reconstruction plate. The tumor was removed along with the reconstruction plate and reconstruction was done with Pectoralis major myocutaneous flap [FIGURE 2A, 2B]. Hemostasis (Ac)hieved and closure done in layers. The postoperative recovery was uneventful.

HISTOPATHOLOGY-

Microscopic examination showed ameloblastic islands with palisading of peripheral cells showing reverse polarity. Within the islands marked features of malignancy were seen that is increased mitosis, hyperchromatism, and pleomorphism. Some of the islands lost its arrangement of peripheral palisading columnar cells and exhibited irregularly arranged squamous cells with mitotic figures, vesicular nuclei and prominent nucleoli. [FIGURE 4]

FOLLOWUP-

Then we referred the patient to department of radiotherapy. They have given 25 fractions of 1.8 Gy e(Ac)h for a period of 5 weeks (45 Gy). Then they referred the patient to Department of Pulmonary Medicine and Gastroenterology to evaluate distant metastasis, reports was negative. The patient is on regular follow-up since 1 year and doing well.

DISCUSSION-

In 1983, Shafers introduced the term (AC), which is a rare malignant neoplasm that requires correct histologic diagnosis and aggressive surgical intervention. The tumor cells resemble the cells seen in ameloblastoma, but they show cytologic atypia. Mean time span from symptoms to treatment is shorter for (AC) (11 months) as compared

with benign ameloblastoma (27months) as its progression is faster^{16,17}. Most (Ac), primary type, arise de novo and histologically defined as an apparent malignant neoplasm that exhibits cytological malignancy, including pleomorphism, hyperchromatic nuclei, and high mitotic rates. The rare variety of (Ac), secondary type, is thought to occur from the recurrent or pre-existing ameloblastoma, and it may have the histologic features of a coexisting ameloblastoma^{12, 15, 18}. The clinical features, biologic behavior, and treatment of (AC), secondary and primary types, are almost identical. Because of its rarity, no full proof therapeutic guidelines exist for (Ac). Some clinicians recommend wide resection. However a small number of reports have described radiation therapy and chemotherapy as treatment modality with Ion beam therapy as a new measure.

After reviewing all article on AC present in Pubmed and Cochrane online digital library till date we got that AC mostly occurs in sixth to seventh decade of life. They are seen twice as often in male patients as in female patients. The most commonly involved area is the posterior portion of the mandible²⁶. Involvement of the maxilla by (AC) seems to be less frequent²⁵⁻²⁸. The most common sign described has been swelling, although others include associated pain, rapid growth, trismus and dysphonia²⁶. The radiographic appearance of the lesion was consistent with that of an ameloblastoma except for the presence of some focal radiopacities, apparently reflecting dystrophic calcifications²⁶.

MA is a very rare tumor, defined in 1992 by the World Health Organization (WHO) as “a neoplasm in which the pattern of an ameloblastoma and cytologic features of malignancy are shown by the primary jaw lesion and/or by any metastatic growth. “The term MA was especially controversial because of the former WHO classification, in which the diagnosis was based on the presence of metastasis. Histologically, MA does not show uniform proliferation, but it has cellular pleomorphism. Spindling of the cells is recognized in some solid proliferating areas²⁹.”

The two forms of malignancy described in association with ameloblastomas are the MA and (AC). For a diagnosis of MA, the cardinal feature is metastatic spread, because the histological appearance of the primary and metastatic lesion(s) is indistinguishable from the typical benign ameloblastoma. The most common sites of metastases are reported to be the chest (lungs, pleura, hilar nodes) in 75% of cases and the cervical lymph nodes or spine in 15%.²⁴ The mode of spread of MA is debated as being either by inhalation at the time of surgery, haematogenous or lymphatic spread. The management of MA.^{19,20} Isolated and discrete lung metastases have been treated by open thor(Ac)otomy and wedge resection 6V15 and chemotherapy has occasionally been used successfully^{21,22,23}.

In the present case the lesion is confined clinically and radiographically, within the mandible with evidence of extension of a carcinoma into gingiva, alveolar ridge, floor of mouth, but no evidence of metastasis to distant site. Microscopically, there is carcinoma as well as morphologic features that resembled ameloblastoma, arose in a preexisting ameloblastoma. These clinical, radiographic, pathologic findings support the diagnosis of an (AC) secondary type. Only a limited number of (AC)s secondary type cases have been reported in the English-language literature till 2017, so this present report adds one more case to the scanty number of publication.

REFERENCES

1. Suaimura M, Yamauchi T, Yashikawa K, Takeda N, Sakita M, Miyazaki T. Malignant ameloblastoma with metastasis to the lumbar vertebrae: report of case. *J Oral Surg* 1969; 27:350-6.
2. Shafer WG, Hine MK, Levy BM: A textbook of oral pathology, ed. 3, Philadelphia, 1974, W.B. Saunders Company, p. 254.
3. Slootweg PJ, Müller H. MA or AC. *Oral Surg Oral Med Oral Pathol*.1984;57:168–76. [PubMed: 6366686]
4. Akhrish S, Buchner A, Shoshani Y, Vered M, Dayan D. (AC): report of a new case, literature review, and comparison to ameloblastoma. *J Oral Maxillofacial Surg* 2007; 65:777–83.
5. Waldron CA, Small IA, Silverman H. Clear cell ameloblastoma: an odontogenic carcinoma. *J Oral Maxillofac Surg* 1985;43:707-17.
6. LolAchi CM, Madan SK, Jacobs JR. AC of the maxilla. *J Laryngol Otol* 1995;109:1019-22.
7. Corio RL, Goldblatt LI, Edwards PA, Hartman KS. (AC): a clinicopathologic study and assessment of eight cases. *Oral Surg Oral Med Oral Pathol* 1987;64:570-6.
8. Infante-Cossio P, Hernandez-Guisado JM, Fernandez-Machin P, Garcia-Perla A, Rillon-Mayordomo A, Gutierrez-Perez JL. (AC) of the maxilla: a report of 3 cases. *J Craniomaxillofac Surg* 1998;26:159-62.
9. Sastre J, Munoz M, Naval L, Adrados M. AC of the maxilla: report of a case. *J Oral Maxillofac Surg* 2002;60:102-4.
10. Simko EJ, Brannon RB, Eibling DE. AC of the mandible. *Head Neck* 1998;20:654-9.
11. Barnes L, Eveson J, Reichart P, Sidransky D. World Health Organization classification of tumours; pathology and genetics of head and neck tumours. Lyon: IARC Press; 2005. p. 286-91.
12. Slootweg PJ, Müller H: MA or AC. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 57:168, 1984
13. Cox DP, Muller S, Carlson GW, et al: AC ex ameloblastoma of the mandible with malignancy-associated hypercalcaemia. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 90:716, 2000
14. Cizmecy O, Aslan A, Onel D, et al: AC ex ameloblastoma of the mandible: Case report. *Otolaryngol Head Neck Surg* 130:633, 2004
15. Karakida K, Aoki T, Sakamoto H, et al: AC, secondary type: A case report. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 110:e33, 2010
16. Slootweg PJ, Müller H. MA or AC. *Oral Surg Oral Med Oral Pathol*. 1984;57:168–76.
17. Cox DP, Muller S, Carlson GW, Murray D. AC ex ameloblastoma of the mandible with malignancy-associated hypercalcaemia. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*. 2000;90:716–22.
18. Jensen AD, Ecker S, Ellerbrock M, et al: Carbon ion therapy for AC. *Radiat Oncol* 6:13, 2011
19. Seward GR, Beales SJ, Johnson NW, Sita Lumsden E. A metastasising ameloblastoma associated with renal calculi and hypercalcaemia. *Cancer* 1975; 36: 2277-2285.
20. Madiedo G, Choi H, Kleinman JG. Ameloblastoma of the maxilla with distant metastases and hypercalcaemia. *Am J Clin Pathol* 1981; 75: 585-591.
21. Ueda M, Kosaki K, Kanedo T, Imaizumi M, Abe T. Doubling time of ameloblastoma metastasizing to the lung: report of two cases. *J Craniomaxillofac Surg* 1992; 20: 320-322
22. Lanham RJ. Chemotherapy of metastatic ameloblastoma. A case report and review of the literature. *Oncology* 1987; 44:133-134.
23. Ramadas K, Jose CC, Subhashini J, Chandni SM, Viswanathan FR. Pulmonary metastases from ameloblastoma of the mandible treated with cisplatin, adriamycin and cyclophosphamide. *Cancer* 1990; 66: 1475-1479.
24. Laighlin EH: Metastasizing ameloblastoma. *Cancer* 64:776-780, 1988
25. Slootweg PJ, Müller H. MA or AC. *Oral Surg Oral Med Oral Pathol* 1984; 57(2): 168–76.
26. Corio LR, Goldblatt LI, Edwards PA, Hartman KS. AC: a clinicopathologic study and assessment of eight cases. *Oral Surg Oral Med Oral Pathol* 1987; 64(5):570–6.
27. Lee L, Maxymiw WG, Wood RE. AC of the maxilla metastatic to the mandible. Case report. *J Craniomaxillofac Surg* 1990; 18(6):247–50.
28. Daramola JO, Abioye AA, Ajagbe HA, Aghadiuno PU. Maxillary MA with intraorbital extension: report of case. *J Oral Surg* 1980; 38(3):203–6.
29. Kramer IRH, Pindborg JJ, Shear M. Histological typing of odontogenic tumours. Heidelberg, Germany, Springer-Verlag, World Health Organisation 1992




Fig 1a) Recurrence after 12 year following surgery
 Fig 1b) Intra proliferative growth in the mandibular anterior region
 Fig 2a) Operated site after the resection of tumour
 Fig 2b) Resected surgical specimen
 Fig 3) computed tomographic scan shows bony destruction of left mandible crossing midline
 Fig 4) Histological examination with Hematoxylin-eosin, original magnification x 20