



## STUDY OF VARIOUS RISK FACTORS RELATED TO HYPOTHYROIDISM IN ADULT PATIENTS AT PMCH, PATNA

### Medicine

**Dr. (Prof) Kaushal Kishore**

Professor Department of Medicine, PMCH Patna, Bihar

### ABSTRACT

The aim to assess various contributing factors related to hypothyroidism for adult patients, and find out the relationship between some demographic characteristic (age, gender, level of education) with risk factors.

**Methodology:** A descriptive study was conducted at the outpatient department in department of medicine, PMCH, Patna in period between October 2016 to May 2017 in order to assess the risk factors related to Hypothyroidism for adult patient.

A random sample consisted of (50) patients from suffering hypothyroidism who were systematically admitted to the outpatient consultation were selected one by one. A questionnaire was constructed for the purpose of the study, which is comprised of three parts that include: (1) Socio demographic Characteristics data (2) The past medical history for patient and family (3) Assessment of risk factors related to diet, psychological factors, social factors and medication. Reliability and validity of questionnaire was determined through test re-test ( $r=0.861$ ) of pilot study. Data was collected by the researcher who interviewed those patients and filled out the constructed questionnaire form. Data were analyzed by using descriptive statistical approach (frequency, percentage and mean of score) and inferential statistical approach (standard deviation and correlation coefficient).

**Results:** The study findings indicated that there is highly significant difference between Women and man the incidence of hypothyroidism disease more than female. It is aged between (28-37) years old for hypothyroidism patients, the most of study sample primary graduates and urban setting. Our results also show that there is significant difference between history of thyroid disease patient and non-history of thyroid disease patients.

**Conclusion:** from the study that past history of thyroid disease raises the risk to develop hypothyroidism disease. Recommendations: The study recommends that the further research aimed at established the new department for early detection hypothyroidism for adult patient and education program for nurses to improve the patient quality of life.

### KEYWORDS

Risk factors, Hypothyroidism.

### Introduction

Worldwide about one billion individuals are estimated to be iodine deficient; however, it is unknown how often this leads to hypothyroidism. In large population-based studies in Western countries with deficiency dietary iodine, 0.3–0.4% of the population have overt hypothyroidism. A larger proportion, 4.3–8.5%, has subclinical hypothyroidism (1) is the production of an imbalance of thyroid hormone. Hypothyroidism is simply insufficiency thyroid hormone either imbalance affects the metabolism in the body. Guyton (2000) says “hypothyroidism like hyperthyroidism is probably initiated by autoimmunity against the thyroid gland, but immunity that destroys the gland rather than stimulates it. The thyroid gland regulates the body's metabolism, heart rate, blood pressure, and body temperature, among other functions. Thyroid hormone maintains the rate at which your body uses fats and carbohydrates, burn calories, assist regulate body temperature, influence heart rate and helps in controlling the production of protein. Sometimes the thyroid can produce too much hormones or insufficiency (2) are most probability to develop hypothyroidism than men. In population-based studies, women were seven times more likely than men to have TSH levels over 10 mU/l (3). Many risk factors predispose women and man to hypothyroidism including autoimmune thyroid disease (AITD), type (1) diabetes mellitus, other endocrine deficiencies, Down syndrome, thalassemia major, thyroid ablation and who are treated with pituitary surgery or irradiation, head and neck irradiation, treatment of growth hormone deficiency, cytotoxic therapy, lithium, interferon  $\alpha$ , interferon  $\beta$ , and therapeutic monoclonal antibodies. Patients with a family or client history of thyroid disease, goiter, history of spontaneous abortion in women, Positive thyroid peroxidase antibody, origin from areas of endemic iodine deficiency, Previous thyroid disease or surgery, or any symptoms suggesting at higher risk for hypothyroidism and hyperthyroidism (4). To assess risk factors related to hypothyroidism for adult patients.

### III. Methodology

A descriptive study was carried out to assessment risk factors related to hypothyroidism for adult patients who attended the outpatient department of medicine PMCH Patna. The study was carried out during the period extended from the October 2016 to May 2017. The sample consisted of (50) patient's hypothyroidism. A questionnaire-interview format was designed and developed by the researcher for the

purpose of the study.

All the items were measured on scale of (3) indicates that the always 1, sometimes 2, never 3. Diet domain; daily 3, Weekly 2, monthly 1 for hypothyroidism. The questionnaire consisted of (3) parts. Part I: Demographic Information Sheet. Part II: historical information satisfactory to the patient and family. Part III: Risk factors domain Which include: diet, Psychological domain, social and Medication domain. Rating scale was used to rate the frequency and extension of the problems. The content validity of the instrument was established through a panel of (14) experts. Test-retest reliability was determined through a computation of Pearson Correlations for the scales. Coefficients for the (43) items of risk factors of hypothyroidism were ( $r=0.861$ ) for the total score of risk factors. Using the questionnaire-structured format through interview technique collected the data. The researcher interviewed each patient personally. The data were analyzed through descriptive data analysis and inferential data analysis the data were analyzed through the use of Statistical Package of Social Sciences (SPSS) version 16.0.

### IV. Results

**Table (1): Distribution of the study Samples by Socio-Demographic Characteristics.**

no	variables	groups	frequency	percentage
1	gender	Male		
		Female		
2	Age Groups (Per years)	18- 27 years	10	20
		28- 37 years	16	32
		38- 47 years	12	24
		48 - 57 years	9	18
		58- 67 years	3	6
3	Residence	Urban	44	88
		Rural	6	12
4	Marital status	Single	5	10
		Married	40	80
		Widow	4	8
		Divorce	1	2

5	Level of Education	Illiterate	5	10
		Read & write	3	6
		Primary Graduate	15	30
		Intermediate Graduate	10	20
		Secondary Graduate	6	12
		Diploma Graduate	3	6
		College Graduate	7	14
6	Occupation	Govern. Employee	11	22
		Free job	2	4
		Retired	1	2
		Housewife	35	70

Table (1) revealed that 96% of the study samples were females, more age (28-37) years old for hypothyroidism, high percentage of them were primary graduated hypothyroidism 30%, 70% of them were housewife for hypothyroidism,(88%) study sample from urban setting and 38% have barely sufficient income for hypothyroidism .

**Table (2): Distribution of Patients and Family according to Hypothyroidism Related to Clinical History.**

LIST	CLINICAL HISTORY		
		Frequency	Percentage
1st	Clinical history for patient		
1	High cholesterol in the blood	6	12
2	Heart disease	8	16
3	Hypertension	15	30
4	Previous surgery of thyroid gland	12	24
5	Diabetes mellitus	5	10
6	Use of radioactive iodine	6	12
7	rheumatoid arthritis	26	52
2nd	Clinical history for family	28	56

Table (2) shows that highest percentage (56%) had rheumatoid arthritis as past medical history and 56% of patients had family history of hypothyroidism

**Table (3): Social and medication domain for adult patients with Hypothyroidism.**

Sl. no	Social domain	Always	Sometimes	Never	MS	ASS
1	Do take advantage of your free time?	18	11	21	2.06	M
2	Participate in social activities	34	5	11	1.54	L
3	Do you like to be relationships with others?	43	4	3	1.2	L
4	Do you deal easily with your colleagues ?	42	5	3	1.22	L
5	Are increasing problems during menstrual ?	10	5	35	25	L
6	Feel anxiety when you stay at home.	4	38	8	2.08	M
7	Prefer to stay at home away	2	24	24	2.24	H

**Table(4) shows that the mean of score are high on items3rd (5,7), 4th (2,5,6,7),moderate and low on the remaining items.**

Sl. no	Medication domain	Always	Sometimes	Never	M.S	ASS
1	Antacids drugs	15	11	24	2.18	M
2	Iron and product	6	18	26	2.4	H
3	Contraceptive drugs	7	18	25	2.36	M
4	Sedatives and opiates	8	17	25	2.34	M
5	Anti-thyroid drug	2	3	45	2.86	H
6	Cardiovascular drugs	9	10	31	2.44	H
7	Diuretic drugs	3	3	44	2.82	H

M.s=mean of score (1-1.69= low,1.7-2.39= moderate,2.4-3= high)

**Table (5) Correlation coefficient between (gender, age, level of education) with risk factors domains for hypothyroidism patients.**

Correlations	Gender	Age(years)	Level of education	Risk factors
Gender	1	.101	.058	.162
Age(years)	.101	1	.136	.028
Level of education	.058	.136	1	.144
Risk factors	.162	.028	.144	1

\*\* Correlation is significant at the 0.01 level (2-tailed). \*\*Correlation is significant at the 0.05 level (2-tailed).

This table (5) show that there is weak negative relationship between contributing factors with gender (r=-.162) , and there is positive moderate relationship between contributing factors with level of education(r=.144) , there is relationship between contributing factors with age(r=.028).

**V. Discussion**

Through the course of the data analysis of the present study, the finding showed that the majority of the study samples for hypothyroidism (96%) females (32%) were (28- 37) years old. This finding agree with other researcher who stated that diagnosed hypothyroidism was ( 2.5% ) in females and (0.6%) in males, hypothyroidism( 4.8%) in females and (0.9%)for males(5).

Seventy percent of the study samples group is housewife as occupation and Thirty eight percent are barely sufficient for hypothyroidism as monthly income. This finding disagrees with other researcher who reported that a very high percentage (54.4%) of the patients with thyroid disease is disability retired (6).

Regarding residence, majority (88%) of the study was urban for hypothyroidism. This finding were in good agreement with that obtain by other researcher which reported that the finding of participants had (19%) participants from urban areas, (81%) participants from rural areas (7).

The finding of the study sample shows that (44%) of the patient had family history for hypothyroidism. This findings can be supported by another study the findings who reported that there was no significant difference between those who had history of thyroid problem in their family and those who did not have history of thyroid problem in their family in cases as well as in control group(8).

The result of accurate study that show diet domain in table (3)for hypothyroidism patients are high effect in (butter and pineapple).The mean of score is low effect in (white meat), (Cheese) and (Lettuce). The mean of score is moderate effect in( Red meat ),(Milk) , (Spanish) and (Green pepper). The finding of the diet domains show that moderate effect for hypothyroidism patient. The researcher believes diet domain the patient have bad dietary habit and no had diet information about nutrition value that effect on health statuses.

This finding can be supported by another study the findings which reported that we get iodine from foods (butter, pineapple) and it is found in commercially processed iodized salt and preservatives used in bread products. High iodine can inhibit thyroid hormone production

by increasing cellular resistance to T3 and may lead to iodine induced thyroid deficiency, swelling, tenderness and goiter(9).

Our finding about psychological domain are similar to those reported that of the other researcher who stated there was predominance of Major depressive disorder 24.2%, followed by anxiety disorders (23%). The p-value of the comparison between the case and the control patients is 0.0001 which is highly significant (10).

The mean of score of social domain in table (4) for hypothyroidism patients are high on item (Are increasing problems during menstrual cycle days?) and item (Prefer to stay at home away from the eyes of the people). The mean of score is moderate item (Do take advantage of your free time?), item (Feel anxiety when you stay at home) and low on the remaining items. This finding agrees with other researcher which reported that the patients of thyroid disease were significantly more anxious and perceived higher social support from family when compared with the control group (11).

The mean of score of medication domain in table (4) for hypothyroidism patients are high on item (Iron and product), item (Anti-thyroid drugs), item (Cardiovascular drugs) and item (Diuretic drugs). The mean of score is moderate item (Antacids drugs), item (Contraceptive drugs) and item (Sedatives and opiates).

These results were similar to those results obtained by other researcher who stated that many different drugs affect thyroid function. Most of these medications act at the level of the thyroid in patients with normal thyroid function, or at the level of thyroid hormone absorption or metabolism in patients requiring exogenous levothyroxine (12).

The findings of the study sample hypothyroidism patients indicated that there is a weak negative relationship between contributing factors with gender ( $r = -.162$ ), and there is a weak positive relationship between contributing factors with level of education ( $r = .144$ ), there is no relationship between contributing factors with age ( $r = .028$ ). Table (5)

## VI. Conclusions

Our data indicate that the contributing factors domains of patients with hypothyroidism is moderate decline in all domains except medication domain is high of risk factors domains.

## References

- [1] Garber, JR., Cobin, RH., Gharib, H., Hennessey, JV., Klein, I., Mechanick, JI., Pessah-Pollack, R., Singer, PA., Woeber, KA for the American Association of Clinical Endocrinologists and the American Thyroid Association Taskforce on Hypothyroidism in Adults. "Clinical Practice Guidelines for Hypothyroidism in Adults". *Thyroid*, Vol.22, NO.12, (2012), P.P:1200-1235.
- [2] Huether, S., & McCance, K.: *Understanding Pathophysiology* (4th ed.) (2007), St. Louis: Mosby.
- [3] Cooper, DS., Braverman, LE.,; Werner & Ingbar's the thyroid: a fundamental and clinical text (10th ed.). Philadelphia: Wolters Kluwer/Lippincott Williams & Wilkins Health. (2012), P. 552. ISBN 145112063X
- [4] Patrici, A.; American Thyroid Association. "Hypothyroidism." *Clinical Thyroidology for Patients*. 2:3. (2009). 30 Sept 2009 <[http://thyroid.org/patients/ct/volume2/issue3/ct\\_patients\\_v23\\_5.html](http://thyroid.org/patients/ct/volume2/issue3/ct_patients_v23_5.html)>.
- [5] Bjuro T., Holmen, J., Krue, E., Midthjell, K., et. al.: Prevalence of thyroid disease, thyroid dysfunction and thyroid peroxidase antibodies in a large, unselected population. The Health Study of Nord-Trøndelag (HUNT), *European Journal of Endocrinology*, Vol.143, (2000), pp. 639-647.
- [6] Vasile, P.; Screening and psychological evaluation of patients with thyroid dysfunction, "Babeş-Bolyai" University Cluj-Napoca faculty of psychology and educational Sciences, 2011, p.11. Viruses and thyroiditis; update, Rachel Desailoud and Didier Hober, *Virology Journal*, Vol.6, (2009), P.5.
- [7] Munifa, Toto, S., Untung, S., Widodo.: Eating pattern and smoking as risk factors for hyperthyroidism. public health postgraduate program faculty of medicine gadjarmada university yogyakarta. (2011), NIM: 09/293533/PKU/11009.
- [8] Raghda, A.; Assessment of Thyroid Function in Pregnant Women From Rimal Health Center, Gaza City, The Islamic University- Gaza, deanery of higher education, faculty of Science, master of biological sciences, medical technology 12/2010-1\1432.
- [9] Brinker, Francis. *Herb Contraindications & Drug Interactions*. 3rd Ed. Sandy, OR: Eclectic Medical Publications, 2001.
- [10] Sheikh, S., Raheel, M., Mohammad, M., Tasleem, A.; Psychiatric Manifestations in thyroid disorders, *International Journal of Clinical Cases and Investigations*. Vol.5, NO. 3, (2013), P.P:84-98.
- [11] Petra M.; Psychosocial Factors in Patients with Thyroid Disease, Thyroid and Parathyroid Diseases -New Insights into Some Old and Some New Issues, Dr. Laura Ward (Ed.), ISBN: 978-953-51-0221-2012.
- [12] Deborah, M.; clinical expression, pathophysiology consequences and general health status in elderly individuals with subclinical thyroid dysfunction, University of Birmingham Research Archive, June 2011.