



## PROSPECTIVE STUDY OF CALCULOUS CHOLECYSTITIS

## General Surgery

**Dr. Sharad M. Patel** Professor Of Surgery, Smt. S.C.L. Hospital, Ahmedabad, Gujarat

**Dr. Naman K. Patel\*** Assistant Professor Of Surgery, Smt. S.C.L. Hospital, Ahmedabad, Gujarat  
\*Corresponding Author

**Dr. Pranavkumar Parthsarathi** 3<sup>rd</sup> year Surgery Resident, Smt. S.C.L. Hospital, Smt. N.H.L. Municipal Medical College, Ahmedabad, Gujarat

**Dr. Parth Prajapati** 2<sup>nd</sup> Year Surgery Resident, Smt. S.C.L. Hospital, Smt. N.H.L. Municipal Medical College, Ahmedabad, Gujarat

**Dr. Raj Depani** 1<sup>st</sup> Year Surgery Resident, Smt. S.C.L. Hospital, Smt. N.H.L. Municipal Medical College, Ahmedabad, Gujarat

## ABSTRACT

**Introduction:** Gallstone disease is one of the common problem affecting the gastrointestinal system. Factors like obesity, sex, diet, age, ethnic background play a role in development of gallstone diseases. In this study we evaluated the common age group affected by gallstone and common presenting symptoms and biochemical nature of the stone.

**Methodology:** 100 patients of clearly documented cases of Gallstone diseases admitted in the surgical units of Smt. S.C.L. Hospital, between June 2017 to June 2019 constitute the material of this study.

**Results:** The 100 patients of gallstone diseases studied, ranged between 22 and 82 years of age. The mean age being 46.5 years. Right hypochondrial pain with epigastric pain was the presenting complaints in majority (30) cases. 22 cases presented with right hypochondrial pain alone and 20 cases had nausea and vomiting along with it. 6(6%) cases had pure cholesterol stones, 54(54%) cases had phosphate containing stones and 40 (40%) had oxalate containing stones.

**Conclusion:** Most of the cases of calculous cholecystitis presented in the fourth decade with females being more commonly affected. Pain in abdomen is the single most important symptom of gallstone disease. We found that pure cholesterol stones are less common.

## KEYWORDS

bile, cholesterol, cholecystitis, gallstone, infection

## I. INTRODUCTION:

Gall stones are one of the most common problems affecting the gastrointestinal tract. There has been a marked rise in the incidence of gallstones in the western countries. The past few decades have shown a surge in the number of Indian population presenting with the same. Obesity, pregnancy, dietary factors, age, gender and ethnic background are all associated with increased risk of developing gall stone disease. Women are more likely to develop gall stones than men.

## 1.1 Common Anomalies and Variations:

Absent gall bladder – extremely rare, autopsy incidence of 0.03% have been reported.<sup>[2]</sup> Double gall bladder, duplication of gall bladder with two separate cavities and two separate cystic ducts has an incidence of approximately 1 in 4000. Pathological process such as cholelithiasis and cholecystitis may involve one organ while the other is spared.<sup>[1]</sup>

Variation in size and shape of gall bladder a. Bilobed gall bladder, b. Fundal diverticulum, c. Phrygian cap, d. Hour glass gall bladder.

## 1.2 Physiology:

The gallbladder stores and concentrates hepatic bile in fasting state. The bile is delivered into the duodenum in response to a meal. The absorptive capacity per unit area of the gallbladder mucosa is greatest compared to any other structure in the body. This allows the bile to be concentrated by 5 to 10 fold by absorption of water and electrolytes which changes the bile composition<sup>[3,4]</sup>. Active NaCl transport by the gallbladder epithelium plays a key role in the concentration of bile. The solute absorption generates an osmotic force which causes passive absorption of water. The concentration of bile affects the solubility of calcium and cholesterol – the two important components of gallstones. The absorption of calcium is not as efficient as the absorption of sodium or water, this causes an increase in sodium concentration. The concentration of bile increases the tendency for cholesterol nucleation. Most patients remain asymptomatic throughout their life and may only be detected incidentally on ultrasound and CT scan. Nearly 3% of these individuals become symptomatic (biliary colic) per year. Of which, 3 to 5% of them develop complications per year.

Laparoscopic cholecystectomy has now become the standard of care in

treatment of gall stone disease. However, the timing of cholecystectomy, its relevance to complications and use of prophylactic antibiotics has always been a subject of debate.

## 1.3 Etio pathogenesis Of Gallstones:

Our knowledge of the etiopathogenesis of gall stone formation remains incomplete although there has been considerable progress on the mechanism involved during the past few decades.

## 1.4 Bile Concentration:

Water is a major component of bile. Significant net water absorption occurs during bile transfer through the bile ducts and during prolonged storage in the gallbladder. As a result, bile water content decreases from 97% weight in the bile ducts to 90% weight in the gallbladder. This threefold to fourfold concentration of bile enhances cholesterol crystallization and gallstone formation considerably.<sup>[5]</sup> The accepted current classification recognizes three types of gallstones; cholesterol, black pigment and brown pigment stones.<sup>[7]</sup>

## 1.5 Cholesterol Stones:

Pure cholesterol stones account for less than 10% of all stones and occur as single large stones with smooth surfaces, though they contain variable amount of bile pigment and calcium they are always more than 70% cholesterol by weight. Most cholesterol stones are radio lucent and less than 10% are radio opaque. The primary event in formation of cholesterol stones is supersaturation of bile with cholesterol. Cholesterol gallstones do not commonly harbour bacteria and are not associated with infected bile.<sup>[3]</sup>

## 1.6 Black Pigment Stones:

They account for 25% of stones in the west, but more prevalent in Asian countries. They are composed of bilirubin polymers without calcium palmitate and varying amounts of cholesterol (3-25%) and a matrix of organic material. They are associated with infections in 20% of patients. They are multiple, small, irregular and dark green to black in colour. Most of them occur secondary to hemolytic disorders, such as hereditary spherocytosis and sickle cell disease, and in those with cirrhosis.

### 1.7 Brown Pigment Stones:

They are less than 1 cm in diameter, brownish yellow and soft. They are often associated with infections of the biliary tract. They contain calcium bilirubinate, calcium palmitate and only small amounts of cholesterol bound in a matrix of organic material.

### 1.8 Etiology:

There is increased prevalence of gall stones in females and frequency of gallstones increases with age in both the sex. Certain risk factors increase the prevalence of gallstones and some induce symptomatic disease in patients. Female sex, Obesity, Age, Genetic and ethnic factors, Highly refined, fibre depleted, high animal fat diet, Diabetes Mellitus, Ileal disease and resection, Haemolytic states, Infections of biliary tract, Parasitic infections, Cirrhosis, Cystic fibrosis.

### Indications:<sup>[8]</sup>

Symptomatic Cholelithiasis, Acute or chronic cholecystitis – with or without stones, gall bladder polyps >1 cm size, Gall bladder carcinoma, Torsion of gall bladder, Traumatic rupture of gall bladder or cystic duct, Biliary peritonitis – with or without demonstrable perforation, Internal biliary fistula, Gas in the gall bladder, Non functioning gall bladder. Our Aim of this study is to evaluate the age and sex incidence of gall stone disease, to illustrate various types of clinical presentation in calculous cholecystitis, to study the various modes of management and their results. To study the bacteriology of the bile in calculous gall bladder disease and treatment outcome and biochemical type of gallstones.

### II. Materials And Methods:

100 patients of clearly documented cases of Gallstone diseases admitted in the surgical units of SMT. SCL Hospital between June 2017 to June 2019 constitute the material of this study. A detailed History was elicited in all patients and thorough clinical examination was done in them. After relevant pre-operative investigations and pre anesthetic check-up, cases were operated as laparoscopic cholecystectomy. Preoperative antibiotics were given. The operative findings and postoperative complications were recorded and carefully analyzed. The gallstones were sent for biochemical analysis and the gallbladder for histopathological examination. All patients received antibiotics and routine post-operative care.

Patient was properly examined in the post-operative period to note the development in any complication. Antibiotics were given and subsequently changed according to the bile culture and sensitivity report. In the follow up period attention were given to subject to improvement of the patients with regard to symptoms.

### III. Results:

The 100 patients of gallstone diseases studied ranged between 22 and 82 years of age. The mean age being 46.5 years. The maximum number of cases occurred in the third and fourth decades. Of the 100 cases, 66 (66%) were females, making the female: male ratio 1.9:1. Right hypochondrial pain with epigastric pain was the presenting complaints in majority (30) cases. 22 cases presented with right hypochondrial pain alone and 20 cases had nausea and vomiting along with it. There were 2 cases presenting with nausea and vomiting and 4 cases with epigastric pain alone but, 4 cases with a combination of both. However, right lumbar pain was the only presenting feature in a patient but it was associated with right hypochondrial pain and nausea and vomiting in 10 cases and 2 case respectively. 2 patients presented with right hypochondrial pain, epigastric pain and nausea and vomiting and we had 2 cases presenting with right hypochondrial and lumbar pain with nausea and vomiting. 8 of our cases had jaundice and 2 of them were found to have a CBD calculus. Murphys sign was positive in 36 cases. Fever was also a presenting feature in all 10 cases that were operated as an emergency. 22 of our cases were diabetic and 12 of them hypertensive of which 8 of them had both. 2 patient had bronchial asthma.

### 3.2 Elective Interval:

Many cases of acute cholecystitis were managed conservatively in the initial periods and taken up for surgery after intervals of one month to six months. Majority of the cases were taken up for surgery after an interval of about 1 week.

### 3.3 Management:

All the 100 cases were operated. Cholecystectomy alone was done in 92 cases. Out of the other 8 cases, 6 cases were combined with

appendectomy, and 2 cases were combined with CBD exploration. Of the 100, 96(96%) of them were done laparoscopically and 4(4%) as open procedure. Out of these 4, 2 of the cases were converted from a laparoscopic to open procedure due to severe adhesions (conversion rate 2%), and 2 cases were done for CBD exploration. All the laparoscopic procedures were done through the standard four port.

### 3.4 Stone Analysis:

6(6%) cases had pure cholesterol stones, 54(54%) cases had phosphate containing stones and 40 (40%) had oxalate containing stones.

### III. Discussion:

N. Teckchandani et al. in a comparable study to our series reported a mean age of incidence of 37.74 years (range 18- 65 years). A wide range (22-85 years) is observed in our series also with an average of 46.5 years. Though several studies have reported a higher incidence of gall stone disease in females, Jaime et al.(2010) in a retrospective review of acute cholecystitis have studied more male population than females in an advanced age group (72.15±17.46 years).<sup>[6]</sup> Pain was a presenting feature in most cases in our series except in 2 of the cases who presented as gangrenous cholecystitis presented in a state of drowsiness and had fever, and vomiting. In a majority of the cases pain was in right hypochondrium and epigastric region(30%). 22 % cases had right hypochondrial pain alone and 4% cases had epigastric pain alone.

As most cases were taken after an elective interval for cholecystectomy, fever was not a presenting feature in that. Nevertheless, all cases taken up as emergency had fever. 8 of our cases had jaundice, 6 of them were assumed to be reactive jaundice on the basis of its mild nature, biochemical investigations and further imaging. 2 of the cases were found to have a CBD stone and elevated serum alkaline phosphatase levels and needed CBD exploration. 22 of our cases had diabetes mellitus and itself was an indication for surgery in patients with cholelithiasis.




Most cases in this series underwent laparoscopic or open cholecystectomy after an elective interval from the time of initial attack. Only 20% of the patients had undergone surgery during the initial period. Patients on longer intervals had become symptomatic and two of them needed read missions. 48% of the cases who were operated after 3 months gave history of recurrent symptoms. This was of statistical significance (p=0.017).

Though the operative difficulty was more in cases operated in the initial periods, this did not find statistical significance in our study. However, complications rates were lesser in patients who were operated in the initial period and had operative difficulty. Jaime et al.(2010) concluded that significant proportion of conservative treatment was carried out at the expense of emergency surgery, although in absolute numbers conservative treatment seems to have a higher rate of complications, mortality and hospitalization time.<sup>[6]</sup> C. Skouras et al. after a review of 92 papers on the timing of cholecystectomy concludes that there is strong evidence that early laparoscopic cholecystectomy for acute cholecystitis offers an advantage in the length of hospital stay without increasing the morbidity or mortality.<sup>[8]</sup> Our study also concludes that early intervention reduces complications and length of hospital stay. Our study shows a statistical significance between the positive bile culture and post operative complications. D. Fuks et al.(2013) suggests that in all cases, but especially in severe cases, a sample of bile for bacteriologic study should be obtained.<sup>[9]</sup> Chi square= 12.09, DF=5, P value= 0.03364. The p value is less than 0.05, hence there is a statistical significance between the positive bile culture and post operative complications.

Based on the culture and spectrum of antibiotic sensitivity, 14(14%) cases needed an antibiotic change, whereas, 38(38%) cases were continued on the empirical antibiotic. Of the 14 cases which needed an antibiotic change, 12 of them were open cases among which, 10 had operative difficulty.

2 of the laparoscopic cholecystectomy cases were started on oral antibiotic for a stitch abscess. It was observed that most of the antibiotic changes were needed for open procedures and could be attributed to the post operative complications, operative difficulty or prolonged hospital stay. Nevertheless, D. Fuks et al. reports that

continuation of antibiotic treatment after early cholecystectomy does not seem necessary except in severe cases of acute calculous cholecystitis. This is further supported by Jay Narayan Shah et al(2012) who concluded that already low risk of wound infection following laparoscopic cholecystectomy was not significantly reduced further with the routine use of pre operative antibiotic prophylaxis in uncomplicated patients.



**Types of stone and their frequency**

**IV. Conclusion:**

In our study we have found that mean age of gallstone disease is 46.5 years and the female to male ratio is 1.9:1. Right hypochondrial and epigastric pain is the most common presenting feature with ultrasonogram alone being adequate to diagnose gallstone disease in 80% of cases. Early surgical intervention in cholecystitis reduces complications and length of hospital stay. Bacterial growth in bile can significantly increase the rate of post-operative complications. Pure cholesterol stones are less common. Pure cholesterol stones were not associated with any infected bile.

**REFERENCES:**

- [1]. Mincev M. Bilocular gallbladder. *Orvoskepzes* 1967; 42: 286-98.
- [2]. Olsen DO. Laparoscopic Cholecystectomy In: MacJadyer BV eds. *Laparoscopic surgery of abdomen*. New York: Springer Publication; Laparoscopic surgery of abdomen. New York: Springer Publication;
- [3]. Oddsdottir M, Phaas TH, Hunter JG. Gallbladder and extrahepatic biliarysystem. In: Brucardi CF Eds. *Schwartz's Principles of Surgery*. 9th ed.USA: McGraw Hill Companies Inc; 2009, p. 1135-66.
- [4]. Chari RS, Shah SA. Biliary system In: Townsend CM Eds. *SabistonTextbook of Surgery*. 18th ed. India: Elsevier Publication; 2009, p. 1517-88.
- [5]. van Erpecum KJ. Biliary lipids, water and cholesterol gallstones. *Biol Cell* 2005; 97(11):815–22
- [6]. PurificaciónCaleroGarcía, Jaime Ruiz Tovar, Alfonso SanjuanbenitoDehesa, Alicia CaleroAmaro, MaríaDiezTabernilla, Raquel LatorreFragua, GadaHousari Martín, Enrique Martínez Molina, VirgilioFresneda Moreno. 'Acute cholecystitis: is it still justified to delay surgery?'. *CIR ESP*. 2010;88(2):92–96
- [7]. De U. Evolution of cholecystectomy: A tribute to Carl August Langenbuch. *Indian J Surg* 2004;66:97-100.
- [8]. Palanivelu C. Laparoscopic Cholecystectomy. In: Palanivelu C, editor. *CIGES Atlas of Laparoscopic Surgery*. 2ndedn. New Delhi, India: Vij JP; 2003, p.39.
- [9]. D. Fuks, C. Cossé, J.-M. Régimbeau. 'Antibiotic therapy in acute calculouscholecystitis'. *Journal of Visceral Surgery* (2013) 150, 3—8.