



EFFECT OF OBESITY ON SYMPATHETIC NERVOUS SYSTEM

Physiology

Dr Vinita Ailani	Professor, Department of Physiology, National Institute of Medical Science and Research, Jaipur.
Dr Shikha Saxena*	Assistant Professor, Department of Physiology, National Institute of Medical Science and Research, Jaipur. *Corresponding Author
Mudasir Ahmad Wani	Tutor, Department of Physiology, Tahira Khanam Institute of Paramedical Sciences, Srinagar, Kashmir.

ABSTRACT

Obesity is recognized as a major, worldwide, health problem. Excess weight greatly increases the risk for cardiovascular diseases, diabetes and renal diseases. The present study was conducted in the Department of Physiology, at National Institute of Medical Sciences and Research, Jaipur. 50 obese (25 male and 25 female) and 50 non obese (25 male and 25 female) were included. Sympathetic tests were performed on both groups. All data are expressed as mean± standard deviation. Comparison between groups obese male, non obese male, obese female and non obese female was performed using the student's t-test (t-test for two independent samples / Two-tailed test). Differences was considered significance at $p < 0.05$. It was observed that young obese adults had reduced sympathetic activity. This altered balance of the autonomic nervous system increases the risk of cardiovascular disorders.

KEYWORDS

Cardiovascular diseases and obesity, Obesity, Sympathetic system and obesity

INTRODUCTION

Obesity is a nutritional health problem which is gradually increasing and affecting all population in children, adolescents and adults. It is a vexing problem in the developed economies [1]. According to WHO (World Health Organization [2] 1.9 billion people are overweight and 650 million are obese worldwide. Obesity can be seen as a first wave of a defined cluster of non-communicable diseases called "New World Syndrome" [3] that creates an enormous socioeconomic and public health burden in poorer countries also.

Nutritional population in India is gradually shifting from undernourishment to obesity [4]. Obesity is a condition, which has evolved with the advent of civilization, sedentary life style and high calorie diet [5]. Obesity is also on the rise in our society due to socioeconomic developments leading to change in lifestyle particularly dietary patterns [6].

The obesity can be measured by Body Mass Index (BMI), a person's weight (in kilogram) divided by the square of his or her height (in meters) (QUETELET'S INDEX) [7]. A person with a BMI of more than 25 Kg/m² is considered obese. Other approaches to quantify obesity include [8]: Anthropometry (Skin fold thickness), Densitometry (Underwater weighing), CT OR MRI, and Electrical impedance.

Many disorders occur in obese people with greater frequency. The most important and common of these disorders are hypertension, type 2 diabetes mellitus, dyslipidemia, coronary artery disease, degenerative joint disease and psychosocial disability [9].

Approximately 60% of individuals with obesity are suffering from the metabolic syndrome (including elevated abdominal circumference, high blood pressure, increased blood triglycerides, raised fasting blood sugar and increase in low density protein). Obese patients also have a greater risk of pulmonary functional impairment including sleep apnea, endocrine abnormalities and proteinuria [10].

Obesity increases cardiovascular morbidity and mortality risk through various mechanisms [11]. Various studies have shown that salt intake, blood volume and cardiac output are higher in obese persons than normal subjects [12].

The present study is an attempt to determine sympathetic autonomic functions in young obese. Early establishment of this correlation will help in preventing future cardiac autonomic disturbances like congestive heart failure, coronary artery disease, hypertension etc.

MATERIAL & METHODOLOGY

The present study was conducted among the students of National Institute of Medical Sciences and Research (Jaipur), NIMS Dental College and NIMS College of Paramedical technology at NIMS Jaipur. It is a comparative type of study that effect of obesity on 100 young obese and non-obese adults between age group 18-26 years divided into 4 groups on the basis of obesity and gender.

1. First group, Group 'A' 25 obese boys [OB+ Boys]
2. Second group, Group 'B' 25 obese girls [OB- Boys]
3. Third group, Group 'C' 25 Non-obese boys [OB+ Girls]
4. Fourth group, Group 'D' 25 Non-obese girls [OB- Girls]

Inclusion criteria: Consent participants; Students within the age group of 18-26 years; both gender were included.

Exclusion criteria: Subjects who were not interested; cases of diabetes mellitus, hypertension, cardiovascular or Endocrine disorders; on any drug that may alter autonomic function of their body; chronic smokers and chronic alcoholics.

Tests of Sympathetic Function:

I Orthostatic Test:

Blood pressure response was measured from supine to standing posture. A blood pressure recording of the subject was first taken in supine position. Then immediately the subject was asked to stand for atleast 3 minutes thereafter two recordings of blood pressure were taken 1st after 1 min and 2nd at about 2 to 3 mins. The difference in systolic and diastolic pressure at 1st min and 3rd min of standing from that of lying (baseline) pressure was calculated. A decrease in systolic blood pressure by atleast 20 mmHg or in diastolic blood pressure by atleast 10 mmHg within 3 mins of active standing was considered abnormal as defined by consensus of the American Society and the American Academy of Neurology.

II Cold Pressor Test:

The C.P.T consists of placing the hand into ice water which is a painful stimulus. It has been used to study the autonomic response of many individuals [13]. The afferent fibers (which are stimulated by placing the hand in cold water) and the efferent fibers are sympathetic fibers of the autonomic nervous system.

The subject was asked to immerse his/her hands in a container of ice water for one minute with temperature maintained at 1-4°C, throughout the test. Blood pressure was recorded before CPT, after 1 min of CPT and at 3 minutes after CPT. Difference between the blood pressure at three different stages of cold pressor test was recorded and observed [14].

Statistical Analysis:

Mean differences were tested for significance by Students unpaired 't' test. The statistical significance was assigned at p<0.05.

TESTS FOR SYMPATHETIC FUNCTION:

a) ORTHOSTATIC TOLERANCE TEST:

Table No. 1 (a): Comparison of blood pressure (SBP, DBP & MBP) by Student t test at 2 different phases of orthostatic test in Group A (obese boys) and B (non-obese boys).

	Group A (Obese Boys)	Group B (Non-Obese Boys)	p-value
Resting BP (mmHg)	Mean (±)SD	Mean (±)SD	
SBP	128.08 ± 3.71	118.9 ± 4.51	< 0.001
DBP	84.16 ± 3.64	80.16 ± 4.2	<0.05
MBP	98.42 ± 3.14	93.06 ± 3.56	<0.001
BP after standing			
SBP	121.28 ± 3.55	109.2 ± 4.39	< 0.001
DBP	84.32 ± 3.54	82.72 ± 3.95	0.138
MBP	96.92 ± 2.83	91.54 ± 3.27	< 0.001

Table No. 1 (b): Comparison of mean values of blood pressure (SBP, DBP & MBP) by Student t test at 2 different phases of orthostatic test in Group C (obese girls) & D (non-obese girls).

	Group C (obese girls)	Group D (non-obese girls)	p-value
Resting BP (mmHg)	Mean (±)SD	Mean (±)SD	
SBP	130.08± 3.53	130.08± 3.53	< 0.001
DBP	83.36 ± 2.75	79.28 ± 3.20	<0.05
MBP	98.93 ± 2.58	93.06 ± 3.47	< 0.001
BP after standing			
SBP	123.04 ± 3.42	111.36 ± 4.78	< 0.001
DBP	84.56 ± 2.48	81.52 ± 3.33	< 0.001
MBP	97.17 ± 2.57	91.46 ± 3.49	< 0.001


Table No. 1 (c): Comparison of mean values of BP by Student t test at 2 different phases of orthostatic test in Group A (obese boys) & C (obese girls)

	Group A (Obese Boys)	Group C (Obese Girls)	p-value
Resting BP (mmHg)	Mean (±)SD	Mean (±)SD	
SBP	128.08 ± 3.71	130.08± 3.53	0.057
DBP	84.16 ± 3.64	83.36 ± 2.75	0.933
MBP	98.42 ± 3.14	98.93 ± 2.58	0.409
Standing BP			
SBP	121.28 ± 3.55	123.04 ± 3.42	0.081
DBP	84.32 ± 3.54	84.56 ± 2.48	0.783
MBP	96.92 ± 2.83	97.17 ± 2.57	0.755

Table No. 1 (d): Comparison of mean values of BP by Student t test at 2 different phases of orthostatic test in Group B (non-obese boys) & D (non-obese girls)

	Group B (Non-Obese Boys)	Group D (Non-Obese Girls)	p-value
Resting BP (mmHg)	Mean (±)SD	Mean (±)SD	
SBP	118.9 ± 4.51	120.64 ± 4.49	0.174
DBP	80.16 ± 4.2	79.28 ± 3.20	0.247
MBP	93.06 ± 3.56	93.06 ± 3.47	0.791
Standing BP			
SBP	109.2 ± 4.39	111.36 ± 4.78	0.103
DBP	82.72 ± 3.95	81.52 ± 3.33	0.251
MBP	91.54 ± 3.27	91.46 ± 3.49	0.931

Figure 1: Comparison of mean values of BP at 2 different phases of orthostatic test in Group A (obese boys), B (non-obese boys), C (obese girls) & D (non-obese girls).



COLD PRESSOR TEST (CPT):

Table No. 2 (a): Comparison of mean values of blood pressure (SBP, DBP and MBP) by Student t test at 3 different phases of cold pressor test (CPT) in groups A (obese boys) and B (non-obese boys).

	Group A (obese boys)	Group B (non-obese boys)	p-value
Resting BP (mmHg)	Mean (±)SD	Mean (±)SD	
SBP	128.08 ± 3.71	118.9 ± 4.51	< 0.001
DBP	84.16 ± 3.64	80.16 ± 4.2	<0.05
MBP	98.42 ± 3.14	93.06 ± 3.56	< 0.001
BP after 1 min of CPT			
SBP	133.52 ± 3.33	126.8 ± 4.72	< 0.001
DBP	86.88 ± 3.46	86.08 ± 4.45	0.482
MBP	101.32 ± 3.35	99.65 ± 3.69	< 0.001
BP after 3 mins of CPT			
SBP	128.76 ± 3.33	120.88 ± 3.78	< 0.001
DBP	83 ± 3.43	81.36 ± 4.30	0.144
MBP	98.24 ± 3.03	94.53 ± 3.40	< 0.001

Table No. 2 (b): Comparison of mean values of blood pressure (SBP, DBP & MBP) by Student t test at 3 different phases of cold pressor test (CPT) in Group C (obese girls) & D (non-obese girls).

	Group C (obese girls)	Group D (non-obese girls)	p-value
Resting BP (mmHg)	Mean (±)SD	Mean (±)SD	
SBP	130.08± 3.53	120.64 ± 4.49	< 0.001
DBP	83.36 ± 2.75	79.28 ± 3.20	< 0.05
MBP	98.93 ± 2.58	93.06 ± 3.47	< 0.001
BP after 1 min of CPT			
SBP	135.84 ± 3.50	128.72 ± 4.31	< 0.001
DBP	88.08 ± 2.19	85.12 ± 3.41	< 0.001
MBP	103.83 ± 2.25	99.51 ± 3.80	< 0.001
BP after 3 mins of CPT			
SBP	131.04 ± 2.56	122.3 ± 4.15	< 0.001
DBP	84.16 ± 2.93	80.64 ± 2.75	< 0.001
MBP	99.78 ± 2.40	94.29 ± 2.72	< 0.001


Table No. 2 (c): Comparison of BP at 3 different phases of CPT by Student t test in groups A (obese boys) & C (obese girls)

	Group A (Obese Boys)	Group C (Obese Girls)	p-value
Resting BP (mmHg)	Mean (±)SD	Mean (±)SD	
SBP	128.08 ± 3.71	130.08± 3.53	0.057
DBP	84.16 ± 3.64	83.36 ± 2.75	0.933
MBP	98.42 ± 3.14	98.93 ± 2.58	0.247
After 1 min of CPT			
SBP	133.52 ± 3.33	135.84 ± 3.50	0.020
DBP	86.88 ± 3.46	88.08 ± 2.19	0.150
MBP	101.32 ± 3.35	103.83 ± 2.25	0.003
After 3 mins of CPT			
SBP	128.76 ± 3.33	131.04 ± 2.56	0.024
DBP	83 ± 3.43	84.16 ± 2.93	0.206
MBP	98.24 ± 3.03	99.78 ± 2.40	0.785

Table No. 2 (d): Comparison of BP at 3 different phases of CPT by Student t test in groups B (non-obese boys) & D (non-obese girls)

	Group B (Non-Obese Boys)	Group D (Non-Obese Girls)	p-value
Resting BP	Mean (±)SD	Mean (±)SD	
SBP	118.9 ± 4.51	120.64 ± 4.49	0.174
DBP	80.16 ± 4.2	79.28 ± 3.20	0.247
MBP	93.06 ± 3.56	93.06 ± 3.47	0.791
After 1 min of CPT			
SBP	126.8 ± 4.72	128.72 ± 4.31	0.140
DBP	86.08 ± 4.45	85.12 ± 3.41	0.397
MBP	99.65 ± 3.69	99.51 ± 3.80	0.900
After 3 mins of CPT			
SBP	120.88 ± 3.78	122.3 ± 4.15	0.206
DBP	81.36 ± 4.30	80.64 ± 2.75	0.485
MBP	94.53 ± 3.40	94.29 ± 2.72	0.785

Figure 2: Comparison of mean values of BP at 3 different phases of cold pressor test in Group A (obese boys), B (non-obese boys), C (obese girls) & D (non-obese girls).



DISCUSSION

The mean values of blood pressure response to standing from supine posture (Orthostatic Test) were found to be reduced in obese subjects as compared to the control subjects in present study. The mean values of blood pressure response to cold pressor test (CPT) were decreased in obese subjects as compared to control subjects.

The cardiovascular responses were measured as changes in systolic blood pressure (SBP), diastolic blood pressure (DBP), mean blood pressure (MBP) before, during and after the application of orthostatic tolerance test and cold pressor test in all the four groups (a, b, c & d).

a) Orthostatic Tolerance Test:

When comparison was done between mean values of SBP, DBP & MBP in Group A (obese boys) & B (non-obese boys), a highly significant difference was observed ($p < 0.001$) in SBP and MBP at 2 different phases of orthostatic test but a non-significant difference ($p > 0.05$) in DBP in standing position was observed between Group A (obese boys) & B (non-obese boys) (Table No. 1 (a) and Figure No 1).

When comparison was done between mean values of SBP, DBP & MBP in Group C (obese girls) & D (non-obese girls), a highly significant difference was observed ($p < 0.001$) in SBP, DBP & MBP at 2 different phases of orthostatic test between Group A (obese boys) & B (non-obese boys) (Table No. 1 (b) and figure No 1).

In the present study it was observed that the blood pressure response to standing from supine posture was significantly lower in obese subjects as compared to non-obese subjects. It indicates the decrease activity of the sympathetic nervous system.

Obese subjects was less responsive to blood pressure changes to standing posture, similar result were shown by [15, 16].

The result of the study was contradictory to present study where they observed orthostatic tolerance values were significantly increased in obese subjects [17].

b) Cold Pressor Test (CPT):

When comparison was done between mean values of SBP, DBP and MBP at resting, at 1 min of CPT & at 3 mins of CPT in Group A (obese boys) and B (non-obese boys), a significant difference ($p < 0.05$) was observed between SBP, DBP & MBP before CPT; between SBP after 1 minute of CPT and 3 minutes after CPT but a non-significant difference ($p > 0.05$) was observed between DBP and MBP at 1 minute of CPT and at 3 minutes of CPT in Group A (obese boys) & B (non-obese boys). (Table No 2 (a) and Figure No. 2)

When comparison was done between mean values of SBP, DBP and MBP at resting, at 1 min of CPT & at 3 mins of CPT in Group C (obese girls) & D (non-obese girls), the mean values of SBP, DBP & MBP showed a highly significant difference ($p < 0.001$) in Group C (obese girls) & D (non-obese girls). (Table No. 2 (b) and Figure No. 2)

The result of the present study showed the less increase of blood pressure response to cold pressor test in obese subjects as compared to the control subjects. A lesser increase in the blood pressure after the cold water immersion points towards sympathetic insufficiency in obese subjects [13].

The difference in mean systolic blood pressure recorded before and after cold pressor test was low in the study Group a (obese boys) compared to the control group. Obese subjects exhibit lower sympathetic response on exposure to cold. Our study results were in accordance with reported study of Monterio et al, 2012 [18].

Increased sympathetic activity induced by cold water stress causes release of nor-epinephrine and increase in blood pressure more in obese subject as compare to control, which is contradictory to our studies [19, 20].

Obesity is associated with both sympathetic and parasympathetic nervous system dysfunction [20]. Decrease in the sympathetic activity may result in a disordered homeostatic mechanism thus promoting excessive storage of energy [21].

CONCLUSION

Present study conclude that with irrespective of gender, reduced sympathetic activity predict increased risk for subsequent cardiac events in obese individuals. Further more studies are still needed in future to explore and clinically help in reduce the risks of various diseases due to obesity.

REFERENCES

- Lau DC, Douketis JD, Morrison KM, Hramiak IM, Sharma AM. 2006 Canadian clinical practice guidelines on the management and prevention of obesity in adults and children [summary]. Canadian Medical Association Journal 2007; 176(8): 1-13.
- World Health Organization: Global health risks: mortality and burden of disease attributable to selected major risks, 2009.
- Kapil D, Chaudhary AK. New world syndrome (obesity) gone by guggul: a review. Journal of Ayurveda and Holistic Medicine 2015; 2(9): 31-35.
- Nageswari KS, Sharma R, Kohli DR. Assessment of respiratory and sympathetic cardiovascular parameters in obese school children. Indian journal of physiology and pharmacology 2007; 51(3): 235.
- Youngee K, Suh YK, Choi H. BMI and metabolic disorders in South Korean adults: 1998 Korea national health and nutrition survey. Obesity research 2004; 12(3): 445-453.
- Chhatwal J, Verma M, Riar SK. Obesity among pre-adolescent and adolescents of a developing country (India). Asia Pacific journal of clinical nutrition 2004; 13(3):23-25.
- Garrow JS, Webster J. Quetelet's index (W/H2) as a measure of fatness. International journal of obesity 1985; 9(2): 147-153.
- Wells JC, Fewtrell MS. Measuring body composition. Archives of disease in childhood 2006; 91(7):612-7.
- Aviva M, Spadano J, Coakley EH, Field AE, Colditz G, Dietz WH. The disease burden associated with overweight and obesity. JAMA 1999; 282(16): 1523-1529.
- Samuel K, Burke LE, Bray GA, Blair S, Allison DB, Pi-Sunyer X, Eckel RH. Clinical implications of obesity with specific focus on cardiovascular disease: a statement for professionals from the American Heart Association Council on Nutrition, Physical Activity, and Metabolism: endorsed by the American College of Cardiology Foundation. Circulation 2004; 110(18): 2952-2967.
- Sangun Ö, Dündar B , Köşker M, Pirgon, Ö, Dündar N. Prevalence of metabolic syndrome in obese children and adolescents using three different criteria and evaluation of risk factors. Journal of clinical research in pediatric endocrinology 2011; 3(2): 70-73
- Messeri FH, Christie B, DeCarvalho JG, Aristimuno GG, Suarez DH, Dreslinski GR, Frohlich ED. Obesity and essential hypertension: hemodynamics, intravascular volume, sodium excretion, and plasma renin activity. Archives of internal medicine 1981; 141(1):81-85
- LeBlanc J, Dulac S, Cote J, Girard B. Autonomic nervous system and adaptation to cold in man. Journal of Applied Physiology 1975; 39(2): 181-186.
- Kilgour RD, Carvalho J. Gender differences in cardiovascular responses to the cold hand pressor test and facial cooling. Canadian journal of physiology and pharmacology 1994; 72(10): 1193-1199.
- Emdin M, Gastaldelli A, Muscelli E, Macerata A, Natali A, Camastra S, Ferrannini E. Hyperinsulinemia and autonomic nervous system dysfunction in obesity. Circulation 2001; 103: 513-519.
- Borne P, Hausberg M, Hoffman RP, Mark AL, Anderson EA. Hyperinsulinemia produces cardiac vagal withdrawal and nonuniform sympathetic activation in normal subjects. American Journal of Physiology-Regulatory, Integrative and Comparative Physiology 1999; 276(1):178-R183.
- Chaudhuri A, Borade NG, Tirumalai J, Saldanha D, Ghosh B, Srivastava K. A study of autonomic functions and obesity in postmenopausal women. Industrial psychiatry journal 2012; 21(1):39-41.
- Monteiro G, Chathoth VM, Kador K. Cardiac autonomic response during a cold pressor test in normal and overweight adults. Int J Biomed Adv Res 2012; 3(6): 514-516.
- Velasco M, Gómez J, Blanco M, Rodríguez I. The cold pressor test: pharmacological and therapeutic aspects. American journal of therapeutics 1997; 4(1): 34-38.
- Garg R, Malhotra V, Goel N, Dhar U, Tripathi Y A study of autonomic function tests in obese people. International Journal of Medical Research and Health Sciences 2013; 2(4): 750-755.
- Bray GA. Integration of energy intake and expenditure in animals and man: the autonomic and adrenal hypothesis. Clinics in endocrinology and metabolism 1984; 13(3): 521-546.