



## CLINICAL STUDY OF SMALL BOWEL PERFORATION

## General Surgery

**Dr Mahesh M Pukar**

Professor, Department of General Surgery, Smt. B.K. Shah Medical Institute & Research Centre, Pipariya, Vadodara, Gujarat 391 760, India.

## ABSTRACT

**Background and Objectives:** Bowel perforation is most common cause of surgical emergency. Perforation of the small intestine causing peritonitis is the most common abdominal surgical emergency encountered. The objective of the study was to analyze the Clinico Pathological Relations of Small bowel perforations to better understand the condition. **Materials and Methods:** This study was prospective clinicopathological study starting from June 2018 to December 2018 carried out in Dhiraj General Hospital, Vadodara in General Surgery Department. 30 patients presenting with features of small bowel perforation were studied. **Conclusion:** Perforation commonly occurs in young males. Pain, abdominal distension along with Generalized tenderness was commonly found in all. Nonspecific inflammation was most common non traumatic cause of ileal perforation after blunt abdominal trauma. Stress, smoking and NSAIDS were common risk factors involved in causation of bowel perforation

## KEYWORDS

Small Bowel; Perforation; Tuberculosis.

## Introduction

Intestinal perforation is the third most common cause for explorative laparotomy as an emergency. Small Bowel Perforation is a perforation in the wall of small intestine anywhere starting from duodenum to ileum. It leads to spillage of contents of small bowel into Peritoneal cavity causing peritonitis. It is a very acute condition. It needs to be treated as early as possible.

Small bowel Perforations are common in occurrence. Upper bowel perforation can be either free or contained.

1. Free perforation Bowel contents spill freely into the abdominal cavity, causing diffuse peritonitis (eg. duodenal or gastric perforation).
2. Contained perforation free spillage is prevented as contiguous organs wall off the area of perforation (as when a duodenal ulcer penetrates into the pancreas).

Lower bowel perforation (eg. in patients with acute diverticulitis or acute appendicitis) results in free intraperitoneal contamination.

## Frequency

Small bowel injuries following blunt abdominal trauma are infrequent in children with an incidence of 17%. Peptic ulcer perforations are a common cause of morbidity and mortality with an acute abdomen in adults. Though the rate has fallen in parallel with the general decline in the prevalence of peptic ulcer disease. Duodenal ulcer perforations are 23 times more common than gastric ulcer perforations. About one third of gastric perforations are due to gastric carcinoma.

Approximately 1015% of patients with acute diverticulitis develop free perforation. The overall mortality rate is relatively high (2040%), largely because of complications, such as septic shock and multiorgan failure. In elderly patients, acute appendicitis has a mortality rate of 35% and a morbidity rate of 50%. A major contributing factor to morbidity and mortality in these patients is the presence coexisting medical conditions in elder patients. Bowel injuries associated with endoscopy are not a common cause of perforation.

As perforations related to endoscopic retrograde cholangiopancreatography (ERCP) occur in only about 1% of patients [1].

## Materials and Method

Thirty patients presenting with features of small bowel perforation were studied.

In all the patients complete detailed history was taken including age of the patient, signs and symptoms and their duration and relevant past history.

Complete physical examination including clinical and systemic examination was done. All the routine investigations were done. For pathological Analysis peritoneal fluid examination and Biopsy taken from the Perforation will be sent.

Correlation will be made between Pathological Reports and Clinical presentation of patients.

The outcome will be evaluated as per the predesigned Performa of study.

## Inclusion Criteria

1. All patients, who willing to give consent to participate in this study.
2. All patients operated for Small Bowel Perforations.

## Exclusion Criteria

1. Patients with multiple co morbidity.
2. Patient not willing for study.

Patients were evaluated for the causes of small bowel perforation based on various investigations, clinical presentation and pathological analysis.

Results and Discussion  
Sex and Age Distribution

In the present study nearly two third of the patients (67%) were males with male to female ratio of 2:1 suggesting male preponderance. Most of the patients presented in fourth decade of life with age 31 to 40 years (40%). Mean age was 45.56 years. Overall these findings suggest that, the frequency of small bowel perforation is high among males and are common in third and fourth decade of life. Similarly, Dr. Raja Gopala Rao Akireddy et al. [2] conducted a similar study of 100 patients, in which also male preponderance was noted with 68% of males and male to female ratio of 2.12:1. The commonest age group was 31 to 40 years (39%) and the mean age was 33.97±13.93 years. Most of the patients in our study came from lower socioeconomic class & tribal area. Most of them were uneducated and labourers. Males were usually workers and females were usually housewives. This denotes high levels of stress in males. They were unwary of their health. Most of the people remain empty stomach & doing hard work and for their tiredness they were using NSAIDS with their empty stomach suggesting cause for perforations due to peptic ulcers. Smoking was also major risk factor found in all the male patients studied. Mean age study of various study reveals that perforation commonly occurs in younger age group commonly in 3rd to 4th decade of life, with mean age of 45.56 years which is similar to Allsopp T. et al. study [3] while it was 29.36 in Ansari AG study [4]. As most of the patients are labourers and poorly educated younger population was mainly working as labourers in various places. Most of them were ignorant about their health and presented late. Stress was major triggering factor in men of age group between 30 to 40 years.

Clinical Presentation  
Site of Pain

In our study all the patients of small bowel perforation presented with abdominal pain. In cases of duodenal perforation pain was present over epigastric region in most of the patients. In Ileal and Jejunal

perforations pain was commonly generalised and present in right iliac fossa in rest of the cases. In study conducted by Dr. Sunil Sudharshan H [5] of clinicopathological evaluation, management and outcome of gastrointestinal perforations in 75 patient's similar results were found.

### **Clinical Features**

In this study all the patients presented with abdominal pain (100%). The next common complaint was vomiting 22 cases (73%), abdominal distension 21 cases (70%) and constipation 17 cases (56%). The other complaints were fever 7 cases (23%) and blunt abdominal trauma. Abdominal pain and distension was the commonest presenting symptoms reported in our patients and the same being reported by Shrivastava D et al. [6] in which they studied on 155 patients. All the patients were presented with fever and abdominal pain. 122 (78.70%) patients presented with abdominal distension. Pain was present in all cases due to peritoneal inflammation secondary to bowel perforation in all cases. Abdominal distension was present as sequel of dilated oedematous bowel loops secondary to inflammatory exudative fluid and paralytic ileus as consequences of protective mechanism for limitation of spread of peritonitis. Also, most of the patients presented late as they came from remote rural areas so abdominal distension was already set in.

### **Etiology of Small Bowel Perforation**

Small bowel perforation is a rare event that results from a number of different etiological causes. Causes include, trauma, infection, inflammatory bowel disease, foreign body ingestion, malignancy, iatrogenic causes and medical therapies for other diseases.

#### **• Pathology of Small Bowel Perforation a. Peritoneal Fluid Study**

In our study, most of the patient presented with purulent peritoneal fluid (53%) followed by greenish fluid (23%). Greenish fluid denotes presence of bile in duodenal perforation. Purulent fluid was present due to contamination of peritoneal cavity in cases which presented late for treatment. Dr. Rauf A Wani, Fazl Q Parry and Nadeem A Bhat et al. [7] conducted prospective study on 79 cases of nontraumatic terminal ileal perforation and similar findings were noted.

#### **b. Histopathological Examination**

In our study, on histopathological examination, majority of the patients had nonspecific inflammation 22 cases (73%) followed by acute inflammation 5 cases (17%), malignancy 2 cases (6%) and tuberculous infection 1 case (3%) as causes of bowel perforation. Dr. Raja Gopala Rao Akireddy et al. [2] conducted a similar study of 100 patients in which on histopathological examination, majority of the patients had (83%) chronic inflammatory changes followed by tuberculosis (12%) and malignancy (5%).

#### **c. Site of Perforation**

In the present study, with regard to site of perforation, more than half of the study population had ileal perforation 17 cases (57%) and in the remaining had duodenal 9 cases (30%) and jejunal perforation 4 cases (13%). However Seth S and Agrawal KK, conducted a study of 33 patients & found duodenum being the commonest site followed by ileum and jejunum in their study [8].

#### **d. Tuberculosis**

Coccolini F. et al. [10] studied 119 cases of abdominal TB and suggested ileum as common site of perforation in cases of tuberculosis. In our case study also tuberculous inflammation was found in ileum in one case. Intestinal tuberculosis commonly occurs after ingestion of infected sputum in cases of pulmonary tuberculosis. The ileocecal area and jejunum are the most common sites involved as there are high densities of lymphoid aggregates and physiologic stasis [10].

#### **e. Malignancy**

The different pathologic types of small bowel malignant tumors include adenocarcinoma, carcinoid tumors, leiomyosarcoma, and lymphoma. Adenocarcinoma is the most commonly encountered small bowel malignant tumor [11], which is true in case of our study. Adenocarcinoma is malignant tumour most commonly found in ileum. As lymphatics of small bowel are most numerous in ileum in form of Peyer's patches, malignant spread of cancer cells commonly occurs in ileum, which leads to submucosal enlargement of nodes and ulceration and perforation.

#### **f. Trauma**

In this study six cases (20%) of traumatic small bowel perforation

was noted. Blunt abdominal trauma was the cause in all the bowel perforations. Small bowel is typically compressed against a fixed point, usually the vertebral column. This rapid increase in intraluminal pressure leads to perforation of the bowel wall at the antimesenteric border, where the bowel is usually weaker. Laparoscopic seat belts, commonly used in most cars until the 1980s, were largely held responsible for perforation and SBP [12]. Greater length of small bowel particularly ileum and jejunum also predispose it for trauma along with its mobility.

### **Conclusion**

Perforation commonly occurs in young males. Pain, abdominal distension along with generalised tenderness was commonly found in all. Nonspecific inflammation was most common nontraumatic cause of ileal perforation after blunt abdominal trauma. Stress, smoking and NSAIDs were common risk factors involved in causation of bowel perforation

### **REFERENCES**

1. Stapfer M, Selby RR, Stain SC, et al. Management of duodenal perforation after endoscopic retrograde cholangiopancreatography and sphincterotomy. *Ann Surg.* 2000 Aug;232(2):1918.
2. Dr. Raja Gopala Rao Akireddy et al; Prospective Study of Patients Presenting With Small Bowel Perforations and Outcome in A Rural Hospital In South India; *IOSR Journal of Dental and Medical Sciences (IOSRJDMS)* 2016 Sep;15(9):0116.
3. Allsopp T, Carmody C, Mikhail M. Outcomes of Rurally Treated Small Bowel Perforations Compared to a Tertiary Centre. *The Internet Journal of Surgery* 2012;28(2).
4. Ansari AG, Nagvi SQH, Ghumrao AA, Jamali AH. Management of typhoid ileal perforations: a surgical experience of 44 cases. *Gomal J Med Sci.* 2009;7(1):2730.
5. Dr. Sunil Sudharshan H. Study of clinicopathological Evaluation, Management and Outcome of Gastro Intestinal Perforations; *rguhs*, pg 43.
6. Shrivastava D, Jain AK, Gharde P, Sharma DB, Verma RS. Typhoid intestinal perforation in Central India a surgical experience of 155 cases in resource limited setting. *IJBAR.* 2014;5(12):6004.
7. Rauf A Wani, Fazl Q Parry, Nadeem A Bhat, Mehmood A Wani, Tasaduq H Bhat, Fowzia Farzana, et al. Nontraumatic terminal ileal perforation. *World J Emerg Surg.* 2006; 1: 7
8. Seth S, Agrawal KK. A Review of 51 cases of Duodenal perforation in Rohilkhand region. *Int J Contemporary Med Res.* 2016;3(6):18068.
9. Coccolini F, Ansaloni L, Catena F, Lazzareschi D, Puviani L, Pinna AD. Tubercular bowel perforation: what to do? *Ulus Travma Acil Cerrahi Derg* 2011;17(1):6674.
10. Leung VK, Law ST, Lam CW, et al. Intestinal tuberculosis in a regional hospital in Hong Kong: a 10 year experience. *Hong Kong Med J.* 2006;12:264-271.
11. Kumar A, Patodia M, Sharda VK, Pandove PK, Bal MS. Primary Adenocarcinoma of ileum presenting as Perforation: A Review of literature. *JIMSA* 2013;26(4):2234.
12. Symeonidis N, Ballas K, Psarras K, Lalountas M, Rafailidis S, Pavlidis T, Sakantamis A. Isolated Small Bowel Perforation After Blunt Abdominal Trauma: Report Of 2 Cases. *Ann Ital Chir.* 2013 Nov Dec;84(6): 699703.