



A COMPARATIVE STUDY BETWEEN THE OUTCOME OF PRIMARY REPAIR VERSUS LOOP ILEOSTOMY IN ILEAL PERFORATION AT NMC, SASARAM, BIHAR

Surgery

Dr. Subhash Kumar*	M.B.B.S., M.S. (Surgery), FMAS, FIAGS, Associate Professor, Departments Of Surgery, Narayan Medical College, Jamuhar, Sasaram, Bihar. *Corresponding Author
Dr. N. K. Jha	M.B.B.S., M.S. (Surg.), Professor and Head of Department, Departments Of Surgery, Narayan Medical College, Jamuhar, Sasaram, Bihar.
Dr. Debarshi Jana	Young Scientist (DST), Department of Gynecology and Obstetrics, Institute of Post-Graduate Medical Education and Research, A.J.C. Bose Road, Kolkata-700020, West Bengal, India

ABSTRACT

INTRODUCTION. Ileal perforation peritonitis is a common surgical emergency in the Indian subcontinent and in tropical countries. It is reported to constitute the fifth common cause of abdominal emergencies due to high incidence of enteric fever and tuberculosis in these regions.

METHODS. Sixty proven cases of ileal perforation patients admitted to Surgical Emergency were taken up for emergency surgery. Randomisation was done by senior surgeons by picking up card from both the groups. The surgical management was done as primary repair (group A) and loop ileostomy (group B).

RESULTS. An increased rate of postoperative complications was seen in group A when compared with group B with 6 (20%) patients landed up in peritonitis secondary to leakage from primary repair requiring reoperation as compared to 2 (6.67%) in ileostomy closure. A ratio of 1 : 1.51 days was observed between hospital stay of group A to group B.

CONCLUSION. In cases of ileal perforation temporary defunctioning loop ileostomy plays an important role. We recommend that defunctioning ileostomy should be preferred over other surgical options in cases of ileal perforations. It should be recommended that ileostomy in these cases is only temporary and the extra cost and cost of management are not more than the price of life.

KEYWORDS

INTRODUCTION

Gastrointestinal perforations have been surgical problem since the time immortal. Scientists have found evidence of gastrointestinal perforations in Egyptian mummies. Perforation is said to occur once a pathology which extends through the full thickness of the hollow viscus leading to peritoneal contamination with intraluminal contents. Perforation can occur anywhere in the gastrointestinal tract starting from oesophagus to the rectum.

Ileal perforation peritonitis is a common surgical emergency in the Indian subcontinent and in tropical countries. It is reported to constitute the fifth common cause of abdominal emergencies due to high incidence of enteric fever and tuber-culosis in these regions. Despite the availability of modern diagnostic facilities and advances in treatment regimes, this disease has an abrupt onset and a rapid downhill course with a high mortality if not treated.

Various causes of nontraumatic ileal perforation include bacterial infections (salmonella, yersinia, and tuberculosis), viral infections (cytomegalovirus, human immunodeficiency virus), fungal infection (histoplasma), parasitic infections (A. lumbricoides, E. vermicularis, and E. histolytica), and others (Wagener's granulomatous and drugs (Nonsteroidal anti-inflammatory drugs, e.g., aspirin, paracetamol, mefenamic acid, Ibuprofen, etc.)). In a significant number of cases the cause of perforation is not known and it is called nonspecific ileal perforation. The perforation causes gram-negative aerobic and anaerobic infection leading to peritonitis.

Various operative procedures were advocated by different authors, such as the following:

- (i) simple primary repair of perforation;
- (ii) repair of perforation with ileotransverse colostomy
- (iii) primary ileostomy;
- (iv) single layer repair with an omental patch;
- (v) resection and anastomosis.

Even with such a variety of procedures, ileal perforation still has a high rate of morbidity and mortality. The aim of the present study is to evaluate the outcome of primary repair versus loop ileostomy in cases of ileal perforation by comparing them in terms of postoperative morbidity, mortality and cost-effectiveness, and complications and to find out the ideal procedure. The study will help to establish the criteria for instituting the management modality according to presentation and

severity of the disease and the outcome of these procedures. Effective management of the disease will help in decreasing morbidity and mortality associated with the disease.

MATERIAL AND METHOD

This comparative study was conducted in the Department of Surgery, NMC, Sasaram, Bihar. Sixty patients admitted to Surgical Emergency with acute abdomen were selected for the study. There were not any preoperative selection criteria; the cases which were proven to be cases of perforation peritonitis on the basis of investigations and clinical examination were taken for study and considered for comparative study if laparotomy diagnosed to be case of ileal perforation. These patients were taken up for emergency surgery after resuscitation, and an informed consent was taken. The antibiotics were given in both groups after admission to hospital and before surgery with 3rd generation cephalosporin (cefotaxime, ceftazidime, ceftriaxone, etc.) and metronidazole. These patients were divided into two groups group A and group B. Randomisation was done by senior surgeons by picking up card from both the groups. The surgical management was done as primary repair (group A) and loop ileostomy (group B); comparative study was done between both procedures. All operations were done by group of three experienced surgeons and they all performed the same technique. All the procedures were carried with hand sewn method. In group A primary closure was done in two layers, the inner layer closed with 3-0 poly glycolic acid (vicryl) and outer layer closed with silk 3-0. In group B loop ileostomy was done. Postoperative complications in each group like wound infection, wound dehiscence, intraabdominal abscess, stricture of anastomosis site, faecal fistula, peritonitis, septicemia, ileostomy related complications, paralytic ileus, intestinal obstruction and death and so forth are evaluated.

RESULTS

During the 12-month period of study, 60 patients with ileal perforation were studied. Ileal perforations were most commonly observed in third and fourth decade of life with males more commonly affected (Male : Female: 6.5 : 1) (Table 1). Pain abdomen was the most common clinical presentation (100%) followed by fever, abdominal distension, vomiting, and obstipation. Time since perforation was within 12 hour in 3 cases, between 12 and 24 hour in 23 cases, between 24 and 48 hour in 18 cases, 48 and 72 hour in 7 cases, between 72 and 96 hour in 8 cases, and between 96 and 120 hour in 2 cases. Most of the patients (83.33%) presented within 72 hours of perforation and most cases

were operated within 12 hours of presentation after adequate resuscitation. The average duration of fever was 7.2 days whereas in patients with typhoid perforation, the average duration of fever was 10.41 days ranging from 1 day to 25 days. Fever preceded the abdominal symptoms in these patients. In all the cases in study group, the etiology of perforation was typhoid, circular perforation at antimesenteric border (36.67%), nonspecific (35%), tuberculosis, elliptical perforation at antimesenteric border (18.33%), and trauma (10%). There was not any case having malignant cause of ileal perforation. Complications: wound infection was the commonest complication (36.67%). It was present in about 11 (36.67%) cases each in patients having undergone primary repair of perforation and patients having undergone ileostomy. Ileostomy related complications occurred in 19 patients (63.33%). Peristomal skin excoriation was the most common ileostomy related complication in 10 patients (33.33%) followed by weight loss in 4 (13.33%), retraction in 4 (13.33%), fluid and electrolyte imbalance in 3 (10%), and prolapse in 1 (3.33%). Ileostomy closure related complications occurred in 7 patients (11.67%), wound infection in 6 (20%), anastomotic leak in 2 (6.67%), intra-abdominal collections in 2 (6.67%), wound dehiscence in 4 (13.33%), and reoperations in 2 (6.67%). The complications between the two groups were statistically significant with value 0.026, with chi square test² value 9.24 with degree of freedom Df 3. The average duration of hospital stay in patients having undergone primary closure was 14.3 days compared to 21.53 days in patients with ileostomy, which included ileostomy closure. The average duration of ileostomy before closure was 208.1 days (about 3.6 months). Only two patients having DM and outcome of these patients were good and remaining 58 patients were having no any comorbidities. In all the cases the biopsy was sent and histopathological examination was done and found to be typhoid enteritis 12 (20%), tubercular 6 (10%), and nonspecific inflammation in the remaining 42 (70%) cases.

Table 1: Age distribution in both the groups.

Age group (in years)	Group A		Group B	
	Number of cases	% age	Number of cases	% age
10–20	3	10	4	13.33
21–30	7	23.34	8	26.66
31–40	10	33.33	7	23.34
41–50	4	13.33	7	23.34
51–60	4	13.33	4	13.33
61–70	2	6.67	0	0
Total	30	100	30	100
Range	15–70		16–60	

CONCLUSION

Temporary defunctioning loop ileostomy in cases of ileal perforation plays an important role in reducing the incidence of complications like faecal fistula. This helps reduce mor-tality in patients undergoing surgery for ileal perforations. Ileostomy-specific complications, however, increase the post-operative stay of the patient. These complications can be reduced, if not outright eliminated, by proper fashioning of the stoma and provision of adequate nursing care of the stoma. We recommend that defunctioning loop ileostomy should be preferred over other surgical options in cases of ileal perforations in randomised study. It should be recom-mended that ileostomy in these cases is only temporary and the extra cost and cost of management are not more than the price of life.

REFERENCES

1. P. B. Prasad, D. K. Choudhury, and O. Prakash, "Typhoid perforation treated by closure and proximal side-to-side ileo-transverse colostomy," *Journal of the Indian Medical Association*, vol. 65, no. 11, pp. 297–299, 1975.
2. B. K. Kaul, "Operative management of typhoid perforation in children," *International Surgery*, vol. 60, no. 8, pp. 407–410, 1975.
3. K. P. Singh, K. Singh, and J. S. Kohli, "Choice of surgical procedure in typhoid perforation: experience in 42 cases," *Journal of the Indian Medical Association*, vol. 89, no. 9, pp. 255–256, 1991.
4. P. G. Purohit, "Surgical treatment of typhoid perforation. Experience of 1976 Sangli epidemic," *Indian Journal of Surgery*, vol. 40, no. 5, pp. 227–238, 1978.
5. C. G. Athie, C. B. Guizar, A. V. Alcantara, G. H. Alcaraz, and J. Montalvo, "Twenty-five years of experience in the surgical treatment of perforation of the ileum caused by Salmonella typhi at the General Hospital of Mexico City, Mexico," *Surgery*, vol. 123, no. 6, pp. 632–636, 1998.
6. A. M. Malik, A. A. Laghari, Q. Mallah et al., "Different surgical options and ileostomy in typhoid perforation," *World Journal of Medical Sciences*, vol. 1, pp. 112–116, 2006.
7. R. A. Wani, F. Q. Parray, N. A. Bhat, M. A. Wani, T. H. Bhat, and Farzana, "Nontraumatic terminal ileal perforation," *World Journal of Emergency Surgery*, vol. 24, no. 1, article 7, 2006.
8. A. R. K. Adesunkanmi, T. A. Badmus, F. O. Fadiora, and E. A. Agbakwuru, "Generalized peritonitis secondary to typhoid ileal perforation: assessment of severity using modified APACHE II score," *Indian Journal of Surgery*, vol. 67, no. 1, pp. 29–33, 2005.

8. S. Talwar, R. K. Sharma, D. K. Mittal, and P. Prasad, "Typhoid enteric perforation," *Australian and New Zealand Journal of Surgery*, vol. 67, no. 6, pp. 351–353, 1997.
9. U. Beniwal, D. Jindal, J. Sharma, S. Jain, and G. Shyam, "Com-parative study of operative procedures in typhoid perforation," *Indian Journal of Surgery*, vol. 65, no. 2, pp. 172–177, 2003.
10. R. Bakx, O. R. C. Busch, W. A. Bemelman, G. J. Veldink, J. F. M. Slors, and J. J. B. van Lanschoot, "Morbidity of temporary loop ileostomies," *Digestive Surgery*, vol. 21, no. 4, pp. 277–281, 2004.
11. O. Perez, A. Habr-Gama, V. E. Seid et al., "Loop ileostomy morbidity: timing of closure matters," *Diseases of the Colon and Rectum*, vol. 49, no. 10, pp. 1539–1545, 2006.