



A STUDY ON THE RATE OF OCCURRENCE OF *H. PYLORI* IN DYSPEPTIC PATIENTS IN A TERTIARY CARE HOSPITAL

Pathology

Dr. Tarak Banik Assistant Professor, Malda Medical College, West Bengal, India.

Dr. Prasenjit Kumar Bar* Assistant Professor, Malda Medical College, West Bengal, India. *Corresponding Author

Dr Saikat Mandal Demonstrator, Malda Medical College, West Bengal, India.

Dr. Subhrajyoti Naskar Assistant Professor, Malda Medical College, West Bengal, India.

ABSTRACT

INTRODUCTION: *Helicobacter pylori* (*H. pylori*) chronically infect more than half of all people worldwide. We explore the rate of occurrence of *H. pylori* in dyspeptic patients and its causal association with peptic ulcer disease, non-autoimmune gastritis, non peptic ulcer dyspepsia and gastric carcinoma.

OBJECTIVES: To find out the rate of occurrence of *H. pylori* in dyspeptic patients and to emphasize the diagnosis of *H. pylori* by rapid urease test, serology, Giemsa stain of biopsy specimens and to correlate all these diagnostic tests with biopsy.

METHODS: An observational prospective cohort study was performed on 60 selected cases. Detail clinical history, laboratory tests and histopathological examination were done. Findings were recorded and analyzed statistically.

RESULTS: The comparison of patient's age group, socio-economic status and association of *H. pylori* are statistically significant. Dyspepsia was more common in male (68.3%) and the sex ratio was 2.15:1. The rate of occurrence of *H. pylori* in dyspeptic patients was 68.33% and for functional dyspepsia was 61.29%. Majority of dyspeptic patients were positive for IgG ELISA test, 65% were positive for the biopsy urease test and 55% in modified Giemsa stain for *H. pylori* in tissue sections. Incidence of chronic superficial gastritis, benign gastric ulcer and duodenitis/duodenal ulceration were noted and association with *H. pylori* was statistically significant.

CONCLUSION: Early diagnosis may prevent dyspepsia and complications of *H. pylori* infection like chronic superficial gastritis, benign gastric ulcer, duodenitis/duodenal ulceration even adenocarcinoma.

KEYWORDS

Dyspepsia, *H. Pylori*, Endoscopy, Histopathology

INTRODUCTION:

Dyspepsia is a symptom complex which is characterized by pain, discomfort or nausea referable to upper alimentary tract, which may be intermittent or continuous in nature. Dyspepsia is a major health problem; prevalence ranges from 20-30% in general population [1]. Dyspepsia is mainly divided into two main groups- functional dyspepsia, accounting for 60% of total cases and dyspepsia with organic causes [2]. *H. pylori* (formerly *Campylobacter pylori*) is a gram negative, flagellated, microaerophilic, spiral organism that chronically infects more than half of all people worldwide. In developing countries like India, 70-90% of the population carries *H. pylori*; almost all of these acquire infection before the age of 10 years [3, 4]. Its role has been established in chronic antral gastritis, duodenal ulcer, chronic gastric ulcer, dyspepsia, gastric carcinoma and gastric lymphoma. The risk of gastric cancer has been estimated to be six-fold higher in *H. pylori* infected population than in uninfected population [5]. Biopsy urease test has a sensitivity of 80-99% and a specificity of 92-100% [6, 7]. *H. pylori* specific IgG antibody levels in serum by enzyme linked immunosorbent assays (ELISA) with a high sensitivity and specificity. The present study was undertaken to find out the rate of occurrence of *H. pylori* from endoscopic biopsy based tests (urease test and histology) and serology.

OBJECTIVES:

1. To find out the rate of occurrence of *H. pylori* in dyspeptic patients, using biopsy based tests and serology.
2. Evaluation of different diagnostic tests for *H. pylori*.
3. To find out the spectrum of histopathological changes that occurs in the gastro-duodenal mucosa in cases of dyspepsia

MATERIALS AND METHODS:

Study Type: Cross-sectional, hospital based observational study

Study Design: Prospective cohort study

Study setting/area, Population and Period: The study was conducted in the Department of Pathology, in a tertiary care hospital, from May 2014 to September 2015.

Sampling:

Selection of cohort of patient with dyspepsia

The Institutional Ethics Committee approved the protocol of the study.

60 adult dyspeptic patients attending the gastroenterology clinic and undergoing endoscopy, in our hospital and other hospitals in and around locality, were included in the study. All the patients were enrolled in the study after an informed consent. Patient's personal identification, socio-economic status, detailed clinical history and physical findings were noted.

Sampling technique: Consecutive non probability technique used, consensus sampling.

Patient inclusion Criteria:

1. Dyspeptic patients in the age group of 18 years to 60 years.
2. Patient who had given informed consent.

Patient Exclusion Criteria:

(i) Age < 18 years or > 60 years, (ii) patients who has taken anti-*H. pylori* antibiotics within 1 month, (iii) proton pump inhibitors (PPI) within 2 weeks or H_2 receptor blocker drugs within 1 week prior to endoscopy, (iv) patients with ischemic heart disease, chronic lung disease, diabetes mellitus, thyroid disease and on long standing NSAID therapy, (v) pregnant or lactating females and (vi) patients with known malignancy.

Study Techniques:

Case history and clinical examination, collection & testing of specimens followed by statistical analysis done.

Techniques in details: Case history and clinical examination of patients were done; cases selected maintaining the inclusion and exclusion criteria. 2 ml venous blood collected in plain vial for Anti *H. pylori* IgG ELISA. The serum samples labeled and brought to the laboratory keeping in an ice-lined chamber then allowed freezing in the deep freezer at -20°C for 2 days for future testing. Endoscopic biopsy was taken from the stomach: 4 quadrant antral biopsy, from corpus, fundus and from areas with endoscopic evidence of any other pathology. For biopsy urease test three endoscopic tissue bits-2 bits from the antrum and 1 from the corpus were taken and put in a numbered vial containing freshly prepared urea solution and change of colour was observed. Immediate change of colour from yellow to pink

and considered as positive for urease test. A test tube containing urea solution was taken as control for colour change. Three endoscopic tissue bits collected in 10% buffered neutral formalin for histopathological diagnosis by H & E stain. Modified Giemsa stain was used for demonstration of *H. pylori* in tissue sections; *H. pylori* is seen as spiral shaped bacteria, stained dark blue, seen in or near adherent mucus on the luminal side of the gastric surface and pit epithelial cells.

Data analysis: All data collected were tabulated on a grand chart and the statistical tests were performed by using the Statistical Package for Social Sciences (SPSS), Version 16 software for windows. The tests of significance were done by Chi-square test (X^2) and p value <0.05 was taken as significant.

Statistical methods: Percentage, prevalence, Pearson Chi-Square Test etc. were calculated using SPSS.

RESULT AND ANALYSIS:

Dyspepsia was more common in male (68.3%) than females (31.66%). The sex ratio was 2.15:1. In our study, three diagnostic tests for *H. pylori* were performed, namely- (i) anti-*H. pylori* IgG ELISA, (ii) biopsy urease test, and (iii) modified Giemsa stain for the demonstration of the organism in biopsy specimen. Majority of the patients (65%) were positive for the biopsy urease test. Modified Giemsa stain for *H. pylori* in tissue section was positive in 33 cases (55%).

Table 1: Rate of occurrence of *H. pylori* in dyspepsia.

		+ve for <i>H. pylori</i> -ve for <i>H. pylori</i>	
		(Total 41)	(Total 19)
Urease +ve (n = 39)	Giemsa +ve	31	-
	Giemsa -ve	8	-
Urease -ve (n = 21)	Giemsa +ve	2	-
	Giemsa -ve	19	19

Table 1 shows that the rate of occurrence of *H. pylori* in dyspeptic patients is = (41/60) X 100 % = 68.33%. Case definition: A case of *Helicobacter pylori* infection was defined as those patients who presented with dyspeptic symptoms and were positive for at least one of the two diagnostic tests- biopsy urease test and/or modified Giemsa stain.

Table 2: Analysis of the test of significance between socio-economic status, age, sex and dietary habits.

		<i>H. pylori</i> Positive: Count (%)	<i>H. pylori</i> Negative: Count (%)	Pearson Chi Square Test (X^2)
Socio- Economic Status	Lower	20(80.0%)	5(20.0%)	Value: 6.837, Df: 2, Asymp. Sig. (2- sided): 0.033
	Middle	17(70.8%)	7(29.2%)	
	Upper	4(36.4%)	7(63.6%)	
Age	< 30 Years	15(53.5%)	13(46.4%)	Value: 5.287, Df: 1, Asymp. Sig. (2- sided): 0.0214
	>30 Years	26(81.2%)	6(18.7%)	
Sex	Male	26(63.4%)	15(36.6%)	Value: 1.448, Df: 1 Asymp. Sig. (2- sided): 0.229
	Female	15(78.9%)	4(21.1%)	
Dietary Habits	Vegetarian	8(57.1%)	6(42.8%)	Value: 1.057 Df: 1 Asymp. Sig. (2- sided): 0.303
	Non- Vegetarian	33(71.7%)	13(28.2%)	

Table 2: Shows that highest numbers of dyspeptic patients were in the 'lower' socio-economic group (as per Kuppuswamy's Socioeconomic Status Scale) [8]. 80.0% were positive for *H. pylori* infection which is statistically significant. 41 dyspeptic patients were male and majority of them were positive for *H. pylori*. Positivity for *H. pylori* in female patients (78.9%), which is statistically not significant. Among non-vegetarian cases 71.7% and among vegetarian cases 57.1% were positive for *H. pylori* infection which is statistically not significant.

Table 3: Endoscopic findings in cases of dyspepsia

ENDOSCOPIC FINDINGS	NUMBER OF CASES (% OF TOTAL)	+ve for <i>H. pylori</i> (% OF EACH GROUP)	-ve for <i>H. pylori</i> (% OF EACH GROUP)
Normal UGI study	31 (51.6%)	19 (61.2%)	12 (38.7%)
Non-erosive antral gastritis	9 (15.0%)	9 (100.0%)	0 (0.0%)
Deformed pylorus	5 (8.3%)	3 (60.0%)	2 (40.0%)

Erosive gastritis	4 (6.6%)	4 (100%)	0 (0.0%)
Duodenal ulceration	4 (6.6%)	4 (100%)	0 (0.0%)
Active gastric ulcer	3 (5.0%)	2 (66.6%)	1 (33.3%)
Neoplastic ulcerated lesion	2 (3.3%)	0 (0.0%)	2 (100%)
Oesophagitis	2 (3.3%)	0 (0.0%)	2 (100%)
TOTAL	60	41	19

Table 3 shows that out of 31 dyspeptic patients with a normal upper gastro-intestinal endoscopy, 19 patients (61.2%) were positive for *H. pylori* infection. Hence, the rate of occurrence of *H. pylori* in functional dyspepsia in this study is = (19/31) X 100% = 61.29%.

Table 4: Histopathological findings in dyspepsia

Histopathological Findings	Number Of Cases	+Ve For <i>H. pylori</i> (% Of Each Group)	-Ve For <i>H. pylori</i> (% Of Each Group)
Chronic superficial gastritis	44 (73.3%)	31 (70.4%)	13 (29.5%)
Normal histopathological findings	4 (6.6%)	2 (50.0%)	2 (50.0%)
Duodenitis/duodenal ulceration	4 (6.6%)	4 (100.0%)	0 (0.0%)
Benign gastric ulcer	3 (5.0%)	2 (66.6%)	1 (33.3%)
Well differentiated adenocarcinoma	2 (3.3%)	0 (0.0%)	2 (100%)
Atrophic gastritis	1 (1.6%)	1 (100%)	0 (0.0%)
Intestinal metaplasia	1 (1.6%)	1 (100%)	0 (0.0%)
Barrett's oesophagus	1 (1.6%)	0 (0.0%)	1 (100%)
TOTAL	60	41	19

Table 4 shows that out of 60 cases 41 were positive for *H. pylori* and the commonest histological finding was chronic superficial gastritis followed by normal histopathological findings and duodenitis/duodenal ulceration.

Table 5: Sensitivity and specificity of IgG ELISA

Ig G ELISA	Modified Giemsa stain		Parameter
	Positive (n=33)	Negative (n=27)	
Positive (n=35)	30	5	Sensitivity: 90.91% Specificity: 81.48% Positive Predictive Value: 85.71%, Negative Predictive Value: 88%, Method: Wilson Score.
Negative (n=25)	3	22	

Table 5 shows that IgG ELISA has a high sensitivity (90.91%) and good specificity (81.48%). This present study assesses the sensitivity and specificity of IgG ELISA by taking the histological demonstration of the organism by modified Giemsa stain as the 'Gold standard'.




Fig A: Positive and negative biopsy urease test. Fig B: Gastric and duodenal erosion at endoscopy. Fig C: Chronic superficial gastritis with mild lymphomononuclear cell infiltrates in the lamina propria (H&E X 400). Fig D: *H. pylori* stained by modified Giemsa stain (X 1000).

DISCUSSION:

The present study showed male preponderance and is consistent with studies done by Longman et al. and Raul V Destura et al who reported male: female ratio to be 2:1 [9]. Tests for *H. pylori* by IgG ELISA showed positive in 58.33% cases. A similar study done by Malik A et al. in 1999 showed IgG ELISA positivity in 53.3% cases [10]. Biopsy

urease test was positive 39 (65%) cases. Malik A et al. found 61.2% urease test positivity; while Rao P et al. and Arora U et al. found 76.3% and 72% biopsy urease test positivity respectively [10,11,12]. With modified Giemsa stain for the organism in tissue sections, the bacterium could be demonstrated in 33 cases (55%). Rate of occurrence of *H. pylori* in dyspeptic patients in this study was 68.33%. Rao P et al observed 76.2% positivity for *H. pylori* among patients with gastroduodenal disorders as compared to 9.1% among asymptomatic individuals [11]. Shrivastava UK et al. in their study, showed 65% positivity for *H. pylori* in functional dyspepsia [13]. The socio-economic status analyzed by Kuppuswamy's Socio-economic Status Scale which was updated and modified in 2007 [8]. The 'lower' socio-economic group had the highest number of dyspeptic patients (25 cases); among them 20 cases (80%) were positive for *H. pylori* which is statistically significant. Studies done previously in India and outside, also observed similar association [14,15]. Highest percentage of *H. pylori* infection identified in >30 years age group. In a study by Rajesh Kumar et al. in 2006 in Jammu, India, maximum percentage of *H. pylori* positive cases were in the age group 36-45 years [14]. The present study results are consistent with earlier results. Relation between dietary habits and *H. pylori* showed p-value was 0.303 (p > 0.05) which was statistically insignificant. No previous study has observed such a relationship. The rate of occurrence of *H. pylori* in functional dyspepsia was 61.29% in our study. Similar studies in India, reported 65% prevalence of *H. pylori* in functional dyspepsia [13]. H & E stained tissue sections show 73.3% cases were diagnosed as chronic superficial gastritis. In a similar study, Rajesh Kumar et al. showed chronic superficial gastritis as the most common biopsy finding in dyspepsia [14]. Duodenitis/duodenal ulceration all cases (100%) were positive for *H. pylori*. Dayal VM et al. and Prasad S et al. have reported 75-100% association of *H. pylori* with duodenitis/duodenal ulceration [16,17]. Gastric carcinoma is a known complication of *H. pylori* infection, but any inference could not be drawn from the 2 cases of adenocarcinoma as the population was too small. The sensitivity and specificity of IgG ELISA was calculated. The results were- sensitivity 90.91% and specificity 81.48%. The results are consistent with previous studies [18]. IgG ELISA test cannot be used as a confirmatory test because of low specificity and positive predictive value. Rao P et al. had observed similar results in their study and recommended IgG ELISA as a very good screening test [11].

CONCLUSION:

Dyspepsia was more common in male and highest numbers of dyspeptic patients were in the 'lower' socio-economic group and in >30 years age group. Majority were positive for *H. pylori* infection; diagnosed by biopsy urease test, modified Giemsa stain for *H. pylori* in tissue sections and IgG ELISA test. The rate of occurrence of *H. pylori* in dyspeptic patients was 68.33% and for functional dyspepsia was 61.29%.

Early diagnosis may prevent dyspepsia and complications of *H. pylori* infection like chronic superficial gastritis, benign gastric ulcer, duodenitis/duodenal ulceration even adenocarcinoma. The study was conducted in a tertiary health care with limited resources and smaller number of cases, may not reflect the actual scenario in general population.

Contribution by authors:

1. Dr. Tarak Banik-Concept designing and conducting the study & writing the manuscript.
2. Dr. Prasenjit Kumar Bar and Dr. Saikat Mandal - Conducting the study, statistical analysis and preparing the manuscript suitable for publication.
3. Dr. Subhrajyoti Naskar - Guiding the study procedure, statistical analysis and preparing the manuscript suitable for publication.

REFERENCES

1. Hammer J, Talley MJ. Non Ulcer Dyspepsia. Curr Opin Gastroenterol 1999; 15: 492-6.
2. William L. Hasler. In: Functional Dyspepsia, Harrison's Principles of Internal Medicine. 16th edition. McGraw-Hill, New York. 2005; Vol-I, 222.
3. Dunn BE, Cohen H, Blaser MJ: Helicobacter pylori. Clin Microbiol Rev 1997; 10: 720-741.
4. Megraud F. Epidemiology of H.pylori infection. Gastro Clin North Am 1993; 22: 73-88.
5. The EUROGAST Study Group: 1993. An international association between Helicobacter pylori infection and gastric cancer. Lancet. 1993;341(8857):1359-62
6. Wong BC, Wong WM, Wang WH, Tang VS, Young J, Lai KC, Yuen ST, Leung SY, Hu WH, Chan CK, Hui WM, Lam SK. An evaluation of invasive and non-invasive tests for the diagnosis of Helicobacter pylori infection in Chinese. Aliment Pharmacol Ther 2001; 15: 505-511.
7. Viala CH, Windsor HM, Forbes GM, Chairman SO, Marshall BJ, Mollison LC. Evaluation of a new formulation CLO test. J Gastroenterol Hepatol 2002; 17: 127-130.
8. N. Kumar, C. Shekhar, P. Kumar, A.S. Kundu. Kuppuswamy's Socioeconomic Status

Scale-Updating for 2007. The Indian Journal of Pediatrics. December 2007; Volume 74, Number 12: 215.

9. Raul V Destura, Eternity D Labio, Leah J Barrett, Cirle S Alcantara, Venancio I Gloria, Ma Lourdes O Daez and Richard L Guerrant. Laboratory diagnosis and susceptibility profile of Helicobacter pylori infection in the Philippines. Annals of Clinical Microbiology and Antimicrobials 2004; 3:25.
10. Malik A, Singhal S, Mukharji U. Evaluation of elisa based antibody test against H. Pylori in cases of non Ulcer Dyspepsia. Indian Journal of Medical Microbiology 1999; 17: 137-139.
11. P Rao, A Sarkar, PG Shivananda, G Pai. Comparison of Elisa for antibody detection and biopsy urease test against H. pylori in cases of gastroduodenal disorders. Indian J Med Sci 2001; 55: 366-70.
12. Arora U, Aggarwal A, Singh K. Comparative evaluation of conventional methods and elisa based IgG antibodies detection for diagnosis of Helicobacter pylori infection in cases of dyspepsia. Indian J Med Microbiol 2003; 21: 46-48.
13. Shrivastava Upendra K, Gupta Aman, Gupta Arun, Bhatia Arati. Role of Helicobacter pylori in functional dyspepsia. Indian Journal of Surgery 2004; 66: 341-346.
14. Rajesh Kumar, G. Bano, B. Kapoor, Sunil Sharma, Yudhvir Gupta. Clinical Profile in H. Pylori Positive Patients in Jammu. JK science 2006; Vol 8: No 3
15. Emmanuel Ike Ugwuja, Nicholas Chukwuka Ugwu. Helicobacter pylori In Uninvestigated Dyspepsia In Primary Cares In Abakaliki, Nigeria. Online J Health Allied Scs. 2007; 1: 4.
16. Dayal VM, Kumar P, Kamal J, Shahi SK, Agarwal BR. Triple drug therapy of H.pylori infection in duodenal ulcer disease Indian J Gastroenterol 1997; 16: 46-48.
17. Prasad S, Mathew M, Chandhy G, Prasanna Rajan D, Venkateswaran S, Ramakrishna BS. Prevalence of Helicobacter pylori in Southern Indian controls and patients in Gastroduodenal disease. J Gastroenterol Hepatol 1994; 9: 501-06.
18. J E Crabtree, T M Shallcross, R V Heatley, J I Wyatt. Evaluation of a commercial ELISA for serodiagnosis of Helicobacter pylori infection. J Clin Pathol 1991; 44: 326-328.