



EVALUATION OF INCIDENCE OF DRY EYE AFTER PHACOEMULSIFICATION CATARACT SURGERY

Ophthalmology

Dr. Pari S. Desai	M.B.B.S, M.S.(Ophthalmology) Department of Ophthalmology, Dr. D. Y. Patil Hospital & Research Centre, Pimpri, Pune, Maharashtra, India.
Dr. O. K. Radhakrishnan*	M.B.B.S, M.S. (Ophthalmology) Professor, Department of Ophthalmology, Dr. D. Y. Patil Hospital & Research Centre, Pimpri, Pune, Maharashtra, India. *Corresponding Author
Dr. Shivani Pattnaik	M.B.B.S, M.S.(Ophthalmology) Department of Ophthalmology, Dr. D. Y. Patil Hospital & Research Centre, Pimpri, Pune, Maharashtra, India.

ABSTRACT

Methods: A descriptive longitudinal study conducted in tertiary care hospital during 2016-2018. Total 150 patients undergoing cataract surgery were included. Tear meniscus height (TMH), Tear film break up time (TBUT), Schirmer's test, ocular surface disease index (OSDI) parameters were studied pre-operatively and post-operatively.

Purpose: Evaluation of incidence of dry eye after phacoemulsification cataract surgery

Results: Out of 150, there were 81 (54%) males and 69 (46%) females with mean age of 63.58 ± 8.03 years, 88 (59%) subjects were operated for right eye cataract while 62 (41%) for left eye. Lower mean values of TMH, TBUT & Schirmer's test were observed in operated eye in comparison with non-operated eye after cataract surgery at 1 week, 1 month & 3 month follow up. Difference in mean was statistically highly significant ($p < 0.001$). 18 (12%) cases diagnosed as dry eye after cataract surgery.

Conclusions: Highly significant decreased in value of tear meniscus height, tear film break up time, Schirmer's test occurs after phacoemulsification surgery at 1 week post operatively and showed gradual improvement towards end of 3rd month.

KEYWORDS

Phacoemulsification cataract surgery, tear meniscus height, tear film break up time, Schirmer's test, ocular surface disease index

INTRODUCTION:

Cataract is a very important cause of blindness worldwide. Around 18 million people are blind due to bilateral cataract, accounting for 48% of all causes of blindness.¹ As per The World Health Organization (WHO) report, approximately 37 million people were blind in the world in 2002.¹ Cataract surgery is cost-effective and results in almost immediate visual rehabilitation.² Improvement in visual acuity after cataract surgery results in considerable gain in real life activities and emotional and social life components.³ Though postoperative improvement in vision occurs, many patients are dissatisfied due to tear film dysfunction, poor near vision and reduced contrast sensitivity. Studies suggested that cataract surgery could aggravate dry eye.^{4,5}

Reports from North and Eastern India shows dry eye prevalence which varies between 18.4% and 40.8%.⁶⁻⁹ One small study from high altitude showed a higher prevalence of 54%.¹⁰ In Manual small-incision cataract surgery (SICS) a straight or a curved incision is made on the superior sclera which penetrates the cornea at the level of the schwalbe's line, whereas phacoemulsification cataract surgery involves a clear corneal incision.¹¹ Corneal nerve fibres are located in the stromal layer and originate from the temporal and nasal areas, then bifurcate through the entire cornea, with the most dense fibres located in the central and paracentral areas. Incision at any site of the cornea may cause disruption of nerve fibres consequently reducing corneal sensitivity and reducing the tear production.¹² Dry eye symptoms range from mild transient irritation to persistent dryness, burning, itching, redness, pain, ocular fatigue and visual disturbance. Severe dry eye results in impairment in daily living, work productivity and affect mood and confidence.¹³ The current study will evaluate the dry eye status in patients with senile cataract undergoing phacoemulsification cataract surgery.

MATERIAL & METHODS:

A descriptive longitudinal study was conducted on patients undergoing phacoemulsification surgery for age related cataract in tertiary care hospital in western Maharashtra during period of September 2016 to August 2018. Ethical committee clearance was taken prior to study. Total 150 patients were included in this study, after obtaining informed written consent from patients, each patient was assigned into two groups: Cases group A (Eye scheduled for phacoemulsification cataract surgery) & Control group B (Unoperated eye of the same patient). visual acuity, tear meniscus height (TMH), tear film break up time (TBUT), Schirmer's test, Ocular surface disease index (OSDI) were performed pre-operatively and post operatively at 1 week, 1 month, and 3rd month.

Inclusion criteria:

1. Age of patient > 50 years.
2. Bilateral age-related cataract.

Exclusion criteria:

1. Other eye having pseudophakia.
2. Cataract caused by trauma, uveitis, drug induced etc.
3. Ocular diseases like glaucoma, disorders of lids, conjunctiva, cornea, sclera or using medication for same.
4. History of corneal refractive surgery
5. History of chemical burns, radiation or using contact lens.

Statistical analysis:

The statistical analysis was done by using SPSS software 16 version. Quantitative data was analysed as mean $\pm$ standard deviation (SD) & Qualitative data as frequencies. Unpaired t-test was used to compare normally distributed continuous variable between groups.

RESULTS:

Out of 150, there were 81 (54%) males and 69 (46%) females in the study. Mean age of study sample was 63.58 ± 8.03 years, with the oldest individual being 85 years old and youngest being 51 years of age. 67 (44.67%) patients were from 61-70 years age group followed by 57 (38%) patients in 51-60 years age group. 88 (59%) patients were operated for right eye cataract while 62 (41%) patients were operated for left eye cataract.




Figure 1: Line diagram showing mean values among study samples at different time interval

Lower mean values of tear meniscus height, tear film break up time and Schirmer's test were noted in operated eye for cataract surgery in comparison with non-operated eye after cataract surgery at 1 week, 1 month & 3 month follow up. The difference in values were gradually reduced over period of time but still lower in operated eyes at the end of third month postoperatively. (Figure 1).

Table 1: Tear meniscus height (mm) comparison by unpaired t test (operated vs non-operated eye)

Unpaired t-test for Equality of Means						
Time	t	df	Sig. (2-tailed)	Mean Difference	Std. Error Difference	95% CI of the Difference Lower Upper
Pre-op	.121	298	.904	.0033333	.0275050	-.0507953 .0574620
1 Week	-7.586	294.873	.000	-.2000000	.0263658	-.2518889 -.1481111
1 Month	-4.908	298	.000	-.1366667	.0278445	-.1914635 -.0818699
3 Month	-2.706	297.401	.007	-.0766667	.0283330	-.1324252 -.0209081

Table 2: Tear film break up time (sec) comparison by unpaired t test (operated vs non-operated eye)

Unpaired t-test for Equality of Means						
Time	t	df	Sig. (2-tailed)	Mean Difference	Std. Error Difference	95% CI of the Difference Lower Upper
Pre-op	1.095	298	.274	.253333	.2312628	-.2017818 .7084484
1 Week	-13.108	285.983	.000	-3.35333	.2558319	-3.8568857 -2.8497810
1 Month	-8.982	266.165	.000	-2.35333	.2619977	-2.8691850 -1.8374816
3 Month	-7.272	275.228	.000	-1.87333	.2576119	-2.3804735 -1.3661932

Table 3: Schirmer's test (mm) comparison by unpaired t test (operated vs non-operated eye)

Unpaired t-test for Equality of Means						
Time	t	Df	Sig. (2-tailed)	Mean Difference	Std. Error Difference	95% CI of the Difference Lower Upper
Pre-op	-.146	298	.884	-.086667	.5948777	-1.2573601 1.0840268
1 Week	-7.224	264.771	.000	-5.35333	.7410062	-6.8123480 -3.8943187
1 Month	-5.065	269.194	.000	-3.786667	.7476607	-5.2586727 -2.3146606
3 Month	-4.416	262.601	.000	-3.16000	.7155588	-4.5689631 -1.7510369

On application of unpaired t test meniscus height, tear film break up time and Schirmer's test mean values in operated & non-operated eyes shows significant difference. (Table 1, 2 & 3). OSDI score was lowest at 1 week follow up and gradually improved.

There were 18 (12%) cases diagnosed as dry eye after cataract surgery, while 132 (88%) were normal after surgery at the end of 3rd month post operatively. In this study OSDI (sensitivity 100%, specificity 90.37%) was found more reliable than TBUT (sensitivity 100%, specificity 53.33%) for diagnosis of dry eye taking into consideration that schirmer test is the gold standard.

DISCUSSION:

7 out of 18 patients developed dry eye at the end of 3 months post operatively in 61-70 age group & 6 were from 71-80 years age group. Kamla Dodia et al¹⁴ in their study found that dry eye is associated with higher age (>65 years). Ram J et al¹ showed that tear film stability was affected most commonly in older age group between 66-75 years.

Significant decrease in TMH observed in this study post-operatively, similar observation was found in study done by Cho et al¹⁵ and Gharraee et al.¹⁶ In contrast to this, Sitompul et al¹⁷ did not found any significant difference in TMH on follow up visit compared to preoperative values. Li and colleagues⁵ reported that misuse of topical eyedrops with preservatives is one of the important reasons for dry eye after cataract surgery. Surface irregularities at incision site may cause faster tear film break up. Kasetsuwan N et al¹⁸ evaluated dry eye following cataract surgery in 92 patients. TBUT deteriorated 7 days post phacoemulsification cataract surgery. After 3 months, there was a gradual improvement in TBUT values. Sinha M et al¹⁹ showed deterioration in TBUT at 1 week, 1 month and 3 months. However, Ram et al⁴ in 23 patients reported no difference in TBUT pre and post phacoemulsification cataract surgery which contrasts with our study. This difference may be because of small sample size of their study.

Gradual improvement in Schirmer's test value after 3 months postoperatively was observed by Cetinkaya et al²¹ and Kasetsuwan N et al¹⁸ like this study findings. Cho et al¹⁵, Gharraee et al¹⁶ and Li et al⁵ also observed that Schirmer's test values were reduced on post-operative day

30 and it was statistically significant.

This study shows deterioration of OSDI score following cataract surgery, similar observation was found in Kasetsuwan N et al¹⁸, Bhattacharjee S et al²¹ & Sinha M et al.¹⁹ Study by Li et al⁵ did not find statistically significant difference in OSDI score pre and post cataract surgery. Calvin W Roberts et al²² showed that use of cyclosporine 0.05% emulsion twice daily for one month prior and one month following cataract surgery reduces dry eye occurrence post operatively.

CONCLUSION:

Phacoemulsification cataract surgery affects tear film adversely. Tear meniscus height, tear film break up time, Schirmer's test and OSDI score values showed deterioration following phacoemulsification cataract surgery. Deterioration peaked at 1 week post-operatively and showed gradual improvement towards end of 3 months postoperatively. Incidence of dry eye after phacoemulsification cataract surgery in our study was 12% at 3 months postoperatively. Evaluation of patients pre and post phacoemulsification cataract surgery should be done for identification of unhealthy ocular surface which will ensure promising results postoperatively.

Recommendation:

After phacoemulsification cataract surgery, dry eye symptoms can develop immediately, and the severity can peak on seventh day post-operatively. Phacoemulsification cataract surgery worsen dry eye symptoms post operatively in patients with pre-existing dry eye. It is of paramount importance that ophthalmologists assess dry eye before and after phacoemulsification cataract surgery to ensure good quality of vision and patient satisfaction. When dry eye is diagnosed pre or post operatively, surgeon should add topical preservative free lubricating drops and in exceptional circumstances topical cyclosporine drops. One should use ophthalmic viscosurgical devices (OVDs) on corneal surface during phacoemulsification cataract surgery to reduce the trauma induced by surgery and BSS irrigating solution flushing, especially in the patients who are diagnosed with dry eyes preoperatively.

REFERENCES:

- Resnikoff S, Pascolini D, Etya'ale D, Kocur I, Pararajasegaram R, Pokharel GP, Mariotti SP. Global data on visual impairment in the year 2002. Bulletin of the world health organization. 2004; 82:844-51.
- Lansingh VC, Carter MJ, Martens M. Global cost-effectiveness of cataract surgery. Ophthalmology. 2007 Sep 1;114(9):1670-8.
- Lamoureux EL, Fenwick E, Pesudovs K, Tan D. The impact of cataract surgery on quality of life. Current opinion in ophthalmology. 2011 Jan 1;22(1):19-27.
- Ram J, Gupta A, Brar GS, Kaushik S, Gupta A. Outcomes of phacoemulsification in patients with dry eye. Journal of Cataract & Refractive Surgery. 2002 Aug 1;28(8):1386-9.
- Li XM, Hu L, Hu J, Wang W. Investigation of dry eye disease and analysis of the pathogenic factors in patients after cataract surgery. Cornea. 2007 Oct 1;26: S16-20.
- Gupta SK, Gupta V, Joshi S, Tandon R. Subclinically dry eyes in urban Delhi: an impact of air pollution? Ophthalmologica. 2002;216(5):368-71.
- Sahai A, Malik P. Dry eye: prevalence and attributable risk factors in a hospital-based population. Indian journal of ophthalmology. 2005 Apr 1;53(2):87.
- Gupta N, Prasad I, Jain R, D'Souza P. Estimating the prevalence of dry eye among Indian patients attending a tertiary ophthalmology clinic. Annals of Tropical Medicine & Parasitology. 2010 Apr 1;104(3):247-55.
- Basak SK, Pal PP, Basak S, Bandyopadhyay A, Choudhury S, Sar S. Prevalence of dry eye diseases in hospital-based population in West Bengal, Eastern India. Journal of the Indian Medical Association. 2012 Nov;110(11):789-94.
- Gupta N, Prasad I, Himashree G, D'Souza P. Prevalence of dry eye at high altitude: a case controlled comparative study. High altitude medicine & biology. 2008 Dec 1;9(4):327-34.
- Linebarger EJ, Hardten DR, Shah GK, Lindstrom RL. Phacoemulsification and modern cataract surgery. Survey of ophthalmology. 1999 Sep 1;44(2):123-47.
- Kohlhaas M. Corneal sensation after cataract and refractive surgery. Journal of Cataract & Refractive Surgery. 1998 Oct 1;24(10):1399-409.
- Friedman NJ. Impact of dry eye disease and treatment on quality of life. Current opinion in ophthalmology. 2010 Jul 1;21(4):310-6.
- Dodia K, Bapat S, Chudasama RK. Dry eye risk factors after phacoemulsification cataract surgery at a secondary care hospital. International Journal of Health & Allied Sciences. 2013 Oct 1;2(4):242.
- Cho YK, Kim MS. Dry eye after cataract surgery and associated intraoperative risk factors. Korean Journal of Ophthalmology. 2009 Jun 1;23(2):65-73.
- Gharraee H, Mousavi M, Daneshvar R, Hosseini M, Sazande S. Effect of clear corneal incision location on tear film following phacoemulsification surgery. Iranian Journal of ophthalmology. 2009 Jan 1;21(3):2934.
- Sitompul R, Sancoyo GS, Hutauruk JA, Gondhowiardjo TD. Sensitivity change in cornea and tear layer due to incision difference on cataract surgery with either manual small-incision cataract surgery or phacoemulsification. Cornea. 2008 Sep 1;27:S13-8.
- Kasetsuwan N, Satitpitakul V, Changul T, Jariyakosol S. Incidence and pattern of dry eye after cataract surgery. PLoS one. 2013 Nov 12;8(11):e78657.
- Sinha M, Sinha A, Chowdhury B. Comparative evaluation of dry eye following cataract surgery: a study from north india. IOSR Journal of Dental and Medical Sciences. 2014;13(6):13-8.
- Cetinkaya S, Mestan E, Acir NO, Cetinkaya YF, Dadaci Z, Yener HI. The course of dry eye after phacoemulsification surgery. BMC ophthalmology. 2015 Dec;15(1):68.
- Bhattacharjee S, Ram A, Ferdows SS, Pan DC, Nanda IA. KEYWORDS Phacoemulsification, Dry Eyes, Corneal Surgery, Cataract Surgery. EVALUATION OF DRY EYE SYMPTOMS AFTER PHACOEMULSIFICATION. 2018 Feb 15(97284).
- Roberts CW, Carniglia PE, Brazzo BG. Comparison of topical cyclosporine, punctal occlusion, and a combination for the treatment of dry eye. Cornea. 2007 Aug 1;26(7):805-9.