



## THYROID SWELLINGS- A CLINICOPATHOLOGICAL STUDY

### ENT

**Dr. Anchal Gupta\*** Senior Resident, Department of ENT, Head and Neck Surgery, SMGS Hospital, Government Medical College, Jammu, Jammu and Kashmir, India. \*Corresponding Author

**Dr. Padam Singh Jamwal** Professor, Department of ENT, Head and Neck Surgery, SMGS Hospital, Government Medical College, Jammu, Jammu and Kashmir, India

### ABSTRACT

**Background:** Diseases of the thyroid continue to be a common clinical problem having a prevalence rate of 4 - 7% in the general population.

**Materials and methods:** This retrospective study was conducted in the Department of ENT and Head & Neck Surgery, SMGS Hospital, Jammu from January 2019 to January 2020. A thorough history taking, meticulous clinical examination, routine hematological and biochemical investigations were done in all the patients. General examination of the patient was done and looked for thyroid functional abnormality followed by local examination was carried out to locate and identify the site of the swelling, shape, size and consistency of the thyroid swelling and clinical diagnosis was made. All thyroid swelling cases were subjected to FNAC. The type of thyroidectomy is based on the size and type of thyroid whether neoplastic or non-neoplastic. All the specimens were sent for histopathological examination.

**Results:** Out of 60 patients in the study, 56 were females and only 4 were males. The most common age group was 31-40 years seen in 24(40%) patients. The most common presenting symptom was swelling in the neck seen in 58(96.67%) patients. The most common diagnosis on FNAC was colloid goiter seen in 40(66.67%) patients. The most common was benign nodular colloid goiter seen in 35(58.33%) patients. **Conclusion:** Fine needle aspiration is a very useful and indispensable test in the diagnosis of thyroid lesions.

### KEYWORDS

FNAC, thyroid, swelling.

### INTRODUCTION

Thomas Wharton first coined the term "thyroid" because of the organ's close proximity to the thyroid cartilage (120-200 A.D). The word thyroid is derived from the Greek "thyros" meaning "shield" because it was originally considered to protect the larynx.<sup>[1]</sup>

Normal thyroid gland is impalpable. Enlargement of thyroid gland is the most common manifestation of thyroid disease. The enlargement may be either generalized or localized which again may be toxic or nontoxic. The nontoxic goiter is further divided on etiological basis as endemic goiter and sporadic goiter. The endemic goiter is defined as one where more than 10% of population shows thyroid enlargement.<sup>[2]</sup> Lesion of thyroid are predominantly confined to females in the ratio of 5:1. The fine needle aspiration method for studying the thyroid was first developed in Sweden in the Rudiunhelmet hospital of Stockholm in the 1950s.<sup>[3]</sup> Frable in 1983 used the FNAC as a means of diagnosing most thyroid masses as either neoplasm or goitre nodules.<sup>[4]</sup> The main aim of the study is to analyze the clinical presentation of thyroid swellings and to study the role of FNAC in the diagnosis of thyroid swelling.

### MATERIALS AND METHODS

This retrospective study was conducted in the Department of ENT and Head & Neck Surgery, SMGS Hospital, Jammu from January 2019 to January 2020. A thorough history taking, meticulous clinical examination, routine hematological and biochemical investigations were done in all the patients. General examination of the patient was done and looked for thyroid functional abnormality followed by local examination was carried out to locate and identify the site of the swelling, shape, size and consistency of the thyroid swelling and clinical diagnosis was made.

All thyroid swelling cases were subjected to FNAC as out-patient procedure after explaining the details of the procedure to the patient and taking an informed written consent.

All patients were explained about the lesions based on the FNAC report and were advised surgery. Patients who were willing for surgery were included in the study. The type of thyroidectomy is based on the size and type of thyroid whether neoplastic or non-neoplastic.

All the specimens were fixed in 10% buffered formalin and sent for histopathological examination.

### OBSERVATIONS AND RESULTS:

60 patients were included in the study

### Age and Sex distribution of patients:

Out of 60 patients in the study, 56 were females and only 4 were males. The youngest patient in our study was 14 years old and eldest patient was 68 years old. The most common age group was 31-40 years seen in 24(40%) patients. The overall average age was 32.5 years. In females it was 33.4 years whereas in males it was 28 years.




Figure 1: Gender wise distribution of patients.




Figure 2: Age wise and gender wise distribution of patients.

### Clinical presentation of patients.

The most common presenting symptom was swelling in the neck seen in 58(96.67%). Next most common symptom was pressure symptoms seen in 4(6.67%) patients. 3(5%) patients presented each with pain and neck nodes whereas 1(1.67%) each presented with dysphagia and voice change.[Figure 3]




Figure 3: Distribution of patients according to symptoms.

**FNAC findings**

After thorough general and local examination FNAC was done in all the 60 patients. The most common diagnosis was colloid goiter seen in 40(66.67%) patients. Next most common diagnosis was follicular neoplasm followed by atypical adenoma, thyroid cyst, carcinoma and thyroiditis respectively in decreasing order.[Table 1]

Table 1: Distribution of patients according to FNAC findings.

FNAC Findings	No. of patients	%
Colloid goiter	40	66.67%
Thyroid cyst	3	5%
Thyroiditis	2	3.33%
Follicular neoplasm	8	13.33%
Atypical adenoma	4	6.67%
Papillary Carcinoma	3	5%

**Type of surgery**

All the 60 patients included in the study underwent thyroidectomy. 3 patients who were diagnosed as papillary carcinoma on FNAC underwent total thyroidectomy along with neck dissection depending on levels of lymph nodes involved. Figure 4 shows distribution of patients according to type of thyroidectomy they underwent.




Figure 4: Type of thyroidectomy.

**Histopathological diagnosis**

All the thyroidectomy specimens were sent for histopathological examination. Table 2 shows distribution of patients according to histopathological diagnosis.

The most common was benign nodular colloid goiter seen in 35(58.33%) patients followed by thyroiditis seen in 12(20%) patients, follicular adenoma seen in 8(13.33%) patients, papillary carcinoma seen in 3(5%) patients and follicular carcinoma seen in 2(3.33%).

Table 2: distribution of patients according to histopathological diagnosis

Histopathological diagnosis	No. of patients	%
Benign nodular colloid goiter	35	58.33%
Thyroiditis	12	20%
Follicular adenoma	8	13.33%
Papillary carcinoma	3	5%
Follicular carcinoma	2	3.33%

**DISCUSSION**

Thyroid lesion usually present with swelling of thyroid gland with or

without hyper or hypofunction of thyroid gland. The incidence of thyroid lesions is increasing in recent years. Thyroid surgery is nowadays associated with low morbidity and extremely low mortality. Out of 60 patients in the study, 56 were females and only 4 were males with female: male ratio of 14:1. The most common age group was 31-40 years seen in 24(40%) patients. There is a wide variation of this ratio among different studies, Likhari KS et al had a ratio of 11.5:1, whereas Sengupta A et al had a ratio of 3.8:1.<sup>[5,6]</sup> This is in agreement with the study conducted by Sengupta A et al with mean age of 35.39 years and maximum incidence among age group 31-40 years.<sup>[6]</sup>

The most common diagnosis was colloid goiter seen in 40(66.67%) patients. Next most common diagnosis was follicular neoplasm followed by atypical adenoma, thyroid cyst, carcinoma and thyroiditis respectively in decreasing order.

The most common histopathological diagnosis was benign nodular colloid goiter seen in 35(58.33%) patients followed by thyroiditis seen in 12(20%) patients, follicular adenoma seen in 8(13.33%) patients, papillary carcinoma seen in 3(5%) patients and follicular carcinoma seen in 2(3.33%). These findings are similar to findings by Rout K et al and Wahid FI et al.<sup>[7,8]</sup>

**CONCLUSION**

Thyroid swellings are common in females they occur in 3rd and 4th decade most commonly. FNAC is very useful in the diagnosis.

**Funding:** No funding sources

**Conflict of interest:** None declared

**REFERENCES**

1. Medvei VC. A history of endocrinology. In: Kovacs K, Asa Science. 1998. p.1.
2. Maitra A. Thyroid gland. In: Kumar V, Abbas AK, Fausto N, Aster JC, editors Robbins and Cotran Pathological Basis of Disease. 8th Ed. Philadelphia: Saunders Co; 2010:1107-1126.
3. Galera-Davidson, Gonzalez-Campora R. Thyroid. In: Marluce bibbo, David Wilbur, editors. Comprehensive Cytopathology. 3rd ed. Philadelphia: Saunders Elsevier; 2008:p.633
4. Frable WJ, Frable MA. Thin needle aspiration biopsy: The diagnosis of head and neck tumours revisited. Cancer 1979; 43:1541-48.
5. Likhari KS, Hazari RA, Gupta Sg, Shukla U. Diagnostic accuracy of fine needle aspiration cytology in thyroid lesions: A hospital based study. Thyroid research and Practise. 2013;10(2):68-71.
6. Sengupta A, Pal R, Kar S, Zaman FA, Sengupta S, Pal S. Fine needle aspiration cytology as diagnostic tool in thyroid enlargement. J Nat Sci Biol Med. 2011;2(1):113-8.
7. Rout K, Ray CS, Behera SK, Biswal R. A comparative study of FNAC and histopathology of thyroid swellings. Indian J Otolaryngol Head and Neck Surg. 2011;63(4):370-2.
8. Wahid FI, Hussain M, Khan A, Ahmadkhan I. Diagnostic yield of fine needle aspiration cytology in the diagnosis of Thyroid Nodule and its comparison with national and international studies. ISRA Med J. 2012;4(4):230-4.