



THYROID DISORDERS AND POLYCYSTIC OVARY SYNDROME

Gynaecology

Dr Suman Sharma Assistant Professor, Department Of Gyan. & Obst, RNT Medical College, Udaipur, Rajasthan.

ABSTRACT

BACKGROUND: Polycystic ovarian syndrome (PCOS) is the most common form of chronic anovulation associated with androgen excess; perhaps occurring in 5-10% of reproductive women. **METHODS:** Observational study done in the Department of Obstetrics & Gynaecology. 50 cases of women with PCOS based on Rotterdam's criteria and an equal number of age-matched controls (women without PCOS) were included in the study. **RESULTS:** T4 level was significantly lower in PCOS group with mean free T4 level 0.85 ± 0.67 ng/ml in PCOS group v/s 1.93 ± 0.83 ng/ml in control group (p-value = 0.001). TSH level was significantly higher in PCOS group with mean TSH level 8.18 ± 6.38 mU/L in PCOS group v/s 3.42 ± 1.19 mU/L in control group (p-value = 0.001). **CONCLUSION:** High prevalence of thyroid disorders in PCOS patients thus points towards the importance of early correction of hypothyroidism in the management of infertility associated with PCOS.

KEYWORDS

Women, T4, T3, TSH, PCOS

INTRODUCTION

Polycystic ovarian syndrome (PCOS) is the most common form of chronic anovulation associated with androgen excess; perhaps occurring in 5-10% of reproductive women. PCOS is viewed as a heterogeneous disorder of multifactorial etiology. It is also associated with increased metabolic and cardiovascular risk factors. These risks are linked to insulin resistance and compounded by the common occurrence of obesity, although insulin resistance is also present in non-obese woman with PCOS. During the reproductive years, PCOS is associated with important reproductive morbidity including infertility, irregular uterine bleeding and increased pregnancy loss.²

Dysfunction and anatomic abnormalities of the thyroid are among the most common diseases of the endocrine gland. Abnormalities in the supply of thyroid hormone to the peripheral tissue are associated with alteration in a number of metabolic processes. Early stages of thyroid dysfunction (before symptoms are obvious) can lead to subtle change in ovulation and endometrial receptivity, which may have profound effect on fertility. Infantile hypothyroidism if untreated, leads to sexual immaturity. Untreated juvenile hypothyroidism causes a delay in the onset of puberty followed by anovulatory cycles. In adult woman, severe hypothyroidism may be associated with diminished libido and failure of ovulation. Primary ovarian failure can also be seen in patients with Hashimoto's thyroiditis as a part of autoimmune polyglandular syndrome. Rarely, in primary hypothyroidism, secondary depression of pituitary function may lead to ovarian atrophy and amenorrhoea.³

MATERIAL & METHODS

Type of Study: Observational study

CASE: Women with PCOS which were diagnosed by Rotterdam's Criteria were cases.

Inclusion Criteria

- Age group - 13-45 years.
- Giving written informed consent

Exclusion Criteria

- Women on OCPs
- Women on steroids
- Hyperprolactinemia
- Congenital Adrenal Hyperplasia
- Cushing's Syndrome
- Virilizing tumor of ovary
- Vitiligo
- Endometriosis

CONTROL: Women of the same age group visiting OPD with problems unrelated to Rotterdam's Criteria of PCOS were controls.

Inclusion Criteria

- Age group - 13-45 years.
- Giving written informed consent

Exclusion Criteria

- Menstrual irregularity
- Hyperandrogenism
- With polycystic ovaries
- Insulin resistance
- Inflammatory and autoimmune disease
- Metabolic abnormalities

RESULTS

Table-1. Risk of Thyroid Disorder in PCOS and Control Group

Group	With Thyroid Disease		Without Thyroid Disease	
	No.	%	No.	%
PCOS	19	38.00	31	62.00
Control	3	6.00	47	94.00

38.00% PCOS cases were present with thyroid disorder.

Table-2. Statistical Value of Thyroid Specific Variable of Case and Control Group

Variables	Cases		Controls		p-value
	Mean	SD	Mean	SD	
T3	2.42	1.23	2.08	1.08	>0.05
T4	0.85	0.67	1.93	0.83	<0.001
TSH	8.18	6.38	3.42	1.19	<0.001

T4 level was significantly lower in PCOS group with mean free T4 level 0.85 ± 0.67 ng/ml in PCOS group v/s 1.93 ± 0.83 ng/ml in control group (p-value = 0.001). TSH level was significantly higher in PCOS group with mean TSH level 8.18 ± 6.38 mU/L in PCOS group v/s 3.42 ± 1.19 mU/L in control group (p-value = 0.001)

DISCUSSION

Patients with PCOS often have defective progesterone secretion which leads to an increased estrogen to progesterone ratio. Oestrogen can increase the expression of IL-6 in T cell and inhibitory action of progesterone may lead to over stimulated immune system and makes these patients more prone to autoimmune disorder.⁴

T4 level was significantly lower in PCOS group with mean free T4 level 0.85 ± 0.67 ng/ml in PCOS group v/s 1.93 ± 0.83 ng/ml in control group (p-value = 0.001). TSH level was significantly higher in PCOS group with mean TSH level 8.18 ± 6.38 mU/L in PCOS group v/s 3.42 ± 1.19 mU/L in control group (p-value = 0.001). Similar results were reported by Sinha U et al (2013).⁵ In our study mean serum TSH level was found to be significantly higher in PCOS group and in control group. Significant difference was found between two groups. Similar correlation between TSH and Anti-TPO antibody level was reported by Janssen OE et al (2004)¹

Pinto et al analysed the relationship between selected clinical and metabolic parameters in women with PCOS and subclinical hypothyroidism. Out of the 168 women studied, 19 had subclinical

hypothyroidism. Considering the metabolic parameters, serum low-density lipoprotein cholesterol and PRL levels were significantly higher in the women with SCH (122.6 ± 25.6 mg/dL and 17.7 ± 7.7 ng/mL, respectively) compared with those with normal thyroid function (105.6 ± 33 mg/dL and 14 ± 10.3 ng/mL, respectively).⁶

Janssen, et al observed a prevalence of autoimmune thyroiditis (biochemically) in 26.9% of their 175 patients whereas Ozdemir et al among 107 patients found a prevalence of 30.5%. In his study, Thyroid pathologies were observed in half of the patients with PCOS.^{1,7}

CONCLUSION

High prevalence of thyroid disorders in PCOS patients thus points towards the importance of early correction of hypothyroidism in the management of infertility associated with PCOS.

REFERENCES

1. Janssen OE, Mehlmauer N, Hahn S, et al. High prevalence of autoimmune thyroiditis in patients with polycystic ovary syndrome. *Eur J Endocrinol.* 2004; 150(3): 363–369.
2. Azziz R, Sanchez L, Knochenhauer E, Moran C, Lazenby J, Stephens K et al. Androgen excess in women : Experience with over 1,000 consecutive patients. *The Journal of Clinical Endocrinology and Metabolism.* 2004; 89(2): 453–462.
3. Stein I, Leventhal M. Amenorrhoea associated with bilateral polycystic ovaries. *American Journal of Obstetrics and Gynecology.* 1935; 29(2): 181–191.
4. Du D, X Li. The relationship between thyroiditis and polycystic ovary syndrome: a meta-analysis. *Int J Clin Exp Med.* 2013; 6(10): 880–9.
5. Sinha U, Sinharay K, Saha S, Longkumer SN, Baul SN, Pal SK. Thyroid disorders in polycystic ovarian syndrome subjects: A tertiary hospital based cross-sectional study. *Eur J Endocrinol.* 2004; 150(3): 363–369.
6. Benetti-Pinto CL, Berini Piccolo VR, Garmes HM, Teatin Juliato CR. Subclinical hypothyroidism in young women with polycystic ovary syndrome: An analysis of clinical, hormonal, and metabolic parameters. *Fertil Steril.* 2013; 99:588–92.
7. Ghosh S, Kabir SN, Pakrashi A, Chatterjee S, Chakravarty B. Subclinical hypothyroidism: A determinant of polycystic ovary syndrome. *Horm Res.* 1993; 39:61–6