



A STUDY TO ASSESS THE ROLE OF A RENIN ALDOSTERONE SYSTEM GENE (RAS) POLYMORPHISM IN THE EXISTENCE OF TOXAEMIA OF PREGNANCY.

Anatomy

Dr. T. Praveen

Assistant. Professor, Department of Anatomy, Shadan institute of medical sciences, Teaching hospital & Research centre, Hyderabad, Telangana, (India).

Dr. Nelapati Swetha Vinela*

Assistant. Professor, Department of Bio chemistry, Ayaan institute of medical sciences, Teaching hospital & Research centre, Kanakamamidi, R.R. dist., Telangana, (India).
*Corresponding Author

ABSTRACT

Toxaemia of pregnancy is diagnosed in India around 6-10% of all deliveries and is associated with 22% of perinatal foetal deaths and cause 30% of maternal death. Angiotensin Converting Enzyme (ACE) plays a vital role in the Renin Aldosterone System (RAS) which regulates blood pressure by converting Angiotensin-I into a powerful vasoconstrictor Angiotensin-II. The aim of the present study was to explore out the ACE gene I/D polymorphism in cases of toxaemia of pregnancy and normotensive mothers in south India population. Study was done in 240 cases of Toxaemia of pregnancy and 240 controls (Normotensive) pregnant women who were admitted in Department of obstetrics and gynaecology, Shadan institute of medical sciences, teaching hospital & Research centre, Telangana. 5ml venous blood was collected in EDTA vials and processed for ACE gene I/D polymorphism with using conventional polymerase chain reaction (PCR). A Deletion polymorphism (D allele) has found to be associated with elevated ACE activity leading to hypertension in pregnant women. On genotype analysis, the gene frequency for I/I, I/D and D/D polymorphism in controls group were found 45%, 34% and 21% respectively. Among case group, the frequency of I/I, I/D and D/D polymorphism were noted 16%, 32% and 52% respectively. On comparison of the individual frequencies of the three genotypes, it was found that D/D genotype were more prone to develop disease with an odd's ratio of 6.8866% (D/D vs I/I) and 2.5506% (D/D vs I/D). On statistical comparison of D/D genotypes between the controls and cases, the cases showed a significant increase of frequency (p-value=0.0001), an odd's ratio of 3.9960 was obtained against (I/D+I/I) polymorphism.

KEYWORDS

Toxaemia of Pregnancy, Angiotensin Converting Enzyme (ACE), Insertion/Deletion polymorphism(I/D), polymerase chain reaction (PCR), significant p-value.

INTRODUCTION

Hypertensive disorders are creating complications during pregnancy (Toxaemia of pregnancy) which are common and forming deadly traits along with haemorrhage and infection.^[1] Pre-eclampsia (PE) is a disease occurs during the pregnancy which is specified by the inception of hypertension and the presence of protein in the urine in large amount.^[2] Pre-eclampsia is considered if one or more of the following criteria are present: Blood pressure 140 mm Hg or higher systolic or 90 mm Hg or higher diastolic after 20 weeks of gestation in a woman with previously normal blood pressure. Proteinuria: 0.3g or more of protein in a 24-hours urine collection (usually correspond with 1+ or greater on a urine dipstick test) known as mild preeclampsia.^[1] When systolic blood pressure of 160 mm of Hg or higher or 110 mm of Hg or higher diastolic on two occasions at least six hours apart in a woman on bed rest, the condition is known as severe preeclampsia. It is associated with proteinuria and oliguria, cerebral or visual disturbances, pulmonary oedema of cyanosis, epigastric pain or right upper quadrant pain, impaired liver function, thrombocytopenia and foetal growth restriction known as Eclampsia.^[1]

Toxaemia of pregnancy is a key cause of large number of maternal deaths and thereof foetal deaths. Maternal hypertension (toxaemia of pregnancy) is diagnosed in 6-10% of all deliveries which is associated with 22% of perinatal foetal deaths and 30% of maternal death.^[3]

Angiotensin converting enzyme (ACE) is a monomeric, membrane-bound, zinc and chloride dependent peptidyl dipeptidase that catalysis the conversion of decapeptide angiotensin I to the octapeptide angiotensin II by removing a carboxy terminal dipeptide. ACE has been well known for a key part of renin-angiotensin system (RAS) that regulates blood pressure in pregnant women.^[4] ACE plays a vital role in the RAS which regulates blood pressure. ACE gene High activity can contribute to hypertension because of its vasoconstriction effect.^[5,6]

Angiotensin-converting enzyme (ACE) gene is located on chromosome 17q23 position. It is 21 kb in size and consists of 26 exons and 25 introns.^[7] ACE gene insertion/deletion (I/D) polymorphism defined by the presence of an extra 287 base pair sequence inside intron 16 of the ACE gene. A deletion polymorphism (D allele) has been reported to be associated with elevated ACE activity in plasma and serum.^[7] Some investigators have been reported a strong association between the ACE gene D and I allele polymorphism in pregnant women from various geographical regions.^[7] Allele DD genotype is

associated with higher levels of ACE gene in serum and II genotype is concerned with its lower levels, ID genotype is associated with its intermediate levels of ACE gene in serum. Moreover, it has been assumed that I allele has a sequence similar to a silencer role, which might explain the reason; why the D allele is associated with increased risk of preeclampsia or PIH.^[7]

MATERIALS AND METHODOLOGY

The present study was done in 480 pregnant mothers divided into two groups. 240 samples were taken from case (patients) suffered from Toxaemia of Pregnancy. Another 240 samples were collected from normal pregnant women as control (Normotensive). All the cases and controls samples taken from pregnant women who were admitted in the department of obstetrics and gynaecology, Shadan institute of medical sciences, Telangana (India). All the cases and controls pregnant women have filled written consent form for willing to give their samples for this study. Inclusion criteria: Antenatal mothers diagnosed with pregnancy induced hypertension with their blood pressure of 140/90 mm of Hg or more in to case group. Beside this standard questionnaires were prepared to get the past and present medical/surgical history of cases and controls. In questionnaires several parameters were taken such as history of renal, liver failure, seizures, mother who has the hypertensive disorder before the pregnancy and other medical problems. The permission has taken from the institution ethical committee prior to conduction of this study.

5ml venous blood was collected in EDTA vials and stored in -20°C temperature till the processing of ACE gene I/D polymorphism with simple polymerase chain reaction (PCR).

Following Primers were used for the detection of ACE gene I/D polymorphism in this study:

Forward primer: 5'-CTGGAGACCCCATCCTTCT-3' Tm- 59°C
Reverse primer: 5'-GATGTCGCCATCACATTCGTCAGAT-3' Tm- 58°C

Primers were synthesized from Bangalore Genei and reconstituted with sterile TE buffer (0.1 mM) and further dilution was done with double distilled water as manufacturer's instruction. Primers final (working) concentration was 10 picomoles. The PCR conditions were 95°C for 3 min, 35 cycles of 95°C for 30s, 52°C for 30s, 72°C for 1.20 min and final extension was at 72°C for 5 min. Then PCR product was run with 1% agarose gel containing ethidium bromide. Genotype was

determined by fragment size under UV light in gel documentation system (Bio-Rad) and PCR product was directly submitted for gene sequencing to Chromous Biotech Pvt. Ltd (Bengaluru).

RESULTS

All the DNA samples (240 cases and 240 controls) were amplified and analyzed for I/D polymorphism of the ACE gene. In this study, D and I alleles were identified at 190 bp and 490 bp respectively, leading to the confirmation of the three types of polymorphism D/D, I/D and I/I in ACE gene.




Fig 1, Agarose gel picture, Single band present at 190bp was represented as D homozygous while single band present at 490bp was denoted as I homozygous and presence of two bands denoted as heterozygous conditions.

With the use of genotype analysis, the gene frequency for I/I, I/D and D/D polymorphism in controls group were found 45%, 34% and 21% respectively. Among case group, the frequency of I/I, I/D and D/D polymorphism were noted 16%, 32% and 52% respectively.

Table 1: Distribution Of The Polymorphisms Between The Controls And Cases.

Type Of Ace Gene Polymorphism	Controls (240)	Toxaemia Of Pregnancy (240)	X ² Value	P-Value
I/I	108(45.0%)	38(15.83%)	66.501	0.0001
I/D	80(33.33%)	76(31.66%)		
D/D	52(21.70%)	126(52.55%)		

Table 2: Distribution Of Genotype Between Controls And Cases

Type of ACE polymorphisms	Controls (240)	Toxaemia Of Pregnancy (240)	Odds ratio (95% CI)	P-Value
D/D	52 (21.70%)	126 (52.55%)	3.9960	0.0001
IX (ID + II)	188 (78.33%)	114 (47.99%)	(2.175-7.016) Z - 4.822	

With the use of biostatistics Odd's ratio was calculated for cases/controls: 3.9960.

Table 3: Distribution Of Genotype Between Dd And Ii Genotype Among Controls And Cases.

Genotype	Controls (240)	Toxaemia Of Pregnancy (240)	Odds ratio (95% CI)	P value
D/D	52 (21.70%)	126 (52.55%)	6.8866	0.0001
I/I	108(45%)	38 (15.83%)	(3.439-13.789) Z - 5.447	

On comparison of the individual frequencies of the three genotypes, it was found that D/D genotype were 6 times more prone to develop Toxaemia of pregnancy than I/I genotype with odd's ratio 6.8866.

Table 4: Distribution Of Genotype Between Dd And Id Among Controls And Cases.

Genotype	Controls (240)	Toxaemia Of Pregnancy (240)	Odds ratio (95% CI)	P value
D/D	52 (21.70%)	126 (52.55%)	2.5506	0.0040
I/D	80(33.33%)	76 (31.70%)	(1.348-4.823) Z - 2.881	

Similarly, with the comparison of the individual frequencies it was found that D/D genotype was 2 times more prone to produce Toxaemia of pregnancy than I/D genotype with Odd's ratio 2.353.

First Submitted sequence

TACGCCAATTTTATTCCAGCTCTGAAATTCTCTGAGCTCCCC
TTACAAGCAGAGGTGAGCTAAGGGCTGGAGCTCAAGGCAT
TCAAACCCCTACCAGATCTGACGAATGTGATGGCCACATCA

GGTTTACGCCAATTTTATTCCAGCTCTGAAATTCTCTGAGCT
CCCCTTACAAGCAGAGGTGAGCTAAGGGCTGGAGCTCAAG
GCATTCAAACCCCTACCAGATCTGACGAATGTGATGGCCAC
ATCAGATTCGTGAGATCTGGTAGGGGTTTGAATGCCTTGAG
CTCCAGCCCTTAGCTCACCCTCTGCTTGTAGGGGGAGCTCAA
AAAAATTTCAAAGCTGGAACAAAATTGGCGAAACACATAA
AAGTGACTGTTTGGGACGAGCTAGAGAAATGGGAAAA
AGGATGGGAGTGGTCTCCAGGA

Fig 2, the obtained ACE gene sequence of 490bp (upper) band.

```

Query 246 CATCACATTGTCAGATCTGGTAGGGGTTTGAATGCCTTGAGCTCCAGCCCTTAGCTCAC 303
          |||
Sbjc 61630153 CATCACATTGTCAGATCTGGTAGGGGTTTGAATGCCTTGAGCTCCAGCCCTTAGCTCAC 1630094

Query 304 CTCTGCTTGTAGGGGAGCTCAAAAATTTCAAGCTGGAACAAAATTGGCGAAACCA 361
          |||
Sbjc 61630093 CTCTGCTTGTAGGGGAGCTCAGAAATTTCAAGCTGGAATTTAAATTTGGCGAAACCA 61630034

Query 362 TAAAAGTGACTGTTTGGCGAGGCTAGAGAAATGGGAAAGGATGGGAGTGCTCT 415
          |||
Sbjc 61630033 TAAAAGTGACTGTATAGGCGAGGCTAGAGAAATGGGAAAGGATGGGAGTGCTCT 1629974

Query 416 CCAG 419
          |||
Sbjc61629973 CCAG 61629970
  
```

Fig 3, Homology sequence alignment of 490bp (upper) band was indicated the Insertion (I) polymorphism of ACE gene in population. The red mark nucleotide has replaced by black colour nucleotide.

AGCTAAGGGCTGGAGCTCAAGGCATTCAAACCCCTACCAG
ATCTGACGAATGTGATGGCCACATCAGGTTTCGCCATTTAT
TCCAGCTCTGAAATTTCTCTGAGCTCCCTTACAAGCAGAT
GTGAGCTAAGGGCTGGAGCTCAAGGCATTCAAACCCCTAC
CAGATCTGACAATGTGATGGCCACATCA

Fig 4, the obtained ACE gene sequence of 192bp (lower) band.

```

Query 1 GGTITGOCCTTTTATTCAGCTCTGAAATTTCTCTGAGCTCCCTTACAAGCAGAGGTG 60
          |||
Sbjc 61630037 GGTITGOCCTTTTATTCAGCTCTGAAATTTCTCTGAGCTCCCTTACAAGCAGAGGTG 61630096

Query 61 AGCTAAGGGCTGGAGCTCAAGGCATTCAAACCCCTACCAGATCTGAC-AATGTGATGGCC 120
          |||
Sbjc 61630097 AGCTAAGGGCTGGAGCTCAAGGCATTCAAACCCCTACCAGATCTGACGAATGTGATGGCC 1630156

Query 121 AC 122
          ||
Sbjc 61630157 AC 61630158
  
```

Fig 5, Homology sequence alignment of 192bp (lower) band was indicated the Deletion (D) polymorphism of ACE gene in studied samples.

Statistical analysis with comparison of D/D genotypes between the controls and cases, the cases have showed a significant increase of frequency (p value=0.0001) and odd's ratio of 3.9960 was obtained against (I/D+I/I) polymorphism. This has showed that cases with D/D genotypes are three times more prone for the development of Toxaemia disorder. Similarly, statistical comparison of the individual frequencies among three genotypes, it was found that D/D genotype were found more prone to develop disease with an odd's ratio of 6.8866% (D/D vs I/I) and 2.5506% (D/D vs I/D).

DISCUSSION

According to (Urban2015) it has been mentioned in his study by the analysis of 168 samples that preeclampsia toxaemia women genotype DD of the I/D polymorphism of the ACE gene has been associated with the risk of preeclampsia.^[8] Rahimiet al., 2012 also recorded in their study of pre-eclamptic women they have mentioned the association between D/D genotype of ACE I/D polymorphism and risk of preeclampsia.^[9] However, (Dimri, et al., 2011) in his study of north Indian women had noticed DD genotype of ACE gene I/D polymorphism appears to predispose to severe pre-eclampsia but not non-severe pre-eclampsia.^[10] Correspondingly, (Chen, et al., 2011) in their meta analysis study had mentioned about the subgroup analysis by ethnicity which increased PIH risk was found among both Asians and Caucasians. In this report it has been mentioned a significantly elevated risk among Indian and Asian countries population due to D allele of ACE gene polymorphism.^[11] Although, Uma et al., 2010 in have not been precisely supported that association between the DD genotype of the ACE gene during the later onset of the disease.^[12]

In the report of Biller et al., 2000 it has been recorded that the ACE

genemay regulate the level of human serum ACE concentration. Human serum ACE is at the highest level of activity in DDgenotype carriers in during the disorder of Toxemia^[13]. Correspondingly, Balet *al.*, 2002 had been found the frequency of the D allele was higher in cases than that in the control group. It had also recorded in the study about the serum concentration of ACE was higher in the PIH group than that in the control group.^[14]It had also mentioned by Kauret *al.*, 2005 about the frequency of the DD genotype in the PIH group which had around 60%, while it was only 30% in the control group. The study had showed the statistical difference up to significant level. And the frequency of the D allele in patients with PIH (74%) had been found higher than that the control group (56%). And also has mentioned about the association of ACE I/D polymorphism with PE with OR of 3.35 for the DD genotype and 2.24 for D allele.^[15] In another report presented by Mrozikiewicz *et al.*, 2000 about the frequency of distribution of the D and I alleles of the ACE gene were inconsistent in cases and control groups, and had confirmed that the D allele may be a risk factor for PIH.^[16]

The data generated by present study coincides with results of various co-workers in this field and showing a strong association of D/D genotype of ACE gene I/D polymorphism and toxemia of pregnancy in the north Indian population. The main finding of this study was to explore out the D allele of the ACE I/D polymorphism which was may be major risk factor for toxemia of pregnancy in the studied populations.

CONCLUSION

Based on the data generated by this study, it may conclude that D/D genotype were more prone to develop disease with an odd's ratio of 6.8866% (D/D vs I/I) and 2.5506% (D/D vs I/D) respectively. This genotype polymorphism has asignificant association between D allele of the ACE gene and increased risk for toxemia of pregnancy in the north Indian population. Further researches with a larger dataset is needed for better results to prevent study bias. This study is novel and not available in literatures till date.

REFERENCES

1. ACOG practice Bulletin. Diagnosis and management of preeclampsia and Eclampsia. obsetgynaecol; 2002;99: 159-167.
2. Eiland, Elosha, Nzerue, et al., "Preeclampsia 2012". J. of Pregnancy. (2012); 1-7.
3. Fernandodarias. In Practical guide to high risk pregnancy and delivery, Mosby, Harcourtasia private Ltd. 2000; 2nd edition: 184-185.
4. Skeggs, LT., Kahn, J. Rand Shumway, N.p. Preparation and function of hypertension converting enzyme. THE Journal of Experimental Medicine. 1956; 103, 295-299.
5. K.K. Griendling, T.J. Murphy and R.W alexander. Molecular biology of the preeclampsia syndrome. Journal of perinatal medicine. 2008; Vol.36: No.1, pp. 38-58.
6. C.Hubert, A. M. Houot, P. Corvol, and F.Soubier. Structure of the angiotensin I converting enzyme: Two alternate promoters correspond to evolutionary steps of a duplicated gene,"the Journal of biological chemistry. 1993; vol.87: no.6, pp1816-1828.
7. C MAndo, p.Antonazzo, S.Tabano, Angiotensin converting enzyme and adducing-1 polymorphism in women with preeclampsia and gestational hypertension. Reproductive sciences. 2004; Vol. 16.; No.9, PP.127-131.
8. Katarzyna Kusmierska-Urban. Association of ACE I/D and AGT M235T Gene polymorphism with the Gestational hypertension and the fetal Growth. Obsteric and gynaecology International Journal.2015; March-April.
9. Zohreh Rahimi, Ziba Rahimi, hadiMo Zafari. Preeclampsia and angiotensin converting enzyme (ACE) I/D and angiotensin II type-1 Receptor (AT1R) A1166C polymorphism: association with ACE I/D polymorphism. Journal of the Renin-ngiotensin-aldosterone system. 2012; 14 (2):174-180.
10. Aggarwal S, DimriN, Tandon I, Preeclampsia in North Indian Women: The contribution of genetic polymorphism. J Obsteric Gynaecology Res; 2011;37:1335-1341.
11. Zhen Chen, FangXu, Yonggang Wei. Angiotensin converting enzyme insertion/deletion polymorphism and risk of pregnancy hypertensive disorders: a Meta-Analysis. Journal of the Renin-ngiotensin-aldosterone system. 2011;13(1); 184-195.
12. Uma R, Forsyth SJ, Struthers. Polymorphism of angiotensin converting enzyme Gene in early-onset and late-onset preeclampsia. J Materna fetal Neonatal Med.2010; 23: 874-879.
13. Biller H, Ruprecht B, Gaede KI, et al. Gene polymorphisms of ACE and the angiotensin receptor AT2R1 influence serum ACE levels in sarcoidosis. Sarcoidosis Vasc Diffuse Lung Di. 2009;26(2): 139-146.
14. Bai H, Liu X, Liu R., et al. Angiotensinogen and angiotensin-I converting enzyme gene variations in Chinese pregnancy induced hypertension. Hua Xi Yi Ke Da Xue Xue Bao; 2002; 33: 233-237.
15. Kaur R, Jain V, Khuller M, et al. Association of angiotensin converting enzyme gene polymorphism with pregnancy induced hypertension. Acta Obstet Gynecol Sci.; 2005; 84: 929-933.
16. Seremak-Mrozikiewicz A, Drews K, Malewski Z, et al. Polymorphic genotypes of the angiotensin-converting enzyme in severe pregnancy-induced hypertension. Arch Perinat Med.2000; 6: 22-24.