



CADAVERIC STUDY OF ANATOMICAL VARIATION OF LUMBRICALS IN HAND

Anatomy

Dr. Praveen Kumar

Tutor Deptt of Anatomy ANMMC Gaya.

Dr. Chandra Kiran Tutor Deptt of Anatomy ANMMC Gaya.

Dr. Nimisha Madhu

Tutor Deptt of Anatomy ANMMC Gaya.

Dr. Chandan Kumar*

Junior Resident Deptt of Anatomy ANMMC Gaya. *Corresponding Author

ABSTRACT

Background- To study the innervation pattern of lumbricals, proper knowledge about anatomy and variation prevalent in intrinsic musculature of hand is mandatory for various surgical procedure like cleft hand surgery.

Materials and methods- The study was carried on 54 hands sexes of different age groups used for routine dissection in anatomy department at ANMMC GAYA BIHAR. The distribution patterns of the muscular branches of median and ulnar nerve were studied.

Results- in the present study, variations in innervations of lumbricals from the normal pattern were found. In 3 hands (5.5%), the second lumbricals was solely supplied by a branch from of ulnar nerve. The third lumbricals was supplied by median nerve in 1 hand (1.85%) and by both median and ulnar in 4 hands.

Conclusion- clinician and hand surgeons should be aware of such a variation in innervations of lumbricals during various surgical procedures.

KEYWORDS

Lumbricals, Misplaced, insertion, and split insertion.

INTRODUCTION

The human hand has several intrinsic muscles which are responsible for the highly specialized and intricate movements that are characteristic of the human hand. Among all those muscles, the lumbricals have a significantly greater role to play in the intricate movements of the fingers though small in size.

The lumbricals of the upper limb are four small muscles resembling earthworms (*Ascaris lumbricoides*) and hence named so. In all anatomical descriptions they are conventionally numbered from the lateral to the medial side. The lumbricals take their origin in the palm from the four tendons of the flexor digitorum profundus and pass distally along the radial side of the corresponding metacarpophalangeal joints, in front of the deep transverse metacarpal ligament. Each muscle forms a narrow tendon and runs in a facial canal and on reaching the dorsal surface of the proximal phalanx, joins the radial margin of the dorsal digital expansion as the distal wing tendon.

The first and second lumbricals are unipennate, and are supplied by the median nerve while the third and fourth lumbricals are bipennate and are supplied by a deep branch of the ulnar nerve.

A large number of variations have been described in the literature ranging from complete absence of lumbricals to reduction in their numbers or presence of accessory slips. Neurovascular structures in the hand are variable within a population and that anomalous cases occasionally be encountered. This information may be useful to a surgeon operating in the palm, we undertook a cadaver study to decline more precisely the pathways of innervations of the lumbrical muscles.

MATERIALS AND METHODS

STUDYSITE-

The study was carried on 54 hands of both sexes of different age groups used for routine dissection in Anatomy department at Anugrah narayan magadh medical college Gaya Bihar.

METHODS

Beginning at the palmar wrist crease, the median & ulnar nerves were followed distally, taking care to dissect all the branches & show their anatomical relationships. The lumbrical were also carefully dissected & all areas of innervations noted. Each nerve innervating a lumbrical was traced to its origin from the parent nerve.




Figure 1: Shows the incision for exposure of lumbrical muscles

Observations

In the 54 hands which were dissected, the following observations were made. The first lumbrical was innervated by the median nerve via its proper palmar digital nerve to the lateral side of the index finger in all the cases. The second lumbrical received its nerve supply from the median nerve via its lateral common digital branch for the adjacent sides of index & middle finger in 51 hands. In 3 hands (5.5%), the muscles were solely supplied by the branch from the deep branch of ulnar nerve.

The third lumbrical was innervated by the deep branch of ulnar nerve in 49 hands. In one hand (1.85%) the muscle was solely supplied by the median nerve via its medial common digital branch for the adjacent sides of middle & ring finger. Along with deep branch of ulnar nerve, an additional source of innervation came from medial common digital branch of median nerve in 4 hands (7.4%). The fourth lumbrical received its nerve supply from the deep branch of the ulnar nerve.

In the present study, three types of variation were observed.

Table 1: Types of variations and their innervation and incidence

Variation 1	First, Second and Third lumbricals supplied by Median Nerve. Fourth lumbrical supplied by Deep Ulnar	1 Case (1.85%)
Variation 2	First & Second lumbricals supplied by Median Nerve. Third lumbrical supplied by Median & Deep Ulnar nerve. Fourth lumbrical supplied by Deep Ulnar	4 Cases (7.4%)

Variation 3	First lumbrical supplied by Median nerve. Second, third and Fourth lumbricals supplied by Deep Ulnar nerve.	3 Cases (5.5%)
-------------	--	----------------

VARIATION 1




FIG. 2: Third lumbrical supplied by Median nerve

VARIATION 2




FIG 3: Third Lumbrical supplied by median and deep branch of ulnar nerve

VARIATION 3




FIG. 4: Second lumbrical supplied by deep branch of the ulnar nerve

DISCUSSION

Neurovascular structures in the hand are variable within a population & that anomalous cases occasionally be encountered.

The nerve, supplying the muscles as textbooks describe, not absolutely true. Buchanan (1950): Mentioned that innervation to second lumbrical was inconstant, sometimes derived from deep ulnar & sometimes having a double supply. In the present study, normal pattern was found in 46 hands (85.2%). The median nerve supplied the 1st lumbrical in all cases. In 4 hands (5.5%), Second lumbrical was solely supplied by a branch from deep branch of ulnar nerve. In Mehta & Gartner's series (1961) the median nerve supplied 1st & 2nd lumbricals in all cases. In 3 hands (7.4%), additional branch to 3rd lumbrical came from median nerve (via medial common palmar digital nerve for the middle & ring finger).

Sparing of the innervation to the lumbrical muscles in carpal tunnel syndrome has been described (Desjacques et al, 1980). Unlike the motor branches to the thenar muscles which are superficial (palmar) & hence subject to compression against the flexor retinaculum, the branches to lumbricals are more dorsal & therefore better protected from direct compression. Morphological changes in this variation may be 1) Variation in myotome formation or 2) different path chosen by nerve fibres itself.

Rabischong (1962) noted that the lumbrical muscles have the greatest density of mechanoreceptors than any other muscle, & proposed that they play a critical role in proprioception in addition to their function as IP joint extensors. It would appear to be prudent to protect the innervation of the lumbricals wherever possible & in particular to avoid unnecessary mobilization of the digital nerves during surgical dissection in this area.

CONCLUSION

Clinician and hand surgeons should be aware of such a variation in innervation of lumbricals during various surgical procedures.

REFERENCES

1. R. S. Lauritzen, R. M. Szabo and D. B. Lauritzen; Innervation Of The Lumbrical Muscles; Journal of Hand Surgery, 1996 ,21B: 1:57-58.
2. Mehta, H. J. And Gardner, W. U. (1961). A study of lumbrical muscles in the human hand. Journal of Anatomy, 109: 227-238.
3. Rabischong, P. (1962). L'innervation proprioceptive des muscles lombricaux de la main chez l'homme. Revue de Chirurgie Orthopédique, 48:234-45.
4. Sunderland, S. and Ray, L. J. (1946). Metrical and non-metrical features of the muscular branches of the median nerve. Journal of Comparative Neurology, 85: 191-201.
5. M.L.Ajmani, M.D; Morphological variation of Lumbrical muscles in the Human Hand with Some Observation on its nerve supply; vol.3 No.2, January 2001.
6. Datta AK. Essentials of Human Anatomy. 2nd Ed. Calcutta: Current Books International 1995.
7. Joshi SD, Joshi SS, Athavale SA. Lumbrical Muscles and Carpal Tunnel. Anat Soc India 2005; 54(1): 12-5.