



EVALUATION OF CAUSATIVE FACTORS IN SETTING OF ACUTE BOWEL OBSTRUCTION

General Surgery

Dr. N. Mallikarjunarao

M.B.B.S, M.S., Assistant professor, Dept Of General Surgery, MIMS.

Dr. Sreenidhi Dhambolam*

M.B.B.S., Postgraduate In Dept Of General Surgery, MIMS. *Corresponding Author

KEYWORDS

INTRODUCTION

Acute intestinal obstruction is one of the commonest surgical emergencies in all age groups. It is defined as obstruction in forward propulsion of the contents due to mechanical or neurological causes. Mode of presentation is same in all but underlying cause varies in each age group. Now with better understanding of pathophysiology, improvement in radiological techniques of diagnosis and high degree of refinement in correction of fluid and electrolyte imbalance, introduction of antibiotics to effective bacteriological control, introduction of techniques in gastrointestinal decompression, new surgical principles, as in large bowel obstruction introduction of on table lavage and resection and primary anastomosis has replaced staged procedures and number of days in hospital stay. Improvement in field of anesthesia has all contributed to lowering the morbidity and mortality. The dictum of never let the sun set or rise in small bowel obstruction has made early surgical intervention for intestinal obstruction¹. This in turn has reduced the incidence of strangulation of bowel, which was major cause of mortality in already ill patient. Success in treatment of patient with acute intestinal obstruction depends largely upon early diagnosis, skillful management and appreciation of importance of treating the pathological effects of obstruction just as much as the cause itself.

AIMS AND OBJECTIVES

To study the incidence of tuberculosis in acute intestinal obstruction.
To analyze the modalities of operative management and postoperative complications.

MATERIALS & METHODS

Fifty cases of acute intestinal obstruction that presented to emergency department of Maharaja institute of Medical Sciences & General Hospital, nellimarla between August 2018 & march 2020 were studied.

Inclusion Criteria:

1. Patients of all age groups who attended OPD and emergency department at, Maharaja institute of medical sciences, nellimarla with history and clinical picture suggestive of acute intestinal obstruction.
2. Patients who had hernia with recent onset of irreducibility, pain, vomiting and constipation
3. Intestinal obstruction due to tuberculous etiology is included only on confirmation by histopathology of the lesion.

Exclusion Criteria:

Patients with history of sub-acute intestinal obstruction were excluded from this study.

- All patients with provisional diagnosis of acute intestinal obstruction were assessed clinically in detail as proforma after admission.
- Investigations done included Hb, Blood counts including TC, DC, ESR, X-ray Chest PA view, Plain X-ray abdomen erect film & Ultrasonography abdomen.
- Serum electrolytes were carried out in required cases only.
- All patients were subjected to surgery with objective to relieve the obstruction.

OBSERVATION AND RESULTS

- A clinical study of 50 cases of acute intestinal obstruction were studied during period of August 2018– March 2020 at Maharaja institute of medical sciences, nellimarla.

Analysis is as follows:

Age Distribution

- The study was done in all age groups ranging from new born to 85yrs
- with a mean age of 35 years.

Sex Incidence

Acute intestinal obstruction was commoner in males (72%) than females (28%). Male to female ratio was 2.6:1

Mode Of Presentation & Levels Of Obstruction

Small bowel obstructions (41) outnumbered large bowel obstruction (9).

Levels Of Obstruction

High small bowel obstruction: 10 cases
Low small bowel obstruction: 31 cases
Large bowel obstruction: 09 cases

Analysis Of Symptoms & Signs

Most commonly in 86% of cases there is tenderness, and in 70% of them with pain abdomen and vomiting. Guarding & rigidity seen in 42% & 4% cases respectively. groin swelling seen in 9 patients I.e 18%.

Duration of Pain

Most patients presented with pain lasting more than 2day duration.

Previous History Of Pain

Duration of previous history of pain lasted from 4-7 months.

Etiology & Intraoperative Findings In Obstruction

Small bowel adhesions were the commonest cause of acute intestinal obstruction. Tubercular Strictures accounted for 4% of all cases.

Small Bowel Obstruction - 41 cases (82%)

CAUSES	CASES	PERCENTAGE
Adhesions	13	26
Obstructed Hernias	09	18
Small bowel volvulus	07	14
Bands	06	12
TB Stricture	02	04
Intussusception	02	04
Meckel's Diverticulum	01	02
Meconium Ileus	01	02

Large Bowel Obstruction – 9 cases (18%)

CAUSES	CASES	PERCENTAGE
Neoplasms	3	6
Hirschprung's	3	6
Volvulus	2	4
Intussusception	1	2

Radiological Features

Plain X-ray erect abdomen was done in 39 cases out of 50 cases.

Positive interpretation was when it correlated with exact site of pathology and negative when it did not. in most of the cases multiple air fluid levels are seen.

Strangulation & Causes

The incidence of strangulation was seen in up to 30% patients (15).causes includes volvulus in 6, hernias in 4, adhesions in 2.

Management

Most common operative procedure done was resection & anastomosis (28%), followed by adhesiolysis (22%).

Small Bowel Obstruction -41 Cases

Adhesiolysis in 11 cases, Resection & Anastomosis-11, Band Release-05, Volvulus Derotation-03, Hernia Repair-05, Resection & Hernia Repair-04, Meckel's Diverticulectomy-01, Resection & Stoma-01

Large Bowel Obstruction- 9 cases

Resection & Anastomosis 3 cases

Colostomy 5 cases

Milking of Intussusception in 1 case

COMPLICATIONS

Complications in 50 cases of acute intestinal obstruction, morbidity is 28% & mortality is 12%.

Causes Of Morbidity

Wound sepsis, burst abdomen in 7 cases, faecal fistula in 1, pulmonary infections in 5 cases, septicemia in 3 cases were included as causes of morbidity.

Causes Of Mortality

- Septicemia: 3
- Respiratory Infection: 2
- Multi Organ Failure: 1
- Hypothermia: 1

Presence of strangulation and co-morbid conditions added mortality.

DISCUSSION

Intestinal obstruction is one of the common clinical entity. The mortality is reduced significantly by instituting the treatment at the earliest period. 1–4% of mortality in emergency surgeries is contributed by acute intestinal obstruction¹⁵.

The following were the observations made from the study of 50 patients of acute intestinal obstruction in both children and adults at Maharaja institute of medical sciences, nellimarla.

Incidence

In the present series small bowel obstruction contributed to 82% and large bowel obstruction 18%. This is comparable with reports of Michel³⁷ and Becker³⁸ where small bowel obstruction constituted to 80% and large bowel obstruction constituted 20%.

Age Incidence

The acute intestinal obstruction occurs in all age groups. Maximum incidence was seen between age group of 0–10 yrs (24%). Over all standard deviation is 24.57 years. Earlier studies conducted by Gill and Eggleston¹⁸ has reported 19.04% of cases in age group of 0–10 yrs. In another study by Budharaj³⁹ reported 13% of cases of acute intestinal obstruction below 12 year age. Fuzan¹⁰ reported mean age of 56 yrs.

Sex Incidence


In our study the incidence of intestinal obstruction in males was 36 (72%) and that of females was 14 (28%). Male to female ratio is 2.6:1.0 (3:1) Fuzan¹⁰ and Lee²⁸ reported 2:1 male to female ratio. Budharaj³⁹ reported in his study a ratio of 4:1 between male and female.

Etiology & Mode Of Presentation

In our study the following etiological factors were found,

- Adhesions: 26%
- Hernias: 18%
- Volvulus: 18%
- Bands: 12%
- CA colon: 06%
- Intussusceptions: 06%
- Hirschprung's: 06%
- Tubercular stricture: 04%

- Meckel's diverticulum: 02%
- Meconium ileus: 02%



Adhesions

A total of 26% of cases attributed to adhesions. Among adhesions 24% of the intestinal obstruction was due to postoperative adhesions and remaining 2% was found to be non specific. Majority of incidence was found within 1 year of surgery.

Jain and Prasad⁴⁰ found that adhesions contributed for intestinal obstruction upto 25.5%.

Ti and Young⁴¹ reported that postoperative adhesions and bands contributed upto 23.8% as cause of intestinal obstruction in 62 cases with only postoperative adhesions in 52 patients (19.2%)

Fuzan¹⁰ study in 582 patients found that, in 246 (42.2%) patients the cause for intestinal obstruction was adhesions due to previous operations.

Hernia

A total number of 9 (18%) cases of intestinal obstruction are related to hernia in this study of 50 cases. Out of these 9, 4 (44%) cases are strangulated hernia.

All 9 cases of hernia are in male patients.

All 4 of the strangulated hernia patients underwent resection and anastomosis.

In other five cases only hernia repair was done. All cases were of inguinal hernia.

In the series of Michel G. Sarr (1983) shows hernia related strangulation was present in 42% patients. Budharaj³⁹ studies revealed the etiology for acute intestinal obstruction secondary to obstructed hernia (small bowel and large bowel) accounted for 33%. In his study, the incidence of gangrene was up to 22%.

Volvulus

Volvulus constituted for about 18% in our studies that is 9 cases.

Out of these 9 cases, 7 were small bowel volvulus and 2 cases were of sigmoid volvulus.

A study conducted by Sankaran⁴² reported 24 cases of volvulus in South India among which sigmoid volvulus predominated forming 50% of cases.

Budharaj³⁹ series revealed that 18.2% of intestinal obstruction was due to volvulus & in that 11.9% was due to small bowel volvulus and 6.19% due to large bowel volvulus. Ramachandran¹⁹ in his study quotes that volvulus is the second commonest cause of small bowel obstruction which accounted for nearly 24%.

Gill¹⁸ reported that incidence of volvulus was 25% (36 cases), out of these small bowel volvulus accounted for 23 patients and large bowel volvulus for 13 patients.

Bands

In our study intestinal obstruction due to bands accounted for 12%.

A study series by Gill and Eggleston¹⁸ of 147 cases showed that 6.8% of small intestinal obstruction is due to bands.

Malignancy

In the present study acute intestinal obstruction related to malignancy constituted for 6%(3 cases). Two of the 3 are due to stricturous growth in left colon (one in sigmoid and one in descending colon) and one from right colonic growth.

Ti⁴¹ noted that carcinoma of descending colon and rectum constituted 37.2%. Ascending colon and caecum constituted 9.8%.

Thompson⁴⁴ in his series recorded the incidence of obstructing carcinoma of right colon equals 26% and left colon 69%.

Ramachandran¹⁹ found in his study that sigmoid colon cancer accounted for 6.6% of intestinal obstruction in large bowel, which is nearer to our study.

Fuzan¹⁰ revealed the cause of malignant large bowel obstruction of which ascending colon constituted 3.38% and sigmoid colon constituted up to 27%.

In our study sigmoid colon cancer, descending & ascending colon accounted 33.3% each.

INTUSSUSCEPTION

In our study of 50 cases of acute intestinal obstruction 3 cases were of intussusception (6%), out of these 3 cases, 2 were causing small bowel obstruction and 1 large bowel obstruction. All three were found in < 10yr of age.

Ti⁴¹ revealed his study of 261 patients the incidence of intussusception accounted for 6.3% (17 cases) of intestinal obstruction. In this 17 cases, 14 were infants and 3 adults.

Another series by Kuruvilla⁴⁶ intussusception accounted for 6.3% of the cases of total intestinal obstruction.

Hirschprung's Disease

Hirschprung's disease accounted for 6% (3 cases) in our studies.

In study of Ti⁴¹ Hirschprung's disease accounted for 4.8% of causes of intestinal obstruction.

In study series by Ramachandran¹⁹ Hirschprung's disease constituted for 10.4% cases of the pediatric intestinal obstruction

Tubercular Stricture

The present study accounted for 4% (2cases) of tubercular stricture as a cause for intestinal obstruction. Budharaj³⁹ in review of 242 cases reported that intestinal tuberculosis giving rise to acute intestinal obstruction was seen only in 2.1% of cases.

The study series of Sircar⁴⁷ reported to have 5% of cases of abdominal tuberculosis present with acute intestinal obstruction.

Meckel's Diverticulum

Our study concluded with the incidence of Meckel's diverticulum constituting for 2% (1case) of acute intestinal obstruction.

Budharaja³⁹ reported to have incidence of 1.23% of Meckel's diverticulum causing intestinal obstruction. Ramachandran¹⁹ in series stated about 4.2% of acute intestinal obstruction was due to Meckel's diverticulum.

Meconium Ileus

The incidence of meconium ileus is 1 in 2000 live births 49. In our study it was only one case that caused intestinal obstruction.

Clinical Features – Symptoms & Signs

Maximum presenting symptoms in this study was pain abdomen – 78% (39 cases), vomiting –70% (35 cases), distention abdomen –58% (29 cases), constipation –52% (26 cases).

Asbun⁵⁰ in their retrospective analysis of 105 cases of small bowel obstruction found that incidence of pain abdomen 82%, vomiting 88%, were commoner than constipation (28%) and distention of abdomen (56%).

Budharaj³⁹ in his study reported that, symptoms of in order of

frequency were pain abdomen 95%, distention of abdomen 82%, vomiting 75%, absolute constipation 50% constituting acute intestinal obstruction.

Symptoms & Signs	Present Series	Bhansali S.K. et al
Pain	78 %	96.05%
Vomiting	70 %	89%
Constipation	52 %	55%
Distension	58 %	43%
Tenderness	86 %	33%
Guarding	42 %	—

X-ray Findings

Multiple air fluid levels were the most common finding on x-ray erect abdomen accounting for 56% of positive cases. In Bhansali S.K. et al it was 76.1%.

Management

- All cases were operated in this study.
- Adhesiolysis done in 11 cases.
- Resection and anastomosis was done in 14 cases
 - 2 cases of tubercular stricture of ileum (Ileo-transverse anastomosis)
 - 4 cases of volvulus (small bowel volvulus), which had gangrenous changes and 2 cases of small bowel intussusception.
 - 1 case of band (causing twist of bowel)
 - 1 case of carcinoma colon and 2 cases of sigmoid volvulus.
- Release of bands was done in 5 cases.
- Derotation / undoing of volvulus was done in 3 cases.
- Only hernia repair done in 5 cases.
- Resection and hernia repair done in 4 cases.
- Colostomy done in 5 cases (2 cases of carcinoma, 3 Hirschprung's disease)
- Milking of intussusception in 1 case.

Intraoperative Findings


Adhesions were the most common finding found followed by volvulus and tubercular strictures.

Findings	Present Series	Bhansali S.K. et al
Adhesions	26%	27.4%
Volvulus	14%	1.48%
Strictures	4%	40.7%

Operative Procedure

Surgery	Present Series	Bhansali S.K. et al
Resection	38%	24.4%
Adhesiolysis	22%	17%
Anastomosis	22%	35.78%

Post Operative Complications



Complication	Present Series	Bhansali S.K. et al
Wound Infection	7	5.4
Chest Infection	5	--
Faecal Fistula	1	2.9
Septicemia	3	--
Hypothermia	1	--
Total (%)	34	22.94

Mortality

In our study it is 14% where as in Bhansali S.K. et al it is 11.84%.

CONCLUSION

- The occurrence of acute intestinal obstruction is more in small bowel.
- The incidence of intestinal obstruction is more common in males compared to females.
- Mode of presentation also differs in different levels of intestinal obstruction, small bowel obstruction mainly present with colicky abdominal pain and vomiting, as compared to large bowel obstruction where distention and constipation were predominant symptoms.
- Adhesions accounted for majority of small bowel obstruction (26%).
- Malignancies are common causes of large bowel obstruction.

- Tubercular stricture contributed 4% of all cases.
- Early recognition and timely intervention is important to prevent the bowel going for gangrenous changes.
- Other causes of acute intestinal obstruction in our series were obstructed hernias, volvulus, bands, tubercular stricture, Meckel's diverticulum, intussusceptions and meconium ileus.
- Prognosis was poor in elderly patients and newborns, in patients with co-morbid conditions, presence of strangulated bowel that required resection & anastomosis and those whose presentation to hospital was late.

- [51]. Asbun HJ, Perpinello C, Halasz NA. Small bowel Obstruction and its management. *Int surg* 1989; 74:2-27.
- [52]. 51. Bhansali S.K., Desai A.N & Dhabowala, C.B Tuberculous perforation of the small intestine. *J.Assoc.phys.India*, 16:351- 355,1968.

REFERENCES

- [1]. Winslet MC. Intestinal obstruction. In: Russel RCG, Williams NS, Bull strode CJK, editors. *Bailey & Loves Short practice of Surgery* 23rd edn. 2000. Edward Arnold Ltd NY. 1058-75.
- [2]. Jones RS. Intestinal obstruction. In: Sabiston DC Jr, editor. *Text book of surgery – The biological basis of modern surgical practice* 13th edn. 1986. W.B. Saunders company. 905-13.
- [3]. Oschner A. X-ray diagnosis of ileus – value of Roentgenogram in simple and strangulated obstruction. *Surg. Gynaecol. Obstet.* 1933; 56:719.
- [4]. Donhauser JL, Kelly EC. Intussusception in the adult. *AJS* 1950; 79: 673-77.
- [5]. Wangenstein OH, Rea CE, Smith BA, Schwytzer HC. Experiences with employment of suction in treatment of acute intestinal obstruction. *Surg. Gynecol. Obstet.* 1939; 68:851-68.
- [6]. Adesunkanmi AR, Agbakwuru EA. Changing pattern of acute intestinal obstruction in tropical African population. *E Afr Med J* 1996; 73(11): 727-31.
- [7]. Bass KN, Jones B, Bulkley GB. Current management of small bowel obstruction. *Adv surg* 1997; 31: 1-34.
- [8]. Chatterjee H, Somasekar SN, Ravishankar N, Madhu CP, Sidesh G, Vasantakumar SB. Adult intussusception. *IJS* 2000; 62(3): 210-12.
- [9]. Lopez-Kostner F, Hool GR, Lavery IC. Management and causes of acute large bowel obstruction. *Surg Clin North Am* 1997; 77(6): 1265-90.
- [10]. Fuzan M, Kay make E, Harmancioglu O, Astarcioğlu K. Principal causes of mechanical bowel obstruction in surgically treated adults in Western Turkey. *BJS* 1991; 78: 202-03.
- [11]. Matter I, Assaffy Y. Intussusception in adults. *S Afr Jr Surg* 1997; 35(4): 194-97.
- [12]. Rai S, Chandra SS, Smile SR. A study of risk strangulation and obstruction ingroin hernias. *Aust NZJ Surg* 1998; 68(9): 650-54.
- [13]. Sarkar PK, Sarkar V. Primary resection and anastomosis associated with maximal rectal stretching (MRS) for treatment of acute sigmoid volvulus. *IJS* 2000; 62(2): 122-24.
- [14]. Sourkati EO, Fahal AH, Succman SH et al. Intestinal Obstruction in Khartoum. *East Afr Med Jr* 1996; 73(5): 316-19.
- [15]. Gillespie IE. Small intestine and colon. In: Jamieson, Kay, editors.
- [16]. *Textbook of surgical physiology* 4th edn. Churchill Livingstone. 293-330.
- [17]. Feilding LP. *Textbook of intestinal obstruction*. Churchill Livingstone, NY. 1987. Vol. 13.
- [18]. Carty NJ, Ravichandran D. The management of malignant large bowel obstruction. In: Johnson CD, Taylor I, editors. *Recent advances in surgery* Vol 19. 1996. Churchill Livingstone. 1-16.
- [19]. Gill SS, Eggleston FC. Acute intestinal Obstruction. *Arch Surg* 1965; 91: 389-92.
- [20]. Ramachandran CS. Intestinal volvulus. *IJS* 1989; 66-70.
- [21]. Verma HN, Mishra AP, Prasad DN, Agarwal RKP. Volvulus of caecum. *IJS* 1977; 39: 252-56.
- [22]. Hjelmst A. Volvulus of right colon. *Acta chir Scand* 1959; 118: 455-65.
- [23]. Roggo A, Ottinger LW. Acute small bowel volvulus in adults. *Ann surg* 1992; (2): 135-141.
- [24]. Willets IE, Barclay GR, Banks RE, Rumley A, Allgar V, Stringer MD.
- [25]. Endotoxin, cytokines and lipid peroxides in children with intussusception. *BJS* 2001; 88: 878-83.
- [26]. Nagar RC. Double Intussusception of Jejunum in an adult. *Jr. Ind. Med. Asso.* 1990; 88(12): 340-41.
- [27]. Davis SE, Sperling L. Obstruction of Small Intestine. *Arch Surg* 1969; 99: 424-Fukuoka M, Tachibana S, Harada N, Saito H. Strangulated herniation through a defect in broad ligament. *Surg* 2002; 131: 232-33.
- [28]. Vander wolk WE, Snyder CA, Figg DM. Caecal-Colic adult intussusception as a cause of intestinal obstruction in central Africa. *World Jr. Surg* 1996; 20(3): 341-44.
- [29]. Lee SH, Ong ETL. Changing pattern of intestinal obstruction in Malaysia – a review of 100 consecutive cases. *BJS* 1991; 78: 181-182.
- [30]. Bollinger JA, Fowler EF. Results of treatment of Acute Small bowel Obstruction. *Arch Surg* 1952: 888-954.
- [31]. Dunn JA, Halls JM, Berne JV. Roentgenographic contrast studies in small bowel obstruction. *Arch Surg* 1984; 119: 1305.
- [32]. Begos DG, Sandar A, Modlin IM. The Diagnosis and Management of Adult Intussusception. *AJS* 1997; 173: 88-94.
- [33]. Lopez-Kostner F, Hool GR, Lavery IC. Management and causes of large bowel obstruction. *Surg. Clin. of North Amer.* 1997; 77: 77-86.
- [34]. Chubbuck A, Alponat A, Gonullu NN. Mitomycin-C prevents reformation of intra abdominal adhesions after adhesiolysis. *Surg* 2002; 131(1): 81-83.
- [35]. Tiwary RN, Prasad S. Mesocoloplasty for sigmoid volvulus: a preliminary report. *BJS* 1976; 63: 961-62. [36]. Anderson JR. Acute caecal volvulus. *BJS* 1980; 67: 39-41.
- [37]. Neufang T, Beeker H. Laparskopie bei duennadramileus. *Chirg* 2000; 71: 518-523.
- [38]. Michel ML Jr, Knapp L, Davidson A. Acute intestinal Obstruction; Comparative studies of small intestinal and colic obstruction. *Surg* 1950; 28: 90-110.
- [39]. Becker WF. Acute obstruction of colon – An analysis of 205 cases. *Surg. Gynaec. Obstet.* 1953; 96: 677-682.
- [40]. Budharaja SN, Govindarajulu S, Perianayagum WJ. Acute intestinal obstruction in Pondicherry. *IJS* 1976; 111-117.
- [41]. Jain BL, Prasad N. Intestinal obstruction. *IJS* 1963; 25: 635.
- [42]. Ti TK, Young NK. The pattern of intestinal Obstruction in Malaysia. *BJS* 1976; 63: 963-965. [43]. Sankaran V. Volvulus in South India. *IJS* 1962: 784-790.
- [44]. Mucha P, Jr. Small intestinal Obstruction. *Surg cli of North Am* 1987; 67(3): 597-618.
- [45]. Thompson RW, James BD, Mark MD. Obstructing adenocarcinoma of right side of the colon. *Arch surg* 1968; 96: 100-103.
- [46]. Hsu TC. Comparison of one stage resection and anastomosis of acute complete obstruction of left and right colon. *AJS* 2005; 189: 384-387.
- [47]. Kuruvilla MJ, Challen CR, Rajagopal AK, Rakas JS. Major causes of intestinal obstruction in Libya. *BJS* 1987; 74: 314-315.
- [48]. Sircar S, Taneja VA, Kanasara U. Clinical presentation of abdominal tuberculosis – A prospective study. *Jr IMA* 1996; 94(9): 342-344.
- [49]. Kappor VK, Gupta S, Sikora SS. Acute tubercular abdomen. *IJS* 1991; 53: 71.
- [50]. Grosfeld JL, West KW. Neonatal intestinal obstruction and Intussusception in childhood. In: Zinner MJ, Schwartz SI, Ellis H, editors. *Maingots abdominal operations* 10th edn. Prentice Hall International. INC. 2075-96.