



MYOMA PLACENTA RELATIONSHIP & PREGNANCY OUTCOME

Obstetrics & Gynaecology

Dr Savita Chandra Professor & Head of Department of Obstetrics & Gynaecology, Era's Lucknow Medical College & Hospital, Lucknow, India.

ABSTRACT

During the two years study period, 8557 deliveries took place, of which 114 were pregnancies with myoma giving the incidence as 1.33%. 12 of the 114 pregnancies ended in abortion, i.e. 10.5%. The myoma placenta relationship and the obstetric outcome was analyzed in the remaining 102 pregnancies which continued beyond 22 weeks. Of these, 41.2% had complications. Premature rupture of membranes was the commonest complication seen in 22.5%. No case of preeclampsia or antepartum haemorrhage occurred in the 'no contact' category. Amongst the cases where the placenta was either in contact or superimposed on the fibroid, 85.5% - 87.5% developed one or more complication. The myoma placenta relationship was found to be statistically significant, p value <0.001 . 39.3% were caesarean deliveries. Overall 37.25% were low birth weight neonates. There were two stillbirths, both in the superimposed category.

KEYWORDS

Myoma placenta relationship, pregnancy outcome

INTRODUCTION

The complexity of pregnancy is not fully understood, nor is understood the complexity of the two entities when they coexist that is one, the fibroid and second, the pregnancy. But what is known is the increasing prevalence of coexistence of myoma/fibroid with pregnancy; an apparent increase attributed to diagnostic imaging modalities and a true increase in myoma because of the changing life style of women.

The prevalence of myoma with pregnancy has been reported as 1.6 to 10.7%¹⁻⁵. The vitality and the health of the placenta is critical for both maternal and fetal outcome. Against this background, if a space occupying lesion like a myoma exists in the uterus, it could impact the contour, function and vasculature of the placenta as well as the distensibility of the uterus which in turn could affect the obstetric outcome. With this in mind, the study was planned.

The aim was to determine the prevalence of myoma in pregnancy, and the significance of the myoma placenta relationship, to the maternal & fetal outcome.

Study Design:

This was a two years prospective observational study carried out in the Department of Obstetrics & Gynecology, Goa Medical College. All consecutive cases of pregnancy with fibroid were enrolled. In each case the clinical and the sonographic findings were recorded.

Table 1: Myoma Placenta Relationship & Maternal Complications

MPR [*]	Total Cases	*Cases w/o Complications	APH	PROM	Pre-eclampsia	Malpr ^{**}	Dysfunct. Labor***	PPH	Cases with Complications	MPR Category %	Chi-Square Test
No contact	80	57	0	16	0	5	2	0	23	28.8%	Ref
Contact	14	2	1	6	3	1	1	0	12	85.7%	$\chi^2=16.54$, $P<0.001$
Super-imposed	8	1	1	1	3	0	1	1	7	87.5%	$\chi^2=11.17$, $P<0.001$
Total cases	102	60	2	23	6	6	4	1	42	41.2%	-
% (n=102)	NA	58.8%	2.0%	22.5%	5.9%	5.9%	3.9%	1.0%	41.2%	-	-

*w/o without: **malpresentation: ***dysfunctional labour

On analyzing the maternal outcome, (Table I) it was observed that there was no case of preeclampsia in the 'no contact' category; while 3 of the 14 cases (21.4%) in the 'contact' category and 3 of the 8 cases (37.5%) in the 'superimposed' category developed preeclampsia.

No antepartum haemorrhage (APH) was seen in the 'no contact' category, while it occurred in one of the 14 cases in the 'contact' category i.e. 7.14% and 1 of the 8 cases in the superimposed category i.e. 12.5%. Post partum haemorrhage (PPH) was seen only in the 'superimposed' category. Thus, the risk of APH & PPH was more in the 'contact' and 'superimposed' category.

Interestingly, of the 6 malpresentations in this series, 5 occurred in the 'no contact' category. Of the 23 cases of premature rupture of membranes (PROM), 16 occurred in the 'no contact' category.

Myometrial muscular contraction was excluded by repeat ultrasonography (USG) before enrolling those diagnosed as fibroid by USG in the first trimester. The subjects were followed till the termination of pregnancy.

The myoma placenta relation was categorized as:

a) No contact: Placenta and fibroid were not in contact with each other.

b) Contact: Placenta only partially covering the fibroid.

c) Superimposition: Placenta completely covering the fibroid.

Booked, and referred cases were included. Termination of pregnancy before 22 weeks /fetus weighing less than 500gms was considered as an abortion; beyond 22 weeks to less than 37 weeks was considered as preterm; 37 weeks to 42 weeks was considered as term.

OBSERVATIONS AND RESULTS

During the two years study period, 8557 deliveries took place, of which 114 were pregnancies with fibroids giving the incidence of coexistent fibroid with pregnancy as 1.33%. 12 of the 114 pregnancies i.e. 10.5% ended in abortion. In 102 women pregnancy continued beyond 20 weeks of gestation and their maternal and fetal outcome was analyzed.

Of the 102 pregnancies, in 80 there was no contact between the fibroid and the placenta, in 14 there was contact and in 8 the placenta was superimposed on the fibroid. There was no case of multiple fibroids.

Overall, of the 102 pregnancies, 60 i.e. 58.8% had no maternal complications while 42 cases (41.2%) had maternal complications. 23 out of 80 cases (28.8%) in the 'no contact' category, 12 of the 14 cases (85.7%), in the 'contact' category, and 7 of the 8 cases (87.5%), in the 'superimposed' category developed complications, the difference reached statistical significance, p value 0.001 (Table I).

Table II: Mode of Delivery

Mode of delivery	Total Cases	% (n=102)
Spontaneous Vaginal Delivery	59	58%
Instrumental Vaginal Delivery	3	3%
LSCS	40	39%
Classical Caesarean Section	-	-
Total	102	100%

Of the 102 pregnancies 40 (39.2%) were caesarean deliveries while 62 delivered vaginally. Of the latter, 59 (58%) delivered spontaneously while 3 (3 %) were instrumental vaginal deliveries (Table II) There was no maternal mortality in this series.

Table III: Myoma Placenta Relationship And Weight Of The Newborn

Myoma placenta Relation Category	No. of Cases	Birth weight > 2.5 kg		Birth Weight <2.5 kg		Chi Square Test
		No.	% of Category	No.	% of Category	
No Contact	80	61	76.25%	19	23.75%	Referen ce
Contact	14	3	21.43%	11	78.57%	$\chi^2=16.48$ $p<0.001$
Superimposed	8	-	-	8	100.00%	$\chi^2=19.88$ $p<0.001$
Total Cases % of Total	102 100%	64 62.75%	-	38 37.25%	-	

It is seen from Table III that 38 of the 102 neonates (37.25%) were low birth weight babies. In the 'superimposed' category all 8 neonates (100%) were low birth weight of which two were stillbirths; 11 of the 14 neonates (78.57%) in the 'contact' category while 19 of the 80 babies (23.75%) in the 'no contact' category were low birth weight neonates which was statistically significant, p value <0.001.

DISCUSSION

The exact prevalence of coexisting myoma and pregnancy is not exactly known. The prevalence has been reported as 1.6 to 10.7¹⁻³ while Sheiner et al⁶ have reported the prevalence as 0.1% to 4%.

In our study the incidence of myoma with pregnancy was 1.33% closely comparable to 1.4%. reported by Benson et al⁷. However, our incidence was higher when compared to 0.37% reported by Coronado et al⁸, who attributed their low incidence to under reporting.

The present study showed, that pregnancy with a myoma, was a high risk pregnancy as 42 of the 102 cases (i.e.41.2%) had one or more complications while only 58.8% were without complication (Table I).

With regard to the type of maternal complications, we observed that the myoma placenta relationship was of significance. In our series, APH and preeclampsia occurred where the placenta was either in contact with the fibroid or was superimposed. Similarly, Exacoutous et al² found that 71% of the cases with abruptio placenta had a myoma with placental contact or superimposition in their series. Coronado et al⁸ opined that the risk of abortion of the placenta was four times more when the fibroid was retroplacental. Interestingly, in our study, the majority of PROM occurred in the 'no contact' category i.e. 16 of the 23 cases giving the incidence of PROM as 22.5% (Table I) Zaima et al¹⁰ found 19% risk of PROM which is closely comparable to our result. It is hypothesized that possibly the altered localized distensibility and the mechanical effect of the myoma might be the underlying mechanisms for the increased risk of PROM.

Myomas have been reported to be associated with 10-40% antenatal complications⁹. We found complications in 41.2% of the cases and excluding PPH, 40.2% antenatal complications (Table I).

The Caesarean section rate in our study was 39.2% (Table II). Benson et al⁷ reported a Caesarean section rate of 38% which is closely comparable to our study. Zaima et al¹⁰ found uterine fibroids to be associated with 49% risk of caesarean delivery. Klatsky et al, in 2008, reported that women with fibroids had 3.7 fold increased risk of caesarean delivery¹¹.

Most studies report increased rate of preterm deliveries¹². In this study, 38 of the 102 newborns were low birth weight babies (< 2.5 Kg) of which 17 were preterm i.e. 44.7% of the 102 neonates (Table III).

The present study showed that the myoma placenta relationship is an important determinant of maternal and fetal outcome and that pregnancy with myoma has a higher chance of adverse obstetric outcome.

Limitation of the study was that amongst the enrolled consecutive cases of pregnancy with fibroid there was no case of multiple fibroids or multiple gestation and also it was an observational study.

CONCLUSION

The coexistence of a fibroid, adds a new dimension to the complexity and dynamics of pregnancy since the uterus gets shared between the two entities. This coexistence of the fibroid and pregnancy could alter the dynamics within the endometrium, the decidua, the molecular signaling, the vascular architecture, yet not fully understood and need further research for unravelling the complexities.

Funding: No funding sources

Conflict Of Interest: None declared

Ethical Approval: Approval obtained

REFERENCES

1. Qidwai GI, Caughey AB, Jacoby AF. Obstetric outcomes in women with sonographically identified uterine leiomyomata. *Obstet Gynecol* 2006; 107:376.
2. Exacoustos C, Rosati P. Ultrasound diagnosis of uterine myomas and complications in pregnancy. *Obstet Gynecol* 1993; 82:97.
3. Strobelt N, Ghidini A, Cavallone M, et al. Natural history of uterine leiomyomas in pregnancy. *J Ultrasound Med* 1994; 13:399.
4. Laughlin SK, Baird DD, Savitz DA, et al. Prevalence of uterine leiomyomas in the first trimester of pregnancy: an ultrasound-screening study. *Obstet Gynecol* 2009; 113:630.
5. Stout MJ, Odibo AO, Graseck AS, et al. Leiomyomas at routine second-trimester ultrasound examination and adverse obstetric outcomes. *Obstet Gynecol* 2010; 116:1056.
6. Scheiner E., Bashiri A., Levy A., HersHKovits R., Katz M., and Mazor M.:Obstetric characteristics and perinatal outcome of pregnancies with uterine leiomyoma. *J Repro Med* 2004;49:182-186.
7. Benson CB, Chow JS, Chang-Lee W, et al. Outcome of pregnancies in women with uterine leiomyomas identified by sonography in the first trimester. *J Clin Ultrasound*. 2001;29:261-264. [PubMed].
8. Coronado GD, Marshall LM, Schwartz SM. Complications in pregnancy, labor, and delivery with uterine leiomyomas: a population-based study. *Obstet Gynecol*. 2000;95:764-769. [PubMed] [Google Scholar].
9. Cook H., Ezzati M., Segars J.H., McCarthy K.. The impact of uterine leiomyomas on reproductive outcomes. *Minerva Ginecol*. 62(3): 2010; 225- 236 Web of Science®Google Scholar
10. Zaima A, Ash A. Fibroid in pregnancy: Characteristics, complications, and management. *Postgrad Med J*. 2011;87(1034):819-28.
11. Klatsky PC, Tran ND, Caughey AB, Fujimoto VY. Fibroids and reproductive outcomes: a systematic literature review from conception to delivery. *Am J Obstet Gynecol*. 2008;198:357-366. [PubMed]
12. Lai J, Caughey AB, Qidwai GI, Jacoby AF. Neonatal outcomes in women with sonographically identified uterine leiomyomata. *J Matern Fetal Neonatal Med* 2012; 25:710. 37