



STUDY OF DETERMINE THE RELATIONSHIP BETWEEN C-REACTIVE PROTEIN (CRP) AND ERYTHROCYTE SEDIMENTATION RATE (ESR) WITH GALL BLADDER ADHESION ON CHOLECYSTECTOMY

Surgery

Dr. Om Prakash* MBBS, M.S. (Gen. Surgery), Senior Resident, Department of Surgery, All India Institute of Medical Sciences (AIIMS), New Delhi. *Corresponding Author

Dr. [Prof.] Rajinder Parshad M.S. (Gen. Surgery), Professor, Department of Surgery, All India Institute of Medical Sciences (AIIMS), New Delhi.

ABSTRACT

Background: The conversion from laparoscopic cholecystectomy to open cholecystectomy often results from pericholecystic Adhesion. Pericholecystic Adhesion is a result of cholecystitis in the case of cholelithiasis. C-Reactive Protein (CRP) and Erythrocyte Sedimentation Rate (ESR) can be increased in cholecystitis and used as a predictive factor for the degree of difficulty in laparoscopic cholecystectomy. This study aims to determine the relationship between CRP and ESR with gall bladder adhesion on cholecystectomy at AIIMS, Delhi.

Method: A cross-sectional study was conducted among 61 respondents who underwent a cholecystectomy surgery at All India Institute of Medical Sciences, New Delhi during July-December 2019. Parameters assessed in this study were Gender, diagnosis, CRP, ESR, age, hemoglobin, leukocytes count, platelet count, and the relationship between CRP and ESR value with the pericholecystic adhesions. Data were analyzed by SPSS version 22 for Windows.

Results: Most of respondents were female (70.5%), followed by cholelithiasis (73.8%) as primary diagnosis, no Pericholecystic Adhesions found (68.9%), CRP level < 23 mg/dl (77.0%), and ESR level \geq 11 mm/hour (78.7%). In addition, the average age of respondents was 48.44 \pm 10.85 years old, followed by the hemoglobin level (12.43 \pm 1.77 g/dL), leukocytes counts (9,439.67 \pm 3,719.37/ μ l), and platelet counts (319,295.08 \pm 90,587.59/ μ l). CRP level had a significant relationship to the occurrence of pericholecystic Adhesion (OR: 75.400; 95%CI: 7.991- 711.441; p=0.000).

Conclusion: It can be concluded that there was a relationship between CRP and ESR with the occurrence of pericholecystic Adhesion. However, based on multivariate analysis, CRP was more significant compared to ESR on the occurrence of pericholecystic Adhesion.

KEYWORDS

CRP, ESR, Pericholecystic Adhesion, Cholelithiasis, Cholecystectomy

INTRODUCTION

Cholelithiasis is one of the most common diseases that bring patients to the emergency department with abdominal disorders such as epigastric pain, nausea, vomiting, and anorexia. Although the mortality rate of cholelithiasis is low, the economic burden and the impact of health on it is very significant for health care systems around the world due to high morbidity.

About 20% of people with cholelithiasis experience symptoms and can cause complications, such as cholecystitis, cholangitis, obstructive jaundice, and pancreatitis. Cholecystitis is inflammation of the gallbladder whereas 90% of cholecystitis patients are caused by cholelithiasis.

Ultrasonography (USG) is a critical first step and examination for the diagnosis of acute cholecystitis, with a sensitivity of 80-100% and a specificity of 60- 100%. Besides that, leukocyte levels, C-Reactive Protein (CRP), and erythrocyte sedimentation rate (ESR) are also used as laboratory findings to support the diagnosis of acute cholecystitis.

CRP is an acute-phase reactant that increases when inflammation occurs. CRP levels begin to rise after 6-8 hours, depending on the severity of inflammation, and often used in estimating the severity of inflammation, especially intraabdominal. Based on the 2018 Tokyo Guidelines, one of the criteria for the diagnosis of acute cholecystitis is increased CRP levels.

Erythrocyte sedimentation rate (ESR) is a general hematology test, where red blood cells settle in a standard tube within one hour. ESR levels are increased in cases of inflammation, pregnancy, anemia, autoimmune disorders (rheumatoid arthritis and lupus). ESR can also be increased in cases of acute cholecystitis and is used as a predictive factor for the difficulty of laparoscopic cholecystectomy. Based on those mentioned above, this study aims to evaluate the relationship between CRP and ESR with gall bladder adhesion on cholecystectomy.

MATERIAL AND METHODS

A cross-sectional study was conducted to determine the relationship between C-Reactive Protein (CRP) and Erythrocyte Sedimentation Rate (ESR) as a predictive factor for the degree of pericholecystic adhesions in cholecystectomy.

There were 61 eligible subjects enrolled in this study through the consecutive technique at All India Institute of Medical Sciences, New Delhi.

The inclusion criteria used in this study were patients with cholelithiasis who underwent cholecystectomy during July-December 2019 period. Parameters used in this study include gender, diagnosis, the occurrence of pericholecystic adhesions, CRP, and ESR levels. All of the data were presented in mean, standard deviation, number, percentage, minimum, and maximum value.

Data analysis was performed using SPSS version 22. In analyzing CRP and ESR, if the data is normally distributed, the analysis is carried out with the Anova test and continued with the Posthoc test with the Tukey method. However, the difference analysis is carried out by the Kruskal-Wallis test. Data were determined as significant if the p-value less than 0.05.

RESULTS

A total of 61 respondents were enrolled in this study. Most of respondents were female (70.5%), cholelithiasis as primary diagnosis (73.8%), no pericholecystic adhesions (68.9%), CRP level < 23 mg/dl (77.0%), and ESR level \geq 11 mm/hour (78.7%) (Table 1). Besides, the average age of the sample was 48.44 \pm 10.85 years old, with the oldest age was 73 years, and the youngest was 23 years (Table 1). Most of the respondents have the average hemoglobin level about 12.43 \pm 1.77 g/dl, followed by leukocyte counts (9,439.67 \pm 3,719.37/ μ l), platelet counts (319,295.08 \pm 90,587.59/ μ l), and 19.69 \pm 11.83 mm/hour for ESR levels (Table 1).

Table 1 : Baseline Characteristics Of Respondents

Variable	Respondents (n=61)			
	N(%)	Mean \pm SD	Minimum	Maximum
Gender				
• Male	18(29.5)			
• Female	43(70.5)			
Diagnosis				
• Cholelithiasis	45(73.8)			
• Multiple Cholelithiasis	15(24.6)			
• Cholecystitis Cholelithiasis	1(1.6)			
Pericholecystic Adhesions				
• Yes	19(31.1)			
• No	42(68.9)			

C-Reactive Protein (CRP) (mg/dl)			<6	35
• <23 mg/dl	47(77.0)			
• ≥23mg/dl	14(33.0)			
Erythrocyte Sedimentation Rate (ESR) (mm/hour)		19.69±11.83	8	37
• <11 mm/hour	13(21.3)			
• ≥11 mm/hour	48(78.7)			
Age (years)		48.44±10.85	23	73
Hemoglobin (g/dl)		12.43±1.77	8	16
Leukocytes (/ml)		9,439.67±3,719.37	4,600	25,500
Platelets (/ml)		319,295.08±90,587.59	148,000	528,000

Base on the risk analysis evaluation, the recent study found a statistically significant relationship between ESR (OR: 7.200; 95%CI: 0.862-60.107; p=0.047) and CRP (OR: 88.883; 95%CI: 9.774-807.362; p=0.000) levels to the pericholecystic Adhesion (Table 2). The odds ratio shows that the level of ESR ≥ 11 has the chance of Adhesion 7.200 times as well as CRP level ≥ 23 was likely to have adhesion of 88.883 times (Table 2).

Table 2 : The ESR And CRP Variables Are Related To Pericholecystic Adhesion

Variable	Pericholecystic Adhesion		p	OR	95%CI
	Yes (n=61)	No (n=61)			
ESR, n(%)					
≥11	18(29.5)	30(49.2)	0.047*	7.200	0.862-60.107
<11	01(1.6)	12(19.7)			
CRP, n(%)					
≥23	13(21.3)	01(1.6)	0.000*	88.883	9.774-807.362
<23	06(9.8)	41(67.2)			

*Fisher's Exact Test (p < 0,05); OR: odds-ratio; ESR: Erythrocyte Sedimentation Rates; CRP: C-Reactive Protein

Multivariate analysis found that the CRP variable has a significant relationship compared to the ESR variable on the occurrence of pericholecystic Adhesion (p<0.05). In addition, the odds ratio was also found higher compared with other variables as a risk of pericholecystic Adhesion on the CRP variable (≥23) (Table 3).

Table 3 : The Multivariate Analysis Of Several Variables To The Risk Of Pericholecystic Adhesion

Variable	p	OR	95% CI for EXP(B)	
			Lower	Upper
ESR (≥11)	0.527	2.069	0.218	19.629
CRP (≥23)	0.000*	75.400	7.991	711.441

DISCUSSION

In this study, the total sample studied was 61 patients with a mean age of 48.44±10.85 years old with the oldest age of 73 years and the youngest of 23 years. The gender proportion between males (29.5%) and females (70.5%) shows more cases in women. In accordance with the literature states that females and old age are risk factors that cannot be modified. Based on the pericholecystic adhesion, our study found that about 31.1% respondents were affect. This result is quite different from a study in Surabaya that received 75.6% pericholecystic adhesion.

From this study, it was found that there was a significant relationship between CRP and ESR with pericholecystic adhesion (p<0.05) during cholecystectomy. The odds ratio shows that the level of ESR ≥ 11 has a chance of adhesion 7.2 times. Whereas in the CRP variable, there was a significant relationship between CRP and pericholecystic adhesion during cholecystectomy and showed that the CRP level ≥ 23 was likely to have adhesion of 88.8 times.

CRP variable has a significant effect compared to the ESR variable on the occurrence of pericholecystic Adhesion. Then the odds ratio was found to be a greater chance of pericholecystic Adhesion to the CRP variable (≥ 23), which is 75.4 times. CRP is an acute-phase protein with a half-life of 19 hours. CRP is synthesized and released by the liver in response to Interleukine-6 and other proinflammatory cytokines. CRP activates the classic cascade complement and stimulates phagocytosis.

Increased concentrations of CRP in serum can occur in infections, inflammation, trauma, malignancies, and tissue infarction. not specific for certain diseases.

One of the first studies evaluating CRP as a predictor of the severity of inflammation in cholecystectomy (both open and laparoscopic) with OR=0.979 (95% CI: 0.969-0.989). A previous study by Jessica Mok KW et al., evaluated the CRP values in a retrospective cohort as parameters that predict difficult laparoscopic cholecystectomy or conversion and also illustrate CRP cut-off points to predict conversion. They found that patients with CRP ≤ 220 (3.2%) had far less risk of conversion compared to CRP> 220 (61.9%) (P <. 001). In a multiple logistic regression analysis to predict the level of difficult dissection operations, it was found that the OR value of CRP was 1,007 (95% CI: 1,001- 1,012).

Significantly higher CRP, with a CRP level > 200 mg/dL, has a sensitivity of 50% positive and 100% negative as a predictive value for gangrene cholecystitis. In another study, absolute values (CRP> 190 mg/dL) were predictive factors for difficult cholecystectomy laparoscopy; even changes in the CRP interval (increasing> 90 mg/dL) at 48 hours from hospitalization can predict the possibility of pancreatitis, as a complication.

ESR is a general hematology test, where red blood cells settle in a standard tube within one hour. ESR levels are increased in cases of inflammation, pregnancy, anemia, autoimmune disorders (rheumatoid arthritis and lupus). With ESR results >65 mm/hour (specific 66%, sensitivity 70%) found to be statistically significant in laparoscopic cholecystectomy operations which underwent conversion.

CONCLUSION

From this study, it can be concluded that there is a relationship between CRP and ESR with the occurrence of pericholecystic Adhesion. It can also be concluded that CRP has more influence on the occurrence of pericholecystic Adhesion. By using CRP and ESR laboratory results, it is hoped that the surgeon can estimate the presence or absence of pericholecystic Adhesion in cholelithiasis patients.

A surgeon can use the results of the CRP and ESR to predicting the difficulty level of laparoscopic cholecystectomy so that the surgeon can consider the choice of management modality according to his capabilities. By predicting the degree of difficulty in cholecystectomy laparoscopic surgery, a surgeon can educate patients about the difficulty of the operation and the possibility of conversion.

REFERENCES

- Amri RA, Akbar FN, Hendarto H. Proporsi Batu Empedu terhadap Status Gizi (Obesitas) di RSUD Fatmawati Tahun 2015-2016 [Skripsi]. UIN Syarif Hidayatullah: Fakultas Kedokteran dan Ilmu Kesehatan: Jakarta. 2017.
- Balta AY, Suculli I, Filiz AI, Yucel E, Ozdemir Y, Kurt Y. Single Incision Laparoscopic Cholecystectomy: Result of Our First Experience. *Gulhane Med J*. 2011;53(2):136-138.
- Baskara AS, Budiono BP, Wahyudi F. Perbandingan Teknik Operasi Mini Laparotomi Kolesistektomi dengan Laparoskopi Kolesistektomi pada Kolesistolitiasis terhadap Lama Rawat Inap di RSUD Dr. Kariadi Semarang. *Jurnal Kedokteran Diponegoro*. 2014;3(1):1-14.
- Bodman-Smith KB, Melendez AJ, Campbell I, Harrison PT, Allen JM, Raynes JG. C-reactive protein-mediated phagocytosis and phospholipase D signalling through the high-affinity receptor for immunoglobulin G (FcγmAR1). *Immunology*. 2002;107(2):252-260.
- Cole DS, Watts A, Scott-Coombes D, Avades T. Clinical utility of peri-operative C-reactive protein testing in general surgery. *Ann R Coll Surg Engl*. 2008;90(4):317-321.
- Dewi MMW, Herawati S, Mulyantari NK, Prabawa IPY. The comparison of erythrocyte sedimentation rate (ESR) modify Westergren Caretium Xc-A30 and Westergren Manual in Clinical Pathology Laboratory, Sanglah General Hospital, Denpasar, Bali. *Bali Medical Journal*. 2019;8(2):396-399
- Girsang JH, Hiswani, Jemadi. Karakteristik Penderita Koleslitiasis yang dirawat inap di Rumah Sakit Santa Elisabeth Medan pada tahun 2010-2011. *Gizi, Kesehatan Reproduksi dan Epidemiologi*. 2012;1(1):1-9.
- Hung SC, Liao KF, Lai SW, Li CL, Chen WC. Risk factors associated with symptomatic cholelithiasis in Taiwan: a population-based study. *BMC Gastroenterol*. 2011;11:111.
- Jessica Mok KW, Goh YL, Howell LE, Date RS. Is C-reactive protein the single most useful predictor of difficult laparoscopic cholecystectomy or its conversion? A pilot study. *J Minim Access Surg*. 2016;12(1):26-32.
- Lauwrens SS, Mellow PA, Waworuntu GL, Suyanto A. Hubungan Leukositosis dengan Pelekatkan Kandung Empedu pada Pasien Kolesistektomi di Rumah Sakit X [Skripsi]. Fakultas Kedokteran Universitas Katolik Widya Mandala: Surabaya. 2017.
- Litao MK, Kamat D. Erythrocyte sedimentation rate and C-reactive protein: how best to use them in clinical practice. *Pediatr Ann*. 2014;43(10):417-420.
- Littlefield A, Lenahan C. Cholelithiasis: Presentation and Management. *J Midwifery Womens Health*. 2019;64(3):289-297.
- Pinto A, Reginelli A, Cagini L, Coppolino F, Ianora AAS, Bracale R, et al. Accuracy of ultrasonography in the diagnosis of acute calculous cholecystitis: review of the literature. *Crit Ultrasound J*. 2013;5 Suppl 1(Suppl 1):S11.
- Purwanti A, Maliya A, Zulaiha E. Hubungan Gaya Hidup dengan Kejadian Penyakit Cholelithiasis di Ruang Rawat Inap RSI Surakarta [Skripsi]. Fakultas Ilmu Kesehatan Universitas Muhammadiyah Surakarta: Surakarta. 2016.
- Puzianowska-Kuznicka M, Owczarz M, Wiecezowska-Tobis K, Nadrowski P, Chudek J, Slusarczyk P, et al. Interleukin-6 and C-reactive protein, successful aging, and

- mortality: the PolSenior study. *Immun Ageing*. 2016;13:21.
16. Schafer M, Krahenbuhl L, Buchler MW. Predictive Factor for the type of surgery in acute cholecystitis. *The American Journal of Surgery*. 2001;182(3):291-297.
 17. Schirmer BD, Winters KL, Edlich RF. Cholelithiasis and cholecystitis. *J Long Term Eff Med Implants*. 2005;15(3):329-338.
 18. Sproston NR, Ashworth JJ. Role of C-Reactive Protein at Sites of Inflammation and Infection. *Front Immunol*. 2018;9:754.
 19. Stinton LM, Shaffer EA. Epidemiology of gallbladder disease: cholelithiasis and cancer. *Gut Liver*. 2012;6(2):172-187.
 20. Stirling AD, Moran NR, Kelly ME, Ridgway PF, Conlon KC. The predictive value of C-reactive protein (CRP) in acute pancreatitis - is interval change in CRP an additional indicator of severity?. *HPB (Oxford)*. 2017;19(10):874-880.
 21. Temizi A, Ozdemir Y, Aslan A, Bostanci MT, Atasoy G, Caparlar MA. Role of Complete Blood Counts Parameters in Diagnosis of Acute Cholecystitis. *Acta Medica Mediterranea*. 2017;3:411-316.
 22. Yokoe M, Hata J, Takada T, Strasberg SM, Asbun HJ, Wakabayashi G, et al. Tokyo Guidelines 2018: diagnostic criteria and severity grading of acute cholecystitis (with videos). *J Hepatobiliary Pancreat Sci*. 2018;25(1):41-54.