

ASSOCIATION OF SERUM MAGNESIUM LEVEL WITH RESISTANT HYPERLIPIDEMIA IN DIABETIC AND HYPERTENSIVE PATIENTS

Medicine

Dr. Manish Kumar	MBBS, M.D. (Med.), Senior Resident, Department of Medicine, Darbhanga Medical College & Hospital, Laheriasarai, Bihar.
Dr. Syed Yousuf Faisal*	MBBS, M.D. (Med.), Senior Resident, Department of Medicine, Darbhanga Medical College & Hospital, Laheriasarai, Bihar. *Corresponding Author
Dr. Sudhir Chandra Jha	Associate Professor, Department of Medicine, Darbhanga Medical College & Hospital, Laheriasarai, Bihar.
Dr. Debarshi Jana	Young Scientist (DST), Institute of Post-Graduate Medical Education and Research, A.J.C. Bose Road, Kolkata-700020, West Bengal, India.

ABSTRACT

Introduction: Both diabetes mellitus and hypertension are aspects of metabolic syndrome.

Objectives: The aim of this study was to determine the relationship between serum magnesium level with resistant hyperlipidemia in a group of diabetic and hypertensive patients.

Patients And Methods: The present cross-sectional study was carried out on 90 hypertensive and diabetic patients in Medicine Department of DMCH, Laheriasarai, Bihar, (45 hypertensive and 45 diabetic patients). Included patients had high triglyceride levels despite 8 weeks of treatment with lipid-lowering agents.

Results: There was an inverse significant relationship between serum magnesium and triglyceride levels in diabetic patients ($P=0.002$, $r=-0.458$), however, this correlation was not significant in hypertensive patients ($P=0.754$, $r=0.048$).

Conclusion: This study showed, serum magnesium may affect triglycerides levels in diabetic patients, however, our finding requires further investigation with larger population.

KEYWORDS

INTRODUCTION

Diabetes mellitus, commonly, is known to be a metabolic disease characterized by high blood sugar levels over a long-period of time. There are many lipoprotein disorders in patients with uncontrolled diabetes. These metabolic disorders are due to the excessive production of triglyceride-rich lipoproteins of the liver, which is due to low levels of high-density lipoproteins (HDL). These patients also suffer from hypertriglyceridemia, which is due to a decrease in adipose tissue and lipoprotein lipase activity in the muscle tissues. Reduced HDL-C levels are due to plasma lipoprotein lipase (LPL) in catabolism of triglyceride-rich lipoproteins.

Hypertension is a chronic disease, in which arterial blood pressure rises. High blood pressure, can result in significant problems and increases the risk of cardiac and renal disease, stroke and also death.

Hyperlipidemia is common in patients with hypertension, but the causes of this condition are unknown. Additionally, treatment of hypertension with thiazide diuretics increases the risk of hyperlipidemia.

Magnesium (Mg) is the fourth most abundant cation in the body, as well as the second intracellular cation after potassium. For many enzymes involved in the metabolism of carbohydrates and lipids, Mg is an important cofactor. Hypomagnesemia is one of the most common electrolyte disorders in diabetic patients. Numerous studies have shown a relationship between magnesium deficiency with insulin resistance, diabetes, and hyperlipidemia. Additionally, epidemiological studies have shown an inverse relationship between magnesium intake and the level of blood pressure and the incidence of hypertension. It was also shown that hypertensive patients have lower serum and intracellular magnesium than normal people. Mg plays an important role in lipoprotein lipase enzyme activity and removes triglyceride from blood. It is also essential for the activity of the enzyme, lecithin cholesterol acyltransferase (LCAT), which converts free cholesterol into cholesterol esters and causes liver inflammation. Pyrophosphatase, an enzyme involved in lipid metabolism, also requires Mg for its activity. Therefore, in magnesium deficiency, enzymes are not properly efficient and accordingly dyslipidemia may happen. Recently it was shown that normal population with lower Mg levels, had lower HDL-C, higher triglyceride (TG), and also total cholesterol serum values.

Hyperlipidemia increases the incidence of cardiovascular events and subsequent mortalities in patients. Lipid abnormalities such as lowered serum HDL-C levels and increased triglyceride blood levels are common in type 2 diabetes. Accordingly, lipid abnormalities in type 2 diabetes and hypertensive patients, can be considered as a cause for development of atherosclerosis. The relationship between magnesium and cardiovascular risk factors has been proven in previous studies, however, there are many contradictory findings regarding the effect of magnesium on lipid profiles, as well as the control of blood sugar in diabetic patients.

OBJECTIVES

Several studies have also proven the relationship between magnesium and hypertension, however few studies have been conducted on this subject. Serum magnesium is not commonly measured in diabetic and hypertensive patients, while recent investigation showed various metabolic complications have been attributed to hypomagnesemia. Hence this study was designed to detect the relationship between plasma magnesium levels and lipid profiles in diabetic and hypertensive individuals.

MATERIAL AND METHODS

Study Design

The present cross-sectional study was carried out on 90 hypertensive or diabetic patients at Department of Medicine, Darbhanga Medical College and Hospital, Laheriasarai, Bihar from June 2019 to May 2020 (45 hypertensive patients and 45 diabetic patients). Patients had high triglyceride levels despite eight weeks of treatment with lipid-lowering agents. Samples were gradually selected using convenience sampling method. Blood pressure was measured and recorded using an appropriate cuff under standard conditions. A total of 5 cc blood samples was taken from the patients after obtaining their consent letter, then blood cholesterol, triglyceride, HDL-C, low density lipoprotein (LDL-C), fasting blood glucose, and serum magnesium level were measured by standard methods.

Statistical Analysis

Collected, data entered in SPSS version 18 and analyzed using descriptive statistics, Pearson's or Spearman correlation's coefficient test, and also t test according to the distribution of data. Quantitative variables were investigated for normal distribution using Kolmogorov-Smirnov test. All statistical differences were considered significant at the level of P value lower than 0.05.

RESULTS

Out of 90 participants, 45 patients (50%) were included in group 1 (diabetic patients) and 45 (50%) in group 2 (hypertensive patients). The mean±SD of age, triglyceride, cholesterol, HDL-C, LDL-C, and serum magnesium of all participants was 60.41±16.08 years, 216.76±94.51 mg/dL, 281.34±38.31 mg/dL, 38.53±11.15 mg/dL, 201.04±44.93 mg/dL and 1.97±0.45 mg/dL, respectively. We found no statistically significant relationship between magnesium and cholesterol levels in total patients ($P=0.146$, $r=-0.155$). Additionally, the relationship of magnesium with HDL-C ($P=0.228$, $r=-0.12$), and LDL-C ($P=0.751$, $r=-0.34$) was also not significant in all cases. Spearman's correlation analysis also showed no statistically significant relationship between magnesium level and fasting blood glucose in diabetic patients ($P=0.342$, $r=0.145$). Similarly, no significant relationship between serum magnesium with systolic and diastolic blood pressure in group 2 was detected ($P=0.893$, $r=0.21$ and $P=0.347$, $r=-0.143$ respectively).

We also found, a significant correlation between serum magnesium and triglyceride level in group 1 ($P=0.002$, $r=-0.458$; Spearman's test, Figure 1), since in group 2 this association was not significant ($P=0.75$, $r=0.048$).




Figure1. Significant correlation between serum magnesium and triglyceride level in diabetic patients ($P=0.002$, $r=-0.458$; Spearman's test).

Table 1 : Data Of The Patients

	Group	N	Mean	Std. Deviation	Std. Error Mean
Age (years)	Diabetes	45	61.56	15.494	2.310
	Hypertension		59.27	16.755	2.498
TG (mg/dL)	Diabetes	45	204.27	78.639	11.723
	Hypertension		229.24	107.548	16.032
Cholesterol (mg/dL)	Diabetes	45	282.18	37.901	5.650
	Hypertension		280.51	39.141	5.825
HDL-C (mg/dL)	Diabetes	45	35.67	8.868	1.322
	Hypertension		41.40	12.503	1.864
LDL-C (mg/dL)	Diabetes	45	206.69	45.206	6.739
	Hypertension		195.40	44.451	6.626
Mg (mg/dL)	Diabetes	45	2.0916	0.47487	0.07079
	Hypertension		1.8629	0.40823	0.06086

Likewise, no significant correlation between serum magnesium and cholesterol in group one ($P=0.11$, $r=-0.238$) and group two ($P=0.59$, $r=-0.081$) was detected. Accordingly, the association of serum magnesium with LDL-C in group one ($P=0.91$, $r=-0.16$) and group two ($P=0.4$, $r=-0.12$) was not significant. Besides HDL-C in group one ($P=0.64$, $r=-0.071$) and group two ($P=0.47$, $r=-0.1$) were not associated with serum magnesium.

DISCUSSION

The present study was carried out on 90 hypertensive and diabetic patients (45 hypertensive patients and 45 diabetic patients) with had high triglyceride levels despite a treatment period of 8 weeks with lipid-lowering drugs. In this study we found, plasma magnesium had relationship with serum triglyceride level in diabetic patients. However, the serum magnesium level does not correlate with fasting

blood glucose, cholesterol, HDL-C, or LDL-C levels. Our results also showed that serum magnesium levels were not correlated with level of lipids or systolic and diastolic blood pressure in hypertensive patients. Diabetes mellitus is the most common disorder among endocrine disorders, which is associated with hypomagnesemia. Numerous studies have already shown that Mg levels are lower in diabetic patients. Reduced Mg level may cause insulin resistance and insufficient glucose control, and also dyslipidemia in diabetic patients. Previous studies have shown a significant negative correlation between Mg and fasting blood glucose (FBS). However, our study didn't show a significant correlation between Mg and FBS, which requires further larger studies. Marhale et al concluded that diabetes, dyslipidemia, and hypertension were inversely related to serum Mg, however we found no significant correlation between serum Mg with LDL-C, HDL-C, and triglyceride levels in hypertensive patients. We found, a significant correlation between serum Mg and triglyceride levels in diabetic patients. This finding is consistent with the result of a study by Arpacı et al. Various studies have suggested a direct relationship between Mg level and systolic blood pressure, however, results of the present study did not show this relationship while further studies are necessary for this reason.

CONCLUSION

Results of the current study showed that magnesium (Mg) can affect triglycerides levels in diabetic patients suggesting an inverse relationship between serum magnesium and triglyceride levels. However, there is no relationship between the magnesium serum level with fasting blood glucose, cholesterol, HDL-C, or LDL-C levels. Moreover, there is no relationship between serum magnesium levels and the level of lipid profiles or to the level of systolic or diastolic hypertension in hypertensive patients. Considering the effect of serum magnesium level on reducing triglyceride levels in diabetic patients indicated in this study and the need for correction of lipid profiles in these patients, the administration of magnesium supplements in the treatment of diabetic patients may be necessary.

Limitation of the study

Our study conducted on a limited sample size of patients with resistant hyperlipidemia. This study needs further investigation on larger samples and also on all patients with hyperlipidemia without anti-lipid treatment.

REFERENCES

- Pickup JC, Chusney GD, Crook MA, Viberti GC. Hypomagnesaemia in IDDM patients with microalbuminuria and clinical proteinuria. *Diabetologia*. 1994;37:639.
- Abbate SL, Brunzell JD. Pathophysiology of hyperlipidemia in diabetes mellitus. *J Cardiovasc Pharmacol*. 1990;16 Suppl 9:S1-7.
- Alberti KG, Zimmet P, Shaw J. Metabolic syndrome--a new world-wide definition. A Consensus Statement from the International Diabetes Federation. *Diabet Med*. 2006; 23:469-80.
- Ames RP. Hyperlipidemia in hypertension: causes and prevention. *Am Heart J*. 1991;122:119-24.
- Arpacı D, Tocoglu AG, Ergene H, Korkmaz S, Ucar A, Tamer A. Associations of serum Magnesium levels with diabetes mellitus and diabetic complications. *Hippokratia*. 2015;19:153-7.
- Arslanoğlu I, Günöz H, Bundak R, Saka N. Hypomagnesaemia in childhood IDDM and risk of nephropathy. *Diabetologia*. 1995;38:629.
- Conn VS, Ruppert TM, Chase JD. Blood pressure outcomes of medication adherence interventions: systematic review and meta-analysis. *J Behav Med*. 2016; 39:1065-75.
- Corica F, Allegra A, Ientile R, Buemi M, Corsonello A, Bonanzinga S, et al. Changes in plasma, erythrocyte, and platelet magnesium levels in normotensive and hypertensive obese subjects during oral glucose tolerance test. *Am J Hypertens*. 1999;12:128-36.
- Dasgupta A, Sarma D, Saikia UK. Hypomagnesaemia in type 2 diabetes mellitus. *Indian J Endocrinol Metab*. 2012; 16:1000-3.
- Deepti R, Nalini G. Magnesium plays a salient role in the cells. *J Clin Biomed Sci*. 2014;4:341-5.
- Guerrero-Romero F, Rodríguez-Morán M. Low serum magnesium levels and metabolic syndrome. *Acta Diabetol*. 2002;39:209-13.
- Joffres MR, Reed DM, Yano K. Relationship of magnesium intake and other dietary factors to blood pressure: the Honolulu heart study. *Am J Clin Nutr*. 1987;45:469-75.
- Kawano Y, Matsuoka H, Takishita S, Omac T. Effects of magnesium supplementation in hypertensive patients: assessment by office, home, and ambulatory blood pressures. *Hypertension*. 1998;32:260-5.
- Kim DJ, Xun P, Liu K, Loria C, Yokota K, Jacobs DR Jr, He K. Magnesium intake in relation to systemic inflammation, insulin resistance, and the incidence of diabetes. *Diabetes Care*. 2010;33:2604-10.
- Longstreet DA, Heath DL, Panaretto KS, Vink R. Correlations suggest low magnesium may lead to higher rates of type 2 diabetes in Indigenous Australians. *Rural Remote Health*. 2007; 7:843.
- Mahalle N, Kulkarni MV, Naik SS. Is hypomagnesaemia a coronary risk factor among Indians with coronary artery disease? *J Cardiovasc Dis Res*. 2012;3:280-6.
- Mather HM, Levin GE. Magnesium status in diabetes. *Lancet*. 1979;1(8122):924.
- Randall EW, Mathews M, Gadag V, Zhang H, Sun G. Relationship between serum magnesium values, lipids and anthropometric risk factors. *Atherosclerosis*. 2008; 196:413-9.
- Reverter JL, Senti M, Rubiés-Prat J, Lucas A, Salinas I, Pizarro E, et al. Relationship

- between lipoprotein profile and urinary albumin excretion in type II diabetic patients with stable metabolic control. *Diabetes Care*. 1994;17:189-94.
20. Sales CH, Pedrosa LF, Lima JG, Lemos TM, Colli C. Influence of magnesium status and magnesium intake on the blood glucose control in patients with type 2 diabetes. *Clin Nutr*. 2011;30:359-64.
21. Wasada T, Katsumori K, Saeki A, Saito S, Omori Y. Urinary albumin excretion rate is related to insulin resistance in normotensive subjects with impaired glucose tolerance. *Diabetes Res Clin Pract*. 1997; 34:157-62.
22. Whelton PK, Klag MJ. Magnesium and blood pressure: review of the epidemiologic and clinical trial experience. *Am J Cardiol*. 1989;63:26G-30G.