



STUDY OF INCIDENCE OF DEPRESSION IN OLDER ADULTS PATIENTS VISITING TO TERTIARY CARE CENTER IN BIHAR

General Medicine

Dr. Manish Kumar Singh

Senior Resident, Department of General Medicine AIIMS Patna.

Dr. Shiv Kumar

Senior Resident, Department of General Medicine AIIMS Patna.

Dr. Vijay Kumar*

Associate Prof. Department of General Medicine AIIMS Patna. *Corresponding Author

ABSTRACT

Background: Depression is the commonest psychiatric disorder in the elderly. We attempted to analyze the prevalence and correlates of depression in the elderly. Depression prevalence in the geriatric population has been reported to be between 8-20% at the community level and up to 37% in those receiving primary level care.

Methods: An observational study was carried out taking cases from patients attending the geriatric clinic for the first time. Depression was diagnosed by the Geriatric Depression Score short form (≥ 5). Various epidemiological parameters were assessed in 504 subjects (M = 304, F = 200; mean age = 66.47 ± 13.71 years). Patients less than 60 years of age, those suffering from some acute illness or requiring hospitalization and those unable to participate in the study because of other physical limitations such as speech or hearing impairment or severe dementia were excluded from the study.

Conclusion: The prevalence of depression in the geriatric population in our study was moderately high and emphasizes the importance of specialized counseling and pharmacotherapeutic approaches in older patients presenting to the tertiary hospital set-up.

KEYWORDS

Depression, Epidemiology, Geriatric Depression Score, Prevalence.

INTRODUCTION

Depression is the commonest psychiatric disorder in the elderly. We attempted to analyze the prevalence and correlates of depression in the elderly. The geriatric population of the world has quadrupled from 202 million in 1950 to 843 million in 2013. This is projected to increase three fold by 2050. The brunt of this old age population explosion has been borne by relatively less developed countries quite a few of which lie in south and south-east Asia. By 2050, these countries would account for nearly 80% of the world's geriatric population. Depression is the commonest psychiatric disorder in the aged.

Unipolar depressive disorders contribute maximally to the burden of disease in both middle and high income countries and also are among the ten most important causes in low income nations. The burden of disease is a composite measure of life years lost due to premature death and a weighted measure of years spent with disability, pain or impairments.³

The DSM-IV defines major depressive disorder as depressed mood or a loss of interest or pleasure in daily activities for more than two weeks.

⁴ Depression is an especially significant problem in the aged on account of multiple factors such as physical disability, retirement from active life and changing social dynamics. The persistently declining old age support ratio is cause of concern as it means a smaller support group to deal with the varied needs of the depressed old adults. ¹ The ill-effects of depression range from an increase in the rates of heart disease and stroke to suicidal tendencies in patients.

Depression prevalence in the geriatric population has been reported to be between 8-20% at the community level and up to 37% in those receiving primary level care. ⁷ A retrospective analysis of 74 original studies on a combined strength of 487,275 subjects 60 years of age or more determined the median prevalence of depressive disorders in older adults to be 10.3 % (IQR 4.7%-16.0%). The median prevalence of depression among the Indian geriatric population was estimated to be 21.9% (IQR 11.6%- 31.1%). ⁸ On the other hand, community based mental health studies conducted in India have reported depression prevalence in the aged varying widely from just above 10% up to as high as more than 50%.

Almost no data exists on depression prevalence and its correlates in the north Indian population presenting to tertiary centers. Considering the geriatric population at the point of higher medical care has an added advantage while appreciating the fact that the prevalence of depression in such a population would be more. Primary health care systems in north India suffer from unawareness among physicians about geriatric specific problems, the prevalence of untrained quacks imparting

healthcare and many others. On the other hand, the population presenting to tertiary centers is amenable to intervention and compliance of patients with treatment is appreciable and hence it is this group which might be a prime target of therapy. The authors present an observational study on 504 older patients presenting at a tertiary care center in northern India. The study tries to gauge the prevalence of depression and other epidemiological attributes of depression in older adults visiting advanced care hospitals. The study tries to analyze the association of depression with gender, socio- economic status, occupation, marital status, place of residence, use of habit forming substances and insomnia. The study findings may lay the foundation for multi-center hospital based studies which would evaluate the causal role of various modifiable factors in depression and mark potential intervention sites.

MATERIAL AND METHODS

A cross-sectional study was carried out at a tertiary care teaching and research hospital of Bihar. Cases were taken from patients attending the geriatric clinic for the first time. Patients less than 60 years of age, those suffering from some acute illness or requiring hospitalization and those unable to participate in the study because of other physical limitations such as speech or hearing impairment or severe dementia were excluded from the study. The total number of newly registered geriatric outpatients during the period of study amounting to 1084 patients was assessed for eligibility. 280 patients were excluded as they did not fulfill the inclusion criteria or had one or more of the exclusion points: 30 were between 55-60 years age, 221 needed hospitalization or were suffering from some acute illness and 29 had severe dementia. Out of the rest, 300 did not give consent. The response rate of the study was thus 62.7%.

The study proper included 304 male and 200 female patients. General information was collected including name, age, gender, socio-economic status, place of living, educational status, occupation, marital status, insomnia and habit forming substance use, using a pre-decided standard questionnaire. The questionnaire was filled by resident doctor posted in the Geriatric unit on the basis of verbal interview of the patient in patient's local dialect. All patients were interviewed by the same interviewer over the period of study. Patients were segregated into various socio-economic groups based on the modified Prasad classification proposed for the year 2004 (Table 1). Furthermore, categories I and II, III and IV, V and VI in the classification scheme were combined to form high, middle and low socio- economic groups. Habit forming substance use was defined as the almost daily consumption of tobacco by smoking or chewing irrespective of the quantity or the almost daily consumption of alcohol irrespective of the quantity.

Table 1: Proposed Indian Social Classification For 2004 [13, 14]

Social class	Per capita monthly income limits (Rupees: 1 Indian Rupee = 0.016 \$) Prasad (1970) Suggested modification (2004)		
I. Upper high	100 and above	10000 and above	HIGH
II. High	50-99	5000-9999	
III. Upper middle	30-49	3000-4999	MIDDLE
IV. Lower middle	15-29	1500-2999	
V. Poor	15	500-1499	LOW
VI. Very poor or Below poverty line (BPL)	-	Below 500	

Diagnosis Of Depression:

Depression was diagnosed by the Geriatric depression score (GDS short form, 15 questions).¹⁵ This scale was administered to the participants of the study in translated form in Hindi. A score of greater than or equal to 5 was diagnostic of depression. The validity of the GDS in depression diagnosis in the aged compared to standard diagnostic tools such as the ICD-10 and DSM-IV has been established previously.

Sample Size Determination:

Considering the widely variable prevalence values for depression in the Indian aged, a mean prevalence of 25% was assumed. With a 5% allowable error, the sample size required for our study was determined to be 300.²⁰

RESULTS:

504 patients (M=304, F=200) were analyzed. The mean age of the subjects was 66.47 ± 13.71 years. Majority of the patients (40.8%) were in 60-64 years age group. Depression prevalence was 45.0% (M: 40.5%, F: 52%). (Table 2)

Table 2: Age And Sex Distribution Of Geriatric Subjects With Gds Scores

GDS AGE GROUP	<5		5-9		≥10	
	Male	Female	Male	Female	Male	Female
60-64	76 (15)	47 (9)	20 (4)	21 (4)	19 (4)	23 (4)
65-69	43 (8)	27 (5)	14 (3)	19 (4)	19 (4)	14 (3)
70-74	34 (7)	12 (2)	15 (3)	11 (2)	10 (2)	6 (1)
≥75	28 (5)	10 (2)	10 (2)	6 (1)	16 (3)	4 (1)

These variables were further used to develop a model of depression using binomial logistic regression analysis. Table 4 shows the analysis results for significant variables. The variables which were significant as independent predictors of depression were marital status and insomnia. The model was statistically significant ($\chi^2=59.042$, $p<0.001$). The model explained 14.8% of the variance in depression and correctly classified 65.9% of the cases.

Table 3: Multiple Logistic Regression Analysis Showing Independent Predictors Of Depression In The Subjects:

VARIABLES	DEPRESSED n (%)	ADJUST ED OR	95% CI	p-value
MARITAL STATUS				
MARRIED	174(41.2)	1	0.121-	<0.01
UNMARRIED	1(50)	2.050	34.603	
WIDOW	52(65)	2.293	1.349-3.896	
INSOMNIA				
NO	130(37.9)	1		<0.001
YES	97(60.2)	2.512	1.668-3.784	

In addition, the commonest systemic co-morbidity in the depressed patient population was cardiovascular disease including coronary artery disease and hypertension (n=56; 24.67%). The further analysis of systemic co-morbidities in depressed patient population could not be done as study design did not permit.

DISCUSSION:

The present study evaluated the various attributes of depression in the older adult population presenting to the tertiary care hospital level. In developing countries this is often the first level of dedicated geriatric care where a detailed neuropsychological assessment of the patient is performed.

The study involved the use of the GDS for diagnosis of depression. With a cut-off point of ≥ 5 , the GDS has a high sensitivity of 92% and specificity of 81% in diagnosing geriatric depression.²¹ This form of diagnosis was preferred owing to the atypical presentation of depression in the aged which makes the use of gold standard diagnostic tools such as the DSM-IV and ICD-10 difficult. A similar approach has been adopted by Stek and others in an earlier study.²¹ Yokoyama and others in a study of the association of depression and insomnia in Japan made use of the CES-D scale short form for depression diagnosis. Hence, the use of a screening instrument for diagnosis is a valid concept.²² The study sample had more males than females which fits the usual pattern of health care seeking in this predominantly rural belt with a patriarchal society.

The depression prevalence in our study was 45.03%. As has been mentioned earlier, multiple studies from the Indian sub-continent have reported highly variable figures of geriatric depression. A depression prevalence similar to our study was quoted in the AGED study. 46.2% of older nursing home patients were found to be suffering from depression.²³ A study by Rajkumar and others involving 1000 participants aged over 65 years from Kaniyambadi block, Vellore in southern India brought up a depression prevalence of 12.7%. The WHO Study on global ageing and adult health (SAGE) including adults from China, Ghana, India, Mexico, Russia, and South Africa report even lower depression prevalences of only around 6% in males and up to 8% in females. Although these figures are grossly low as compared to the data of the current study, the studies are essentially different in construction. The authors propose that the prevalence of depression in the aged population visiting hospitals is more significant from a therapeutic view point especially in resource starved developing nations where health-care fails to reach the bulk of the population.

Depression is a major concern in older adults and its development depends on multiple social and life-style parameters as well as the existence of systemic co-morbidities. This study mainly analyzed the former. The variables that showed significant p-values on univariate analysis were gender, occupation, educational level, marital status, insomnia and use of habit forming substances. In contrast, Rajkumar et al's work from south India earmarked low income, hunger, cardiac illnesses, transient ischemic attack, past head injury and diabetes as risk factors for geriatric depression. The variables which were significant on multivariate analysis (marital status, insomnia) and those that were expected to affect depression profile based on earlier literature (age, gender, education, use of habit forming substances) are individually discussed. Whereas income is a significant determinant of depression in the WHO study on global ageing, the univariate analysis of depression with respect to socio-economic status in the current work did not find similar results.

Age:

In this study, variation in depression prevalence with age was not evident. This implies that in the older population, depression rates do not vary in specific age sub-groups. The observation may be confounded by the fact that in the older age groups, fewer patients come for treatment to a tertiary center as evident at our center. Hence, depression cases in the very old may be missed. Further, patients with severe dementia were excluded from our study and a higher number of such patients in the older age groups may lead to bias. These explain the apparent conflict of our results with findings published elsewhere where the oldest old have been shown to be at high risk of chronic depression.²¹ None of the studies conducted in India have addressed this unique problem of varying depression rates among different subgroups of the elderly population and it needs to be addressed separately in cross-sectional and cause-effect designs.

Gender:

52% of the female patients were depressed compared to 40.5% of the male patients. Female gender was a significant risk factor for depression on univariate analysis. Sherina reported similar results. Relative lack of care for the female gender in patriarchal societies such as in northern India may be a contributor. The fact that gender was not a significant determinant on multi-variate analysis may stem from the fact that the gender differences are affected by various other socio-economic differentials such as education, relative accessibility of resources.

Although age and gender are non-modifiable factors in depression

prevalence, much of the female gender specific rise in depression is due to socio-economic factors such as differential access to healthcare, social rights and finances. This is an area where administrative and social institutions can intervenemore than the healthcare setup to reduce this added burden in females.

The use of tobacco (chewable/smokable) and alcohol showed positive association with depression on univariate analysis. Jain et al reported a 58.13% prevalence of depression among older individuals inclined towards substance abuse. Recent research has highlighted the epidemic status of substance abuse among older populations. The present study did not use specific diagnostic criteria for substance abuse problems and only considered whether habit forming substances were being used by the subjects. Different levels of use may hence have been clubbed together. This stays a short-coming of the study. Also prescription drug abuse data was not collected. In the Indian subcontinent, chewable tobacco use is very common especially among rural people in the northern part of the country. In some social strata, it is perfectly acceptable for older females to smoke tobacco in the form of 'beedis', often as a remedy for constipation. Alcohol use is less common but still many older individuals carry over this habit from their young adult life. Habit forming substance use is associated with a feeling of loneliness and lower life satisfaction.

The complications of habit forming substance use such as COPD, malignancies and coronary artery disease set in by the time a person reaches the 60+ age group. These may be additives in the pathogenesis of depression. On the other hand, depressed older people may more often resort to intoxicating substance use. However, substance abuse data was recorded using crude criteria and any interpretation with regards to these should be guarded.

Insomnia:

Insomnia was seen to have a positive correlation with depression. The odds of a patient with insomnia having depression was 2.5 times that of one with unimpaired sleep ($p=0.000$). Yokoyama et al reported an association between difficulty initiating sleep and depression in community dwelling older individuals in Japan.²² Sanjay and others found the odds of depression to be nearly 16 times higher in elderly with insomnia. Many studies have proven that insomnia is a risk factor for new onset major depressive disorder. Persistent middle insomnia has been shown to trigger new onset depression. Research has also emphasized that insomnia may perpetuate depression. Alternatively insomnia has been long regarded as a symptom rather than a cause of depression. Since, this study was not of a longitudinal design, cause-effect relationship could not be ascertained but patients with insomnia were identified as a high risk group for co-existent depression.

CONCLUSION:

The present study revealed spousal loss and insomnia as red flag signs in the older adults for a more detailed evaluation of depression. Several other variables such as female gender, education up to the middle school level and use of habit forming substances were found to have minor effect on depression occurrence in the aged on univariate analysis. The prevalence of depression in the geriatric population in our study was moderately high and emphasizes the importance of specialized counseling and pharmacotherapeutic approaches in older patients presenting to the tertiary hospital set-up.

REFERENCES:

- United Nations, Department of Economic and Social Affairs, Population Division. World Population Ageing 2013. ST/ESA/SER.A/348.
- Volkert J, Schulz H, Härter M, Włodarczyk O, Andreas S. The prevalence of mental disorders in older people in Western countries—a meta- analysis. *Ageing Res Rev.* 2013;12(1):339–353.
- World Health Organization. The global burden of disease: 2004 update. Geneva: WHO; 2008.
- American Psychiatric Association. Diagnostic and statistical manual of mental disorders. 4. Washington, DC: Author; 2000. text revision.
- World Health Organisation. Conquering Depression. New Delhi: Regional Office for South-East Asia. 2001 http://whqlibdoc.who.int /searo/2001/ SEA_Ment_120.pdf. [Accessed Mar 10, 2013].
- Jacob KS. Depression: A major public health problem in need of a multi-sectoral response. *Indian J Med Res.* 2012;136(4):537–9.
- World Health Organization. The World Health Report: 2001: mental health: new understanding new hope. Geneva, Switzerland: WHO; 2001.
- Barua A, Ghosh MK, Kar N, Basilio MA. Prevalence of depressive disorders in the elderly. *Ann Saudi Med.* 2011;31(6):620–4.
- Teja JS, Narang RL. Pattern of incidence of Depression in India. *Indian J Psych.* 1970;12:33–9.
- Bagadia VN, Jeste DV, Doshi SU, Shah LP. Depression: A study of demographic factors in 233 cases. *Indian J Psych.* 1973;15:209–16.
- Raju SS. Frequency of depressive disorders in psychiatric clinics in India: A comparative analysis. *Indian J Psych.* 1979;21:176–9.
- Ponnudurai R, Somasundaram O, Balakrishnan S, Srinivasan N. Depression-a study of 80 cases. *Indian J Psych.* 1981;23:256–8.
- Prasad BG. Changes proposed in Social classification of Indian families. *J Indian Med Assoc.* 1970;55: 198–9.
- Agarwal AK. Social Classification: The Need to Update in the Present Scenario. *Indian J Community Med.* 2008;33(1): 50–51.
- Sheikh JI, Yesavage JA. Geriatric Depression Scale (GDS): Recent evidence and development of a shorter version. *Clin Gerontol.* 1986;5(1-2): 165–173.
- Prakash O, Gupta LN, Singh VB, Nagarajara G. Applicability of 15-item Geriatric Depression Scale to detect depression in elderly medical outpatients. *Asian J Psychiatr.* 2009;2(2):63–5.
- Jongenelis K, Pot AM, Eisses AM, Gerritsen DL, Derksen M, Beekman AT, Kluiters H, Ribbe MW. Diagnostic accuracy of the original 30-item and shortened versions of the Geriatric Depression Scale in nursing home patients. *Int J Geriatr Psychiatry.* 2005;20(11):1067–74.
- Ganguli M, Dube S, Johnston JM, Pandav R, Chandra V, Dodge HH. Depressive symptoms, cognitive impairment and functional impairment in a rural elderly population in India: a Hindi version of the geriatric depression scale (GDS-H). *Int J Geriatr Psychiatry.* 1999;14(10):807–20.
- Almeida OP, Almeida SA. Short versions of the geriatric depression scale: a study of their validity for the diagnosis of a major depressive episode according to ICD-10 and DSM-IV. *Int J Geriatr Psychiatry.* 1999;14(10):858–65.
- Arya R, Antonisamy B, Kumar S. Sample size estimation in prevalence studies. *Indian J Pediatr.* 2012;79(11):1482–1488.
- Stek ML, Vinkers DJ, Gussekloo J, van der Mast RC, Beekman AT, Westendorp RG. Natural history of depression in the oldest old: population-based prospective study. *Br J Psychiatry.* 2006;188:65–9.
- Yokoyama E, Kaneita Y, Saito Y, Uchiyama M, Matsuzaki Y, Tamaki T, Munezawa T, Ohida T. Association between depression and insomnia subtypes: a longitudinal study on the elderly in Japan. *Sleep.* 2010;33(12):1693–702.
- Jongenelis K, Pot AM, Eisses AMH, Beekman ATF, Kluiters H, Ribbe MW. Prevalence and risk indicators of depression in elderly nursing home patients: the AGED study. *J Affect Disord.* 2004;83(2-3):135–142.