



## ELECTROLYTE CHANGES IN NEONATES BEFORE AND AFTER PHOTOTHERAPY FOR NEONATAL HYPERBILIRUBINEMIA: A PROSPECTIVE HOSPITAL-BASED OBSERVATIONAL STUDY

### Neonatology

**Arvind Kumar Shukla** MD, Assistant Professor, Department of Paediatrics, Maharaja Suheldev Autonomous State Medical College, Bahraich, India

**Manjari Tiwari** MD, Senior Resident, Department of Paediatrics, Maharaja Suheldev Autonomous State Medical College, Bahraich, India

**Nidhika Pandey** MD, Assistant Professor, Department of Paediatrics, Rama Medical College Hospital & Research Centre, Kanpur, India

### ABSTRACT

**Background:** Neonatal hyperbilirubinemia is one of the most common cause of hospital admission in neonatal intensive care unit and phototherapy remains the mainstay for treatment of neonatal hyperbilirubinemia. The present study aims to evaluate the electrolytes changes in term and preterm neonates undergoing phototherapy. **Materials and Methods:** In present study, a total of 45 neonates were involved. S. Sodium, S. Potassium, S. Calcium, S. Magnesium, S. Phosphorus levels were sent before starting of phototherapy. Phototherapy was given to neonates and when the serum bilirubin levels came 2–3 mg/dl below age specific levels on the normogram, above blood parameters were sent again. **Results:** Our study revealed that there was a significant ( $P < 0.05$ ) decline in serum electrolyte levels after phototherapy. However, all the neonates remained asymptomatic and did not require any intervention. Furthermore, electrolyte changes were seen more in preterm neonates as compared to term neonates. **Summary:** There is significant change in electrolytes in neonates receiving phototherapy. Low birth weight and preterm neonates are at higher risk of developing electrolyte imbalances and they require close monitoring.

### KEYWORDS

Electrolytes, neonatal hyperbilirubinemia, neonates

### INTRODUCTION

Neonatal hyperbilirubinemia is a common physical finding in 1st week of life.[1] It is a common cause of concern to parents and the most common cause of hospital readmission during neonatal period.<sup>1</sup> The clinical symptoms of jaundice is yellowish discoloration of skin due to deposition of unconjugated bilirubin. Clinically jaundice appears in the neonates when bilirubin level goes  $>7$  mg/dl.<sup>2</sup> 60% term and 80% preterm neonates develop jaundice during the 1st week of life, among them only 6.1% of term neonates are found to have a S. Bilirubin  $>12.9$  mg/dL and 3% have a S. Bilirubin  $>15$  mg/dl.<sup>3</sup>

Neonatal hyperbilirubinemia is treated by phototherapy, exchange transfusion or pharmacological agents. Among these three, phototherapy is the mainstay of treatment of neonatal hyperbilirubinemia and it also decreases the incidence of exchange transfusion and bilirubin encephalopathy. Phototherapy has been effectively used as comparatively non-invasive and inexpensive method for treatment of neonatal hyperbilirubinemia.<sup>4</sup>

Today exchange transfusions are very rare and are used only as a rescue therapy to avoid kernicterus in newborns with severe hyperbilirubinemia when phototherapy is not effective. As any treatment has its adverse effects, phototherapy also has some side effects but they are not harmful or severe.<sup>5</sup>

The main side effects are hyperthermia, feed intolerance, loose stools, skin rashes, bronze baby syndrome, retinal changes, dehydration, hypocalcemia, redistribution of blood flow, genotoxicity and electrolyte imbalances.<sup>2</sup>

Very few studies are currently available that describes adverse effects of phototherapy on electrolytes. Most of the studies showed that phototherapy causes hypocalcemia, which is more in preterm (90%) than term (75%).<sup>6</sup> It is known that phototherapy converts bilirubin into water soluble isomers that are easily eliminated through the gastrointestinal tract or lost in urine.<sup>7</sup> This is because bilirubin absorbs light most strongly in blue region of spectrum near 460 nm wavelength. Gallidium nitride light emitting diodes, which delivers high intensity light of narrow wavelength spectrum, have been developed and are increasingly being used to treat neonatal jaundice.<sup>7</sup>

The aim of phototherapy is to avoid increase in previously increased total serum bilirubin levels, to thwart progression to encephalopathy or kernicterus and to stop the total bilirubin from getting higher to a level that would require exchange transfusion. Since there are only few studies elaborating on the effect of phototherapy on serum electrolytes published till now, more studies are required to reach a conclusion

about these effects on the high-risk neonates.

### MATERIALS AND METHODS:

A prospective hospital-based observational comparative study was conducted on 45 eligible neonates, who were admitted in neonatal intensive care unit (NICU), Medical College situated in north India from April 1, 2021 to March 31, 2022. Informed signed consent was taken from parents and the study was approved by Ethical Committee. Our study included both preterm and term neonates who had unconjugated hyperbilirubinemia and required phototherapy. Neonates with positive septic screen were excluded from the study.

The eligible neonates were put under double surface phototherapy with a wavelength of 425–475 nm with a distance of 25–30 cm from skin. Before starting of phototherapy venous samples were sent for serum electrolytes ( $\text{Na}^+$ ,  $\text{K}^+$ ,  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ ,  $\text{PO}_4^-$ ). Repeat blood samples were sent for serum electrolytes at 12 h of phototherapy. Based on response to phototherapy, samples were taken 12 hourly till serum bilirubin level come 2–3 mg/dl below the age-specific levels on the normogram.  $\text{Na}^+$  and  $\text{K}^+$  were determined by Easylite Analyzer.  $\text{Mg}^{2+}$  was determined by Calmagite method using ERBA CHEM V semi autoanalyzer. Phosphorus was determined using ultraviolet Molybdate method, using ERBA 360 auto analyzer. Calcium was determined using Arsenzo method, using ERBA 360 auto analyzer.

Comparative study was done between the before phototherapy and after phototherapy samples. The data were analyzed with Statistical Package for Social Sciences Version 23.0. Co-relation between various parameters was done using Pearson coefficient,  $P < 0.05$  was considered significant.

### RESULTS

In our study, out of 45 neonates, 25 (55.60%) were male and 20 (44.40%) were female [Table 1]. Out of 45 neonates, 17 (37.8%) were low-birth-weight and 28 (62.2%) had normal birth weight [Table 2]. Mean birth weight of neonates was  $2.66 \pm 0.53$  kg. Out of 45 neonates, 16 (35.6%) were preterm neonates ( $<0.001$  [Table 3]).

In our study, we found that 7 (15.4%) newborns, out of 45 newborns developed hypocalcaemia after phototherapy which was found to be statistically significant difference with  $P < 0.001$  [Table 4]. In our study, only 3 (6.7%) neonates, out of 45 neonates developed hypokalemia after phototherapy which was statistically significant with  $P < 0.001$  [Table 4].

### DISCUSSION

This study was a hospital based prospective study of neonates admitted

in NICU with neonatal hyperbilirubinemia. The study was aimed at evaluating the electrolyte changes in preterm and term neonates undergoing phototherapy. In our study, we estimated serum electrolytes of 45 neonates, who had neonatal hyperbilirubinemia and were receiving phototherapy. We compared the electrolyte changes before and after phototherapy and co-related the electrolyte changes with gestational age and birth weight.

There are very less studies which aim at understanding the electrolyte changes in neonates undergoing phototherapy, especially on serum sodium, calcium, magnesium, potassium and phosphorus levels and comparing the effect of phototherapy on electrolytes in preterm and term. Through this study, we endeavoured to look into electrolyte imbalances in neonates receiving phototherapy.

In our study, we studied 45 neonates, out of which 25 (55.60%) were males and 20 (40.40%) were females. In this study, out of 45 neonates, 17 (37.8%) neonates had low-birth-weight and 28 (62.2%) neonates had normal birth weight. The mean birth weight of neonates was 2.66 ± 0.53 kg.

In our study we included 16 (35.6%) preterm neonates and 29 (64.4%) term neonates. Karamifar et al.<sup>8</sup> and Sethi et al.<sup>9</sup> selected preterms whereas Taheri et al.<sup>7</sup> and Eghbalian et al.<sup>10</sup> included only term neonates in their study.

In our study, there was decline in serum sodium levels after phototherapy which was statistically significant i.e., P < 0.001. The mean serum sodium before and after phototherapy was 139.94 ± 1.55 and 138.69 ± 1.96, which was similar to Reddy et al.<sup>6</sup> (139.02 ± 3.12 and 138.16 ± 3.36), Kumar et al.<sup>11</sup> (139.01 ± 3.119 and 138.15 ± 3.35) and Gayatri et al.<sup>1</sup> (141.97 ± 2.795 and 140.66 ± 4.751).

In our study, we found out that out of 45 neonates, 6 (13.3%) neonates developed hyponatremia after phototherapy [Table 5]. Although the neonates remained asymptomatic and no intervention was required for hyponatremia. In our study, we found that mean sodium levels before and after phototherapy in low-birth-weight neonate were 138.52 ± 1.46 and 135.13 ± 1.68 respectively and this difference was significant with a P < 0.001.

Our study found that incidence of hyponatremia in neonates undergoing phototherapy was high in low-birth-weight neonate (17.6%) as compared to normal weight neonates (10.7%) [Table 6]. These results are similar to those of Reddy et al.<sup>6</sup> (17.2% low-birth-weight and 2.6% normal birth weight), Kumar et al.<sup>11</sup> (17.2% low-birth-weight and 2.6% normal birth weight neonates) and Gayatri et al.<sup>1</sup> (18.75% low-birth-weight and 10.32% normal birth weight).

This implies that low-birth-weight neonates are at a higher risk of developing hyponatremia as compared to normal birth weight neonates and continuous monitoring of sodium levels is required. Furthermore, we found that 18.8% preterm neonates developed hyponatremia after phototherapy as compared to term neonates (10.3%) [Table 7]. This is similar to other studies i.e., Reddy et al.<sup>6</sup> (17.6% preterm and 3.1% term neonates developed hyponatremia), Kumar et al.<sup>11</sup> (17.5% preterm and 3.1% term neonates developed hyponatremia post phototherapy), Gayatri et al.<sup>1</sup> (18.3% preterm and 11.02% term neonates).

Hence, with above results and with concordance with other studies, it is implied that preterm neonates are at a higher risk of developing hyponatremia, who are undergoing phototherapy as compared to term neonates undergoing phototherapy. Although neonates developed hyponatremia following phototherapy, but none of the neonates developed any clinical symptoms and remained asymptomatic.

In our study, we found that following phototherapy, there was significant decline in the mean serum calcium levels which was statistically significant (P < 0.001). The mean S. calcium levels before phototherapy was 8.98 ± 0.37 and after phototherapy was 8.25 ± 0.64. In our study, we co-related hypocalcemia following phototherapy with gestational age and found that preterm neonates are at higher risk of developing hypocalcemia as compared to term neonates. 18.8% of preterm neonates developed hypocalcemia as compared to 10.3% of term neonates [Table 7]. These results were similar to studies conducted by Karamifar et al.<sup>8</sup> (22.6% preterm and 8.7% term neonates), Reddy et al.<sup>6</sup> (41.2% preterm and 6.2% term neonates), Gayatri et al.<sup>1</sup> (25.34% preterm and 10.24% term neonates) but

differed from Arora et al.<sup>3</sup> where hypocalcemia was seen more in term neonates in comparison to preterm neonates

In our study, co-relation of hypocalcemia following phototherapy was done with birth weight and it was found that low-birthweight neonates (31.3%) are at a higher risk of developing hypocalcemia as compared to normal birth weight neonates (27.6%). This was similar to studies conducted by Reddy et al.<sup>6</sup> (36.2% in low-birth-weight and 6.2% in normal birth weight) and Gayatri et al.<sup>1</sup> (26.25% in low-birth-weight and 7.94% in normal birth weight).

The neonates who developed hypocalcemia following phototherapy did not show any features clinically and remained asymptomatic [Table 8]. In our study, we found that following phototherapy there was significant decline in serum potassium levels which was statistically significant with P < 0.001. The mean serum potassium levels before phototherapy were 4.81 ± 0.34 and after phototherapy were 4.20 ± 0.43.

This was similar to study by Suneja et al.<sup>12</sup> where mean potassium levels before phototherapy were 6.095 ± 1.4 and after phototherapy were 5.28 ± 1.08 and Gayatri et al.<sup>1</sup> where mean potassium levels before phototherapy were 5.01 ± 0.683 and after phototherapy were 4.63 ± 0.666. In Reddy et al.<sup>6</sup> there was no decline in serum potassium levels, before and after phototherapy and our results were not in consonance with their study.

Hence, actual relationship between serum potassium with phototherapy needs to be evaluated further with large sample size. In our study, total of 3 (6.7%) neonates developed hypokalaemia, out of which 2 (12.5%) were preterm and 1 (3.4%) was term neonate [Table 9]. Study by Reddy et al.<sup>6</sup> is in consonance with our study where only 0.4% neonates developed hypokalaemia following phototherapy. In our study, we found that there was a significant decline in mean serum magnesium and phosphorus levels, but there was no co-relation of decline of these electrolytes with gestational age and birth weight.

**CONCLUSION**

In our study, there was statistically significant decline in mean serum electrolyte after phototherapy. The incidence of hypocalcaemia following phototherapy was 15.4% and it was higher in preterm (25.1%) and low-birth-weight neonates (29.4%) as compared to term (10.3%) and normal birth weight neonates (7.1%). The incidence of hyponatremia after phototherapy was 13.3% and it was higher in preterm (18.8%) and low-birth-weight neonates (17.6%) as compared to term neonates (10.3%) and normal birth neonates (10.7%). The incidence of hypokalaemia following phototherapy was 6.7% and it was higher in preterm neonates (12.5%) and low birth neonates (11.8%) as compared to term neonates (3.4%) and normal birth weight neonates (3.6%). It was also observed that, in most of the neonates, following phototherapy there was significant decrease in serum sodium, potassium and calcium levels, but all the neonates remained asymptomatic and no intervention was required. Hence, all neonates require continuous monitoring while undergoing phototherapy and further studies are required to know about the effect of phototherapy.

**Table 1: Gender distribution of neonates**

Gender	Number	Percentage
Male	25	55.60
Female	20	44.40
Total	45	100

**Table 2: Birth weight (in kg) distribution of neonates**

Birth weight (in kg)	Number	Percentage
LBW (<2.5 kg)	17	37.8
Normal (2.5-4.0 kg)	28	62.2
Total	45	100

LBW: Low birth weight

**Table 3: Gestational age (in weeks) distribution of neonates**

Gestation age (in weeks)	Number	Percentage
< 37 weeks	16	35.6
37-42 weeks	29	64.4
Total	45	100

**Table 4: Comparison of mean value of different study variables before and after phototherapy with birth weight**

Electrolytes	Before phototherapy	After phototherapy	Mean difference	P-value
Sodium	139.94 ± 1.55	138.69 ± 1.96	-1.21 ± 1.41	< 0.001
Calcium	8.98 ± 0.37	8.25 ± 0.64	-0.73 ± 0.27	< 0.001
Potassium	4.81 ± 0.34	4.20 ± 0.43	-0.61 ± 0.09	< 0.001
Magnesium	0.75 ± 0.05	0.72 ± 0.06	-0.03 ± 0.01	0.12
Phosphorus	1.55 ± 0.12	1.52 ± 0.13	-0.03 ± 0.01	0.15

	(Mean±SD)	(Mean±SD)		
LBW (< 2.5 kg)				
S. Sodium	138.52±1.46	135.13±1.68	2.390	<0.001*
S. Potassium	4.64±0.28	4.32±0.38	0.320	<0.001*
S. Calcium	8.14±0.22	7.18±0.32	0.960	<0.001*
S. Magnesium	2.08±0.10	1.96±0.11	0.180	<0.001*
S. Phosphorus	3.68±0.12	3.30±0.10	0.380	<0.001*
Normal Birth weight (>2.5 kg)				
S. Sodium	140.94±1.52	140.19±1.84	0.750	<0.001*
S. Potassium	4.97±0.32	4.49±0.48	0.480	<0.001*
S. Calcium	9.01±0.39	8.46±0.62	0.680	<0.001*
S. Magnesium	2.23±0.13	2.01±0.14	0.220	<0.001*
S. Phosphorus	3.92±0.15	3.68±0.16	0.240	<0.001*

LBW: Low birth weight

**Table 5: Frequency table showing comparison of S. Sodium before and after phototherapy**

S. Sodium before phototherapy	S. Sodium after phototherapy			Total (%)
	<135 mmol/d (%)	135–145 mmol/d (%)	>145 mmol/d (%)	
135–145 mmol/d	39 (6 (100))	30 (88.2)	3 (60)	(86.7)
>145 mmol/d	0 (0)	4 (11.8)	2 (40)	6 (13.3)
Total	6 (13.3)	34 (75.6)	5 (11.1)	45 (100)

**Table 6: Co-relation of post phototherapy serum electrolytes with birth weight**

Electrolytes	LBW (<2.5 kg) n=17 n (%)	Normal Birth weight (≥2.5 kg) n=28 n (%)	Total	P-value
	<b>S. Sodium</b>			
< 135	3 (17.6)	3 (10.7)	6 (13.3)	0.048*
135–145	13 (76.5)	21 (75)	34 (75.6)	
>145	1 (5.9)	4 (14.3)	5 (11.1)	
<b>S. Potassium</b>				
<3.5	2 (11.8)	1 (3.6)	3 (6.7)	0.006*
3.5–5.5	14 (82.4)	22 (78.6)	36 (80)	
>5.5	1 (5.9)	5 (17.9)	6 (13.3)	
<b>S. Calcium</b>				
<6	1 (5.9)	0 (0)	1 (2.2)	0.001*
6–6.9	4 (23.5)	2 (7.1)	6 (13.3)	
7–7.9	5 (29.4)	8 (28.6)	13 (28.9)	
8–8.9	5 (29.4)	11 (39.3)	16 (35.6)	
>9	2 (11.8)	7 (25.0)	9 (20.0)	
<b>S. Magnesium</b>				
<1.6	1 (5.9)	1 (3.6)	2 (4.4)	0.516#
>1.6	16 (94.1)	27 (96.4)	43 (95.6)	
<b>S. Phosphorus</b>				
2.5	0 (0)	0 (0)	0 (0)	1.000#
2.5–4.5	17 (100)	28 (100)	45 (100)	

LBW: Low birth weight

**Table 7: Co-relation of post phototherapy serum electrolytes with gestational age (in weeks)**

Electrolytes	Gestational age (in weeks)			P-value
	<37 weeks n=16 n (%)	37–42 weeks n=29 n (%)	Total	
<b>S. Sodium</b>				
< 135	3 (18.8)	3 (10.3)	6 (13.3)	0.049*
135–145	12 (75.0)	22 (75.9)	34 (75.6)	
>145	1 (6.2)	4 (13.8)	5 (11.1)	
<b>S. Potassium</b>				
<3.5	2 (12.5)	1 (3.4)	3 (6.7)	0.003*
3.5–5.5	13 (81.3)	23 (79.3)	36 (80.0)	
>5.5	1 (6.3)	5 (17.2)	6 (13.3)	
<b>S. Calcium</b>				
<6	1 (6.3)	0 (0)	1 (2.2)	0.011*
6–6.9	3 (18.8)	3 (10.3)	6 (13.3)	
7–7.9	5 (31.3)	8 (27.6)	13 (28.9)	
8–8.9	4 (25.0)	12 (41.4)	16 (35.6)	
>9	3 (18.8)	6 (20.7)	9 (20.0)	
<b>S. Magnesium</b>				

<1.6	1 (6.3)	1 (3.4)	2 (4.4)	0.306#
>1.6	15 (93.8)	28 (96.6)	43 (95.6)	
<b>S. Phosphorus</b>				
2.5	0 (0)	0 (0)	0 (0)	1.000#
2.5–4.5	16 (100)	29 (100)	45 (100)	

**Table 8: Frequency table showing comparison of S. Calcium before and after phototherapy**

S. Calcium before phototherapy	S. Calcium after phototherapy					Total I (%)
	<6 mg/dl (%)	6–6.9 mg/dl (%)	7–7.9 mg/dl (%)	8–8.9 mg/dl (%)	>9 mg/dl (%)	
7–7.9 mg/dl	1 (100)	5 (83.3)	1 (7.7)	0 (0)	0 (0)	7 (15.6)
8–8.9 mg/dl	0 (0)	1 (16.7)	9 (69.2)	6 (37.5)	1 (11.1)	17 (37.8)
>9 mg/dl	0 (0)	0 (0)	3 (23.1)	10 (62.5)	8 (88.9)	21 (46.7)
Total	1 (2.2)	6 (13.2)	13 (28.9)	16 (35.6)	9 (20.0)	45 (100)

**Table 9: Frequency table showing comparison of S. Potassium before and after phototherapy**

S. Potassium before phototherapy	S. Potassium after phototherapy			Total
	<3.5 mmol/dl	3.5–5.5 mmol/dl	>5.5 mmol/dl	
3.5–5.5 mmol/dl	2 (66.7)	32 (88.9)	5 (83.3)	39 (86.7)
>5.5 mmol/dl	1 (33.3)	4 (11.1)	1 (16.7)	6 (13.3)
Total	3 (6.7)	36 (80.0)	6 (13.3)	45 (100)

**REFERENCES**

1. Bezboruah G. Electrolyte imbalances resulting from phototherapy in neonatal hyperbilirubinemia. IOSR J Dent Med Sci 2019;18:51-8.
2. Gregory ML, Martin CR, Cloherty JP. Neonatal hyperbilirubinemia. In: Cloherty JP, Eichenwald EC, Stark AR. Editors. Manual of Neonatal Care. 7th ed., Ch. 26. Philadelphia, PA: Lippincott-Raven; 2012, p. 304-39.
3. Arora S, Narang GS, Singh G. Serum calcium levels in preterm and term neonates on phototherapy. J Nepal Paediatr Soc 2014;34:24-8.
4. Ullah S, Rahman K, Hedayati M. Hyperbilirubinemia in Neonates: Types, Causes, Clinical Examinations, Preventive Measures and Treatments: A Narrative Review Article. Iran J Public Health. 2016;45(5):558-568.
5. Metcalf, Ryan A, Khan, Jenna, Andrews, Jennifer, Maycock, Dennis, Billimoria, Zeenia MD, Pagano, Severe ABO Hemolytic Disease of the Newborn Requiring Exchange Transfusion, Journal of Pediatric Hematology/Oncology: November 2019 - Volume 41 - Issue 8 - p 632-634
6. Reddy AT, Bai KV, Shankar SU. Electrolyte changes following phototherapy in neonatal hyperbilirubinemia. Indian J Sci Res 2015;4:752-8.
7. Alizadeh-Taheri P, Sajjadian N, Eivazzadeh B. Prevalence of phototherapy induced hypocalcemia in term neonate. Iran J Pediatr 2013;23:710-11.
8. Karamifar H, Pishva N, Amirhakimi GH. Prevalence of phototherapy-induced hypocalcemia. Iran J Med Sci 2015;27:166-8.
9. Yadav RK, Sethi RS, Sethi AS, Kumar L, Chaurasia OS. The evaluation of the effect of phototherapy on serum calcium level. Peoples J Sci Res 2012;5:1-4.
10. Eghbalian F, Monsef A. Phototherapy-induced hypocalcemia in icteric newborns. Iran J Med Sci 2002;27:169-71.
11. Kumar PS, Shankar SU. Serum sodium changes in neonates receiving phototherapy for neonatal hyperbilirubinemia. J Evid Based Med Healthc 2015;2:3982-88.
12. Suneja S, Kumawat R, Saxena RA. Effect of phototherapy on various biochemical parameters in neonatal hyperbilirubinemia: A clinical insight. Indian J Neonatal Med Res 2018;6:13-8.