



A STUDY ON THE LIPID PROFILE OF HYPERTENSIVE PATIENTS

General Medicine

Dr. Sunil Kumar Department Of General Medicine, Jhalawar Medical College, Jhalawar, (Rajasthan).

Dr. Arvind Singh Nehra Department Of General Medicine, Jhalawar Medical College, Jhalawar, (Rajasthan).

Dr. Akansha Jain Department Of General Medicine, Jhalawar Medical College, Jhalawar, (Rajasthan).

ABSTRACT

Background- Dyslipidaemia and hypertension are the commonest risk factors for coronary artery disease (CAD). To study the lipid profile in hypertensive patients.

Methods- The present study was carried out on a total of 135 hypertensive patients attending our Hospital and 135 age and sex matched healthy controls. Twelve hour fasting lipid analysis was done for Serum triglycerides (TG), total cholesterol (TC), High Density Lipoprotein (HDL), Low Density Lipoprotein (LDL).

Results- In our study mean age of hypertensive patients was 53.94 ± 8.25 yrs and in control group was 56.02 ± 6.48 yrs. 74 patients were male & 61 patients were female in hypertensive group and 68 patients were male & 67 patients were female in normotensive group. The Mean serum total cholesterol values were highly significant in hypertensive subjects (256.23 ± 50.23 mg/dL) as compared to the healthy control subjects (150.023 ± 20.01 mg/dL). The mean serum TG level was 108.36 ± 20.01 mg/dl in healthy control subjects, and 218.36 ± 30.02 mg/dl in hypertensive patients. This difference was highly significant. The mean values for HDL was 28.02 ± 6.20 mg/dL for hypertensive subjects whereas 42.03 ± 5.78 mg/dl for healthy control subjects, respectively. The Mean serum LDL values were highly significant in hypertensive subjects (160.32 ± 30.20 mg/dL) as compared to the healthy control subjects (90.23 ± 15.21 mg/dL).

Conclusion- All the lipid profile like cholesterol, LDL, TG and HDL were derange in the hypertensive than those in the healthy controls.

KEYWORDS

Hypertension ,Lipid profile, HDL ,TG , LDL.

INTRODUCTION

Dyslipidaemia and hypertension are the commonest risk factors for coronary artery disease (CAD). Recent reports show that borderline hypertension (systolic BP 130-139 and/or diastolic BP 85-89 mmHg) and Stage I hypertension carry a significant cardiovascular risk and there is a need to reduce this blood pressure¹ The reported prevalence of hypertension varies around the world, with the lowest prevalence in rural India (3.4% in men and 6.8% in women) and highest in Poland (68.9% in men and 72.5% in women).² It has been found that men have a higher prevalence of hypertension than women although this changes later in life with substantial increase in the number of females with hypertension after the age of 50 years. Dyslipidaemia (hyperlipidaemia), which is associated with hypertension, has been recognised as independent risk factor for cardiovascular disease, a leading diagnosis for visits to physicians and cause of death.³

MATERIALS AND METHODS

Patients who diagnosed as cases of essential hypertension based on history and on Antihypertensive Medication were included. The selected subjects were further grouped as:

GROUP 1: Healthy control subjects (n=135). It was ensured by routine examination that all the subjects were healthy and there were no signs and symptoms of hypertensive and other disease.

GROUP 2: Hypertensive subjects (n=135). It included the clinically established patients of hypertension. There blood pressure is in range of systolic blood pressure (>140 mmHg) and diastolic blood pressure (>90 mmHg) and have no symptoms of diabetes mellitus.

An informed consent was taken from all the healthy control subjects and patients, under study apprising them the nature and objective of the study. All subjects were studied as outpatient. Participant's examination included interviews for medical and nutritional history.

After on overnight fast of 10-12 hours, fasting blood samples were collected .Blood samples were drawn from anticubital vein of each subject by using aseptic technique. The blood was collected in plain tubes for lipid parameters respectively. Serum was separated after centrifugation and analysed.

RESULTS

In our study mean age of hypertensive patients was 53.94 ± 8.25 yrs and in control group was 56.02 ± 6.48 yrs. 74 patients were male & 61 patients were female in hypertensive group and 68 patients were male

& 67 patients were female in normotensive group. The Mean serum total cholesterol values were highly significant in hypertensive subjects (256.23 ± 50.23 mg/dL) as compared to the healthy control subjects (150.023 ± 20.01 mg/dL). The mean serum TG level was 108.36 ± 20.01 mg/dl in healthy control subjects, and 218.36 ± 30.02 mg/dl in hypertensive patients. This difference was highly significant. The mean values for HDL was 28.02 ± 6.20 mg/dL for hypertensive subjects whereas 42.03 ± 5.78 mg/dl for healthy control subjects, respectively. The Mean serum LDL values were highly significant in hypertensive subjects (160.32 ± 30.20 mg/dL) as compared to the healthy control subjects (90.23 ± 15.21 mg/dL).

Table 1. General Characteristics

Variable	GROUP-1	GROUP-2	P-value
Age in years	53.94 ± 8.25	56.02 ± 6.48	>0.05
Sex (M:F)	74:61	68:67	>0.05
TC (mg/dl)	150.023 ± 20.01	256.23 ± 50.23	<0.001
LDL (mg/dl)	90.23 ± 15.21	160.32 ± 30.20	<0.001
HDL (mg/dl)	42.03 ± 5.78	28.02 ± 6.20	<0.001
TG (mg/dl)	108.36 ± 20.01	218.36 ± 30.02	<0.001

DISCUSSION

The Mean serum total cholesterol values were highly significant in hypertensive subjects (235.36 ± 32.36 mg/dL) as compared to the healthy control subjects (152.36 ± 12.01 mg/dL). The mean serum TG level was 110.36 ± 16.12 mg/dl in healthy control subjects, and 215.23 ± 35.36 mg/dl in hypertensive patients. This difference was highly significant. The mean values for HDL was 27.23 ± 6.02 mg/dL for hypertensive subjects whereas 43.21 ± 5.36 mg/dl for healthy control subjects, respectively. The Mean serum LDL values were highly significant in hypertensive subjects (154.36 ± 31.26 mg/dL) as compared to the healthy control subjects (95.36 ± 16.02 mg/dL).

The significantly higher plasma total cholesterol, triglycerides and LDL-cholesterol in the hypertensive than in the normotensive patients in the present study is in corroboration with earlier studies.⁴⁻⁶

In accordance to our study, Saha MS et al (2006) also reported a statistically highly significant relation in serum TG level in hypertensive subjects (184.77 ± 5.97 mg/dL) as compared to the healthy controls (142.73 ± 6.68 mg/dL).⁷

Kumar NL et al (2010) reported a statistically highly significant relation in serum TG level in hypertensive subjects (180.88 ± 68.5 mg/dL) as compared to the healthy controls (114.7 ± 17.62 mg/dL).⁸

CONCLUSION

All the lipid profile like cholesterol, LDL, TG and HDL were derange in the hypertensive than those in the healthy controls.

REFERENCES

1. Kadiri S. Current concepts in the management of hypertension. Dokita 1999; 26: 93-96.
2. Shahid A, Zuberi SJ, Hasnain N. Lipid pattern in healthy subjects. Pak J Med Res 1985; 24: 33-7.
3. Youmbissi TJ, Djoumessi S, Nouedoui C. Profile lipidique d'un group d'hypertendus camerounais noir Africains. Medicine d'Afrique Noire 2001; 31: 114-118.
4. Ahaneku JE, Nwosu MC, Ahaneku GI, Okugba PC. Utilisation of Clinical chemistry tests with special reference to lipid profile in disease management in a Nigeria setting. East Afr Med J 1999; 76:172-175.
5. Mgonda YM, Ramaiya KL, Swai ABM, Mc-Larty DG, George KM, Alberti M. Insulin resistance and hypertension in non-obese Africans in Tanzania. Hypertension 1998; 31: 114-118.
6. Jarikre AE, Dim DC, Ajuluchukwu JNA. Plasma lipid levels in Nigerian hypertensives: the gender factor. Nig Qtr J Hosp Med 1996; 6: 293-298.
7. Saha MS, Sana NK and Ranajit Kumar Shaha, Serum lipid profile of hypertensive patients in the northern region of the Bangladesh. J. Bio-Sci; 2006;14: 93-98.
8. Kumar NL, Deepthi J, Rao YN, Deedi MK. Study of lipid profile, serum magnesium and blood glucose in hypertension. Biology and Medicine; 2010; 2(1): 6-16.