



A COMPARATIVE STUDY ON OPEN APPENDICECTOMY AND LAPAROSCOPIC APPENDICECTOMY

General Surgery

Dr. Prabhat Kumar

MS Senior Resident, Department of General Surgery, ANMMCH Gaya

Dr. Sanjay Kumar MS, MCH Assistant Professor Department of General Surgery ANMMCH Gaya

Dr. K.K. Sinha MS Associate Professor Department of General Surgery, ANMMCH Gaya

ABSTRACT

To compare open appendectomy with laparoscopic appendectomy in terms of operative time, postoperative recovery, wound infection, hospital stay and cost of the treatment and overall outcomes. This study was conducted in Department of general Surgery, ANMMCH, Gaya Bihar, India over a period of twelve months. Patients undergoing surgery for acute appendicitis were randomly divided into one of the two groups (A or B). In Group-A patients underwent open appendectomy and in Group B laparoscopic appendectomy was performed. Operative time, Post operatively pain chart and wound infection was recorded and, at the time of discharge, number of days in hospital and cost of treatment was calculated. Laparoscopic appendectomy is safe and effective. Wound infection and postoperative pain is significantly lower after this mode of surgery with shorter hospital stay. But the operative time is more and cost of treatment is much more in laparoscopic appendectomy. **Summary:** Laparoscopic appendectomy is as safe and effective as open appendectomy with lesser postoperative analgesia requirement and wound infections, but more operative time and overall cost of treatment. we recommend not to get biased with laparoscopic procedure as majority of Indian peoples were of low socioeconomic status, unnecessarily not to increase the economic burden, as laparoscopic appendectomy is yet to be proved as gold standard.

KEYWORDS

INTRODUCTION :-

Surgical procedure to remove the appendix is called an appendectomy. Appendectomy can be performed through open or laparoscopic surgery. Appendectomy is one of the commonly performed procedures in General surgery. McBurney described the operative technique for right iliac fossa pain using Gridiron incision in 1894. This remained the technique for appendectomy and did not change much until almost a century later, when in 1983, Semm described the first Laparoscopic appendectomy.(1-5) Laparoscopic appendectomy for suspected appendicitis is considered safe and effective.(6,7) It has gained popularity in recent years and has become one of the most widely performed procedures using the laparoscope globally.(8) However, it has not become the gold standard for acute appendicitis. This is, mainly due to the emergency nature of disease often operated with less expert assisting staff in odd hours when laparoscopic equipment, trained staff and supervision may not be available in all hospitals. In the third world countries, cost of equipment is an additional factor among those mentioned above. In our hospital, appendectomy is being performed by both techniques depending on the availability of trained staff for laparoscopy and patients' preferences. We conducted this prospective study to compare the results of open appendectomy with laparoscopic appendectomy in terms of Operative time, postoperative pain, rate of wound infection, hospital stay and cost of treatment and overall outcome.

Patients And Methods:

Between October 2021 to September 2022 (Twelve months) all patients admitted in the surgical unit with right lower quadrant pain were evaluated with the Alvarado score for appendicitis. Those who scored six or above were diagnosed as acute appendicitis, consent taken & enrolled for study and were assigned to group A or B. Group A comprised of patients that underwent open appendectomy while Group B included laparoscopic appendectomy. All patients were informed about the nature of the study and the possibility of conversion to open in case of laparoscopic group. Patients who did not want to undergo the procedure assigned to them were excluded from the enrolment process. Also, the patients whose diagnosis was unclear, had a palpable lump in abdomen and/or Alvarado score at admission was less than 6 were not included. Same surgical team performed the procedures and maximum precaution was taken to use similar techniques in all patients within each group. Use of analgesia, general and local anaesthetic agents was also kept similar for all patients. Antibiotics were given according to the department protocol (Ceftriaxone and salbactam 1.5gm and Metronidazole 500mg intravenous in single preoperative and two postoperative doses). Patients found to have gangrenous or perforated appendicitis during surgery were treated with continuation of antibiotics for 5days. Postoperative pain control for both groups was achieved by, NSAIDS

(Diclofenac sodium injection 75mg) at the end of procedure followed by oral analgesic. Intravenous or intramuscular analgesics were used if needed. Total number of days in Hospital was calculated and patients received verbal instructions to return to normal activity after discharge. First follow-up was done on seventh postoperative day for suture removal and wound assessment.

RESULTS:

Eighty patients with clinical diagnosis of acute appendicitis were randomized into groups A and B, with 40 patients in each group. Group-A patients underwent open appendectomy while patients in Group-B had laparoscopic appendectomy. Patients in the two groups were comparable with regard to their age, sex and clinical features of acute appendicitis. Migratory right iliac fossa pain and tenderness were common clinical features, followed by anorexia; nausea and elevated temperature. The mean operative time in Group A was significantly shorter than Group B patients. Oral liquids were started after 10±2 hours of surgery in open group which was significantly longer than the laparoscopic group (6±1 hours). Similarly, the laparoscopic group initiated solid diet much earlier than the open group. Use of postoperative analgesics, both injectable as well as oral, and antibiotics was significantly less in the laparoscopic group. Mean comparison of postoperative pain by visual analogue scale, was significantly low in Group B, compared with Group A, on day 0, 1 and 2. The hospital stay was 4±0.8 days in Group A and 1.5±0.06 days in Group B. None of the patients in Group B while, 26.67% patients in Group A developed postoperative wound infection at 1 week follow up.

DISCUSSION:

Appendectomy is the treatment of choice for acute appendicitis(9) and is by far the most commonly performed emergency abdominal operation.(10) Although, open appendectomy is considered a safe and effective operation for acute appendicitis with low morbidity, however, variability in the inflammatory process and the location of appendix at times causes operative difficulties. It has also been associated with potential disadvantages like post operative pain, wound sepsis and complications like intestinal obstruction which may delay recovery.(11) With the development of laparoscopic technique, it has emerged as a modus operandi for both diagnosis and treatment of acute appendicitis.(12) Studies have shown the procedure to be effective and with improved cosmesis, reduced postoperative pain, days of hospitalization and early return to work.(13- 17) Laparoscopic appendectomy was first reported in 1983 and has since been considered safe with high accuracy and complication rates as low as zero to 1.4%.(18,19) Besides, laparoscopy preserves the option of leaving a macroscopically normal appendix safely in place, it also allows localization of the area of inflammation, making it possible to

plan an incision if converting to open appendectomy, gives a better view to examine other peritoneal and pelvic organs, minimizing the chances of negative appendectomy and missing other pathologies.(20) Laparoscopic appendectomy has shown to be both feasible and safe in randomized comparisons with open appendectomy,(21) however local data on this subject is scarce. All laparoscopic appendectomy in this study were performed by one consultant surgeon to limit technique bias. Different consultants performed the open appendectomy using Gridiron (73.3%) or Lanz (26.7 %) incision. Mean operative time was 51.8 ± 7.8 minutes (45 to 90mins) for laparoscopic appendectomy and 39.6 ± 5.6 minutes (30-45 minutes) for open appendectomy. Analgesia requirement in the laparoscopic group was much less than in open group which is in line with studies performed in other setups. There was significantly less number of postoperative complications in the laparoscopy group in our study.(18,20) Others have reported infective complications after laparoscopic appendectomy to be around 20 % for gangrenous or perforated cases and 10% in simple acute appendicitis.6 Mean hospital stay was significantly less (1.5 ± 0.06 days) for laparoscopic group compared to open (4.1 ± 0.8 days). The extended hospital stay in the open group may be attributed to delay in mobilization, initiation of oral diet and the psychological effect of having an incision. Other workers have not shown this difference.(22)The significant difference in our study can be an unintentional bias in favour of laparoscopic group, however, prolonged hospital stay after surgery is usual in our not for profit (free of cost) hospital for patients coming from low socioeconomic strata. A difference in the proportion of patients with perforated and gangrenous appendix in open group (3 of each in open while one of each in laparoscopic group) and a higher rate of postoperative complications, like wound infection that could be attributed to the bad condition of appendix to start with, in this group was noted. There is a possibility that since patients with longer history and severe symptoms were not included in the study due to the exclusion criteria of Alvarado scoring, our sample has relatively less number of gangrenous and perforated cases compared to simple acute ones.

REFERENCES:

- [1]. McBurney CN. The incision Made in the Abdominal Wall in Cases of Appendicitis, with a Description of a New Method of Operating. *Ann Surg* 1894; 20: 38-43.
- [2]. Samelson SL, Reyes HM. Management of perforated appendicitis in children-revisited. *Arch Surg* 1987; 122: 691-6.
- [3]. Editorial. A sound approach to the diagnosis of acute appendicitis. *Lancet* 1987; 198-200.
- [4]. Semm K. Endoscopic appendectomy. *Endoscopy* 1983; 15: 59-64.
- [5]. Moberg AC, Berndrem F, Palmquist I, Petersson T, Resch T, Montgomery A. Randomised clinical trial of laparoscopic versus open appendectomy for confirmed appendicitis. *Br J Surg* 2005; 92: 298-304.
- [6]. Harrell AG, Licourt AE, Novitsky YW, Rosen MJ, Kuwada TS, Kercher KW, et al. Advantages of laparoscopic appendectomy in the elderly. *Am Surg* 2006; 72: 474-80.
- [7]. Young JL, Law CM, Lo CY, Cam CM. A comparative study of routine laparoscopic versus open appendectomy. *JSLs* 2006; 10: 188-92.
- [8]. Sweeney KJ, Keane FB. Moving from open to laparoscopic appendectomy. *Br J Surg* 2003; 20: 257-8.
- [9]. Amir M, Raja MH. Timings for surgery of acute appendicitis. *J Coll Physicians Surg Pak* 2000; 10: 295-7.
- [10]. Fingerhut A, Millat B, Borrie F. Laparoscopic versus open appendectomy: time to decide. *World J Surg* 1999; 23: 835-45.
- [11]. Hellberg A, Rudberg C, Kullman E, Enochson L, Fenyo G, Graffner H, et al. Prospective randomized multicenteric study of laparoscopic versus open appendectomy. *Br J Surg* 1999; 86: 48-53.
- [12]. Seymour I, Schwartz editor, principles of surgery, 7th ed. United State of America: Mc graw hill 1999, 1386.
- [13]. Lin HF, Wu JM, Tseng LM, Chen K, Huang SH, Lai LR. Laparoscopic versus open appendectomy for perforated appendicitis. *J Gastrointest surg* 2006; 10: 906-10.
- [14]. Manicini GJ, Mancini ML, Nelson HS Jr. Efficiency of laparoscopic appendectomy in appendicitis with peritonitis. *Am Surg* 2005; 71: 1-4.
- [15]. Novarra G, Ascanelli S, Turini A, Carsoforo P, Tonini G, puzza E. Laparoscopic appendectomy versus open appendectomy in suspected acute appendicitis in female patients. *Ann Ital Chir* 2002; 73: 59-63.
- [16]. Utpal D. Laparoscopic versus open appendectomy in west Bengal, India. *Chin J Dig Dis* 2005; 6: 165-9.
- [17]. Kamran H, Naveed D, Nazir A, Hameed M, Ahmed M, Khan U. The role of total leukocyte count in the diagnosis of acute appendicitis. *J Ayub Med Coll Abbottabad* 2008; 20: 70-1.
- [18]. Marzouk M, Khater M, Elsedak M, Abdelmoghny A. Laparoscopic versus open appendectomy: a prospective comparative study of 227 patients. *Surg Endosc* 2003; 17: 721-4.
- [19]. Nguyen NT, Zainabadi K, Mavandadi S, Paya M, Stevens CM, Root J, et al. Trends in utilization and outcomes of laparoscopic versus open appendectomy. *Am J Surg* 2004; 188: 813-20.
- [20]. Cox MR, McCall JL, Toouli J, Padbury RT, Wiison TG, Wattchon DA, et al. Prospective randomized comparison of open versus laparoscopic appendectomy in men. *World J Surg* 1996; 20: 263-6.
- [21]. Khalid K, Ahmad N, Farooq O, Anjum I, Sial GA. Acute appendicitis laboratory dependence can be misleading audit of 211 cases. *J Coll Physician Surg Pak* 2001; 11: 434-7.
- [22]. Utpal D. Laparoscopic versus open appendectomy: An Indian perspective. *J Minimal Access Surg* 2005; 1: 15-20.