

A STUDY OF THYROID HORMONE PATTERNS IN CHRONIC KIDNEY DISEASE

General Medicine

Dr M. Venkata Madhav

Professor, Department of Medicine, Dr. Pinnamaneni Siddhartha Institute of Medical Sciences, Chinavutapalli, Krishna District, A.P

Dr Ramya Sai Edupuganti*

Postgraduate, Department of Medicine, Dr. Pinnamaneni Siddhartha Institute of Medical Sciences, Chinavutapalli, Krishna District, A.P *Corresponding Author

ABSTRACT

Background: Kidneys have a prominent role in the metabolism, degradation and excretion of thyroid hormones. There is an increased prevalence of hypothyroidism in chronic kidney disease (CKD) patients as the glomerular filtration rate falls. **Objectives:** To study the thyroid function in chronic kidney disease patients and to study the correlation of thyroid dysfunction with respect to staging of CKD. **Materials and Methods:** A prospective Cross-sectional study was done on 100 patients with Chronic kidney disease who were not on dialysis and fulfilled all the inclusion criteria at Dr PSIMS & RF. Total and Free T3, T4 and TSH levels were estimated for those patients. **Results:** Majority of subjects included in our study were in the age group of 60-69 years with male predominance. Out of 100 patients included in our study, 50 patients (50%) were found to have low T3 syndrome; 6 patients (6%) were having hypothyroidism; 44 patients (44%) were having normal thyroid gland function. Most of the patients (n=58) were in stage 5 of Chronic kidney disease. In the present study, as creatinine clearance reduced with each CKD stage, the levels of total T3 and free T4 were also reduced. **Conclusion:** There is a positive correlation between the severity of CKD and thyroid dysfunction. Hence by studying the incidence of thyroid dysfunction in CKD patients and their subsequent treatment, it may help in slowing the progression of CKD.

KEYWORDS

chronic kidney disease, thyroid dysfunction, hypothyroidism

INTRODUCTION:

CKD is defined as abnormalities of kidney structure or function, present for >3 months, with implications for health, and requires one of two criteria documented or inferred for >3 months: either GFR <60 ml/min/1.73 m² or markers of kidney damage, including albuminuria¹.

Thyroid hormones play an important role in regulating the basal metabolic rate, development, protein synthesis and it also influences the function of some other hormones². In the kidney, thyroid hormones play a significant role in its growth and development and for the maintenance of water and electrolytes. On the other hand, kidney has a significant role in the metabolism, degradation and excretion of thyroid hormones³.

The levels of thyroid stimulating hormones in chronic kidney disease patients may be normal or increased but there is a reduced response to thyrotropin releasing hormone⁴. Circadian rhythm and activity are altered, suggesting abnormality in the level of hypophysis⁴. According to various studies, prevalence of thyroid dysfunction is found to be ranging from 13% in early CKD to 70% End stage renal disease^{5,6,7,8}.

This study was conducted to find out the thyroid dysfunction in CKD patients and also to estimate the correlation between thyroid dysfunction with respect to various stages of chronic kidney disease.

MATERIALS AND METHODS:

A single center prospective cross-sectional study with a sample size of 100 CKD patients who attended Dr. PSIMS & RF were included in the study after meeting the inclusion and exclusion criteria and were evaluated for thyroid function.

Inclusion criteria:

Known CKD patients who are above the age of 18 years diagnosed as per the KDIGO 2012 guidelines.

Exclusion criteria:

1. Acute kidney injury
2. Patients on dialysis
3. Previous history of thyroid illnesses.
4. Recent onset of acute illness e.g., surgery, trauma etc.
5. Pregnancy
6. Known liver disease
7. Drugs altering thyroid profile like amiodarone, steroids, dopamine, phenytoin etc.

RESULTS:

The present study included 100 CKD patients out of whom, 74% (n=74) were males and 26% (n=26) were females.




Figure 1: Sex-wise distribution of study population

Majority of patients were in 60-69 yrs. age group (34%, n=34) followed by 50-59 yrs. age group (32%, n=32). Age of the patients ranged from 35-69 years.




Figure 2: Age-wise distribution of study population

The prevalence of various stages of CKD in present study population were 58% (n=58) in stage 5 CKD, 32% (n=32) in stage 4 CKD, 8% (n=8) in stage 3 CKD and 2% (n=2) in stage 2 CKD.




Figure 3: Prevalence of various stages of CKD in study population

In the present study, the prevalence of low T3 syndrome was 50%, TSH in hypothyroidism range was 6% and 44% of the patients had normal thyroid function.




Figure 4: Prevalence of thyroid dysfunction in the study population

In the present study, as creatinine clearance reduced with each CKD stage, the level of mean total T3 also reduced (103.5 in stages 1-3 vs 69.95 in stage 5). This trend is also followed with free T4 (1.25 in stages 1-3 vs 0.94 in stage 5). This difference is statistically significant $P < 0.05$. The difference with TSH was not statistically significant $P > 0.05$.

Table 1: Thyroid dysfunction in various stages of CKD

CKD stage	N	Total T3	Total T4	Free T3	Free T4	TSH
1-3	10	103.5±28.76	7.01±1.07	1.89±0.47	1.25±0.11	1.85±1.81
4	32	91.03±36.06	6.17±1.16	1.89±0.66	1.09±0.24	1.16±0.79
5	58	69.95±23.76	4.73±1.45	1.47±0.54	0.94±0.29	4.54±1.369
p value		<0.001	<0.001	0.02	0.001	0.3

The present study showed a positive correlation between total T3, free T4 and creatinine clearance and it was statistically significant. It also showed a negative correlation between TSH and creatinine clearance which is not statistically significant.

Table 2: Correlation between creatinine clearance and various thyroid hormones levels

Relation with Cr Clearance	r value	p value
Total T3	0.32	0.001
Total T4	0.51	<0.001
Free T3	0.23	0.02
Free T4	0.38	<0.001
TSH	-0.13	0.2

DISCUSSION:

The present study showed to have 16% between 30 and 39 yrs., 18% between 40 to 49 yrs., 32% between 50 and 59 yrs. and 34% between 60 to 69 yrs. age groups. A study done by Aruna Kumari et al showed 10% in 21-30 yrs, 11.66% in 31-40 yrs., 20% in 41-50 yrs., 41.66% in 51-60 yrs. and 16.66% in >60 yrs. age groups⁹. In contrast to the present study, a study done by Hossain et al showed 11% in 20-29 yrs., 20% in 30-39 yrs., 33% in 40-49 yrs. and 36% in 50-60 yrs. age groups¹⁰.

In the present study, 100 patients of CKD were taken among which 74% of them were males and 26% were females. Avasthi et al showed 73.33% males and 26.67% females¹¹. In contrast to the present study, the study done by Gowda et al showed significantly higher proportion of males (78%) than females (22%)¹².

In the present study, the prevalence of various stages of CKD in the study population was 2%, 8%, 32% and 58% of total population in stages 2 to 5 of CKD respectively, with highest proportion of patients in CKD stage 5 group. Avasthi et al showed similar prevalence of CKD with 46.7% in stage 4 and 53.3% in stage 5 population¹¹. In contrast to the present study, higher prevalence (48%) was observed among stage 4 in a study done by Benjamin et al¹³.

In the present study, the prevalence of low T3 syndrome was 50%, TSH in hypothyroidism range was 6% and 44% of the patients had normal thyroid function. Similar results are seen in a study done by Abhinaya et al in which 40% were found to have low T3 syndrome, 16% had TSH in hypothyroidism range and 10% were in subclinical hypothyroidism and 34% had normal thyroid function.¹⁴ In contrast to the present study, majority of the CKD patients had normal thyroid function in a study done by Chonchol et al (84.6% - normal thyroid function, 9.5% - subclinical hypothyroidism and 5.9% - subclinical hyperthyroidism)¹⁵.

The present study showed positive correlation between total T3, free T4 and creatinine clearance and it was statistically significant $P < 0.05$. It also depicts a negative correlation between TSH and creatinine clearance and this was not statistically significant $P > 0.05$. In a study done by Khatriwada et al, a significant positive correlation between free T3, free T4 and eGFR and a significant negative correlation between TSH and eGFR was noted¹⁶. Similar significant results were obtained in another study done by Shin et al 2013¹⁷.

CONCLUSIONS:

Out of the 100 CKD patients taken in the study, males are relatively higher in proportion when compared to females in a ratio of approximately 3:1. The proportion of patients with thyroid dysfunction is 56%. Low T3 syndrome is the most prevalent followed by low T4. The proportion of patients with low T3 and T4 syndromes increased with the staging of the CKD. A significant positive correlation between total T3, free T4 and creatinine clearance was observed in this study. By studying the incidence of thyroid dysfunction in CKD patients and their subsequent treatment, it may help in slowing the progression of CKD.

REFERENCES:

- Levin, A., Stevens, P. E., Bilous, R. W., Coresh, J. et al (2013). Kidney disease: Improving global outcomes (KDIGO) CKD work group. KDIGO 2012 clinical practice guideline for the evaluation and management of chronic kidney disease. Kidney International Supplements, 3(1), 1-150.
- Gupta, A., K., Virmani, S., & Arora, M. (2017). Thyroid dysfunction in patients of chronic kidney disease. International Journal of Advances in Medicine, 4(5), 1333-1337.
- Singh, S., Verma, A., Aryal, G., Thapa, S., Khakurel, S., & Shrestha, K. (2016). Thyroid hormone profile in patients with chronic kidney disease: a single centre study. Journal of Nepal Health Research Council, 14(34), 197-201.
- Iglesias, P., & Diez, J. J. (2009). Thyroid dysfunction and kidney disease. European journal of endocrinology, 160(4), 503-515.
- Kher V. (2002). End-stage renal disease in developing countries. Kidney international, 62(1), 350-362.
- Prabhar, M. R., Chandrasekaran, V., & Soundararajan, P. (2008). Epidemic of chronic kidney disease in India -what can be done? Saudi journal of kidney diseases and transplantation: an official publication of the Saudi Center for Organ Transplantation, Saudi Arabia, 19(5), 847-853.
- Srinath Reddy, K., Shah, B., Varghese, C., & Ramadoss, A. (2005). Responding to the threat of chronic diseases in India. Lancet (London, England), 366(9498), 1744-1749.
- Prajapati, P., Singh, A. P., & Bendwal, S. (2013). Correlation between severity of chronic kidney disease and thyroid dysfunction. Journal of the Indian Medical Association, 111(8), 514-516.
- T. Dr & G. Dr. (2020). A Study of thyroid dysfunction in chronic kidney disease Patients in a tertiary Care Hospital - A Prospective study. International Journal of Advanced Biochemistry Research. 4. 20-26.
- Hossain, M. M., Shah, K., Begum, N., Ahmed, P., Islam, A. M., & Chowdhury, N. I. (2018). Thyroid Functional Status in Chronic Kidney Disease. Bangladesh Journal of Nuclear Medicine, 18(2), 141-144.
- G Avasthi, S Malhotra, APS Narang, S Sengupta. Study of thyroid function in patients of chronic renal failure. Indian J Nephrol. 2001;11(4):165-169.
- Gowda M. Evaluation of thyroid function status in patients with chronic kidney disease. Journal of medical science and clinical research. 2016 Nov;4(11):14235-41.
- Benjamin V, Kirubakaran S, Elavarasan K, Natarajan S. A prospective study on the evaluation of thyroid function status in patients with chronic kidney disease. Journal of medical science and clinical research. 2017 Sep;5(9):28007-14.
- Abhinaya, V. and Kumar, S. M. Thyroid Function Test in Patients with Chronic Kidney Disease. J Pharm Res Int. 2021 Nov 6;33(48A):150-158.
- Chonchol M, Lippi G, Salvagno G, Zoppini G, Muggeo M, Targher G. Prevalence of subclinical hypothyroidism in patients with chronic kidney disease. Clin J Am Soc Nephrol. 2008 Sep;3(5):1296-300.
- Khatriwada S, Rajendra KC, Gautam S, Lamsal M, Baral N. Thyroid dysfunction and dyslipidemia in chronic kidney disease patients. BMC Endocr Disord. 2015 Oct 29;15:65.
- Shin DH, Lee MJ, Lee HS, Oh HJ, Ko KI, Kim CH et al. Thyroid hormone replacement therapy attenuates the decline of renal function in chronic kidney disease patients with subclinical hypothyroidism. Thyroid. 2013 Jun;23(6):654-61