



“ALL-ON-FOUR” TECHNIQUE AS A SAFE APPROACH IN REHABILITATION OF SEVERELY ATROPHIC JAW: A CASE REPORT

Maxillofacial Surgery

Dr. K.Chandana Vani	Senior Resident, Department of Oral and maxillofacial surgery, Government Dental College and Hospital, Hyderabad, Telangana
Dr. Ranjitha Balineni	Senior Resident, Department of Oral and maxillofacial surgery, Government Dental College and Hospital, Hyderabad
Dr. Kodam Narsimha	Senior Resident, Department of oral and maxillofacial surgery, Government Dental College and Hospital, Hyderabad
Dr. Yasaswini Tallaprodatur	Senior Resident, Department of oral and maxillofacial surgery, Government Dental College and Hospital, Hyderabad

ABSTRACT

The severely atrophied maxilla restricts implant placement mainly in the posterior region when using conventional techniques. The “All-on-4” concept could be a good alternative consisting in placing four implants in the anterior part of the fully edentulous jaw to support provisional, fixed and immediately loaded prosthesis. The two most anterior implants are placed axially, and the two posterior implants are placed in an angled position. This approach can be considered a clinically effective treatment with affordable time and cost, allowing a more simple procedure, less morbidity and higher patient quality of life. This clinical case describes the steps to restore an atrophic maxilla following the “all-on-four” concept, with immediate load.

KEYWORDS

All on four implants, Resorbed ridge, Edentulous jaw

INTRODUCTION:

The severely atrophied maxilla and mandible represents a constant challenge and restricts implant placement in the posterior region when using conventional techniques. This limitation is due to the reduced quantity and quality of available bone, as well as the presence of maxillary sinus pneumatization. Different approaches have been developed to rehabilitate atrophic arches. Among proposed techniques we can count bone graft and maxillary sinus elevation, but these types of interventions are not easily accepted by patients because of their invasive nature, postsurgical morbidity, and high cost. The “All-on-4” concept could be a good alternative consisting in placing four implants in the anterior part of the fully edentulous jaw to support provisional, fixed and immediately loaded prosthesis. The two most anterior implants are placed axially, and the two posterior implants are placed in an angled position [3, 4] to avoid the maxillary sinus, reduce the cantilever length, improve bone anchorage and increase polygonal area for prosthesis support [5,6]. The present clinical report describes a method to restore an atrophic ridges with the “All-on-Four” technique, using the concept of immediate load.

Case Report:

A 32-year-old man, edentulous for a long period of time due to periodontal disease, reported to the institution, seeking for a fixed maxillary prosthesis. His past medical history was uneventful. Preliminary impressions were made with alginate in both dental arches. Jaw relation record was obtained. After esthetic try-in, dentures were fabricated and the maxillary prosthesis was duplicated in order to obtain a multi-functional radiological and surgical guide. Panoramic radiograph and cone beam tomography revealed an advanced alveolar bone resorption in the posterior maxilla and mandible (Fig1). Considering the presented situation, the All-on-four concept with immediate loaded prosthesis was suggested to rehabilitate upper and lower jaw.

Surgery was performed under local anaesthesia. A full thickness crestal incision was made along with anterior releasing incision in the area of the midline. A high speed diamond round bur with copious external irrigation, was used to identify the anterior wall of the maxillary sinus. Once the anterior wall of the sinus was located, surgical sequence for implant placement was undertaken. The two most posterior implants with the diameter of 3.75 mm and the length of 13 mm were placed anterior to the mesial wall of the sinus, tilted mesially. Then, two anterior implants with dimension of 3.7 mm diameter and 11.5 mm length were placed in the anterior maxilla parallel to the midline of the jaw. Anterior implants are typically emerged at the lateral or central incisor position, posterior implants are

placed in the second premolar or first molar region. This arrangement allowed for good implant anchorage, short cantilever length, and large interimplant distance. In mandible, the two most posterior implants with the diameter of 4.2 mm and the length of 15 mm were placed anterior to the mental foramen, tilted mesially. Then, two anterior implants with dimension of 3.5 mm diameter and 13 mm length were placed in the anterior mandible parallel to the midline of the jaw [fig 2-3]. After the implant placement, primary closure is done using 3-0 vicryl suture and patient is kept under standard antibiotic therapy.

The definitive prosthesis was fabricated 4 months post-surgically. It consisted of metal-acrylic resin implant-supported fixed prosthesis with Cr-Co framework and acrylic resin prosthetic teeth. Multiunit impression coping [Fig3-4] were attached to the multi-unit abutments and splinted with wire-bars and a low shrinkage autopolymerizing resin. The impression was taken with polyether impression material (3M ESPE Impregum soft) using the pick-up technique. Once the impression was taken, multiunit analogs were placed into the impression and the working model was obtained. After trying the screw-retained metal framework on multiunit abutments and evaluating their passive fit, they were returned to the laboratory for tooth set-up at the previously established vertical dimension occlusion. The final prosthesis was then fabricated and delivered after adjusting the occlusion. The patient was satisfied with his extraoral and intraoral esthetics, function and phonetics (Fig5,6). He was scheduled for regular follow-up.



Fig 1: Post operative orthopantomogram




Fig 2: maxillary and mandibular multiunit abutment




Fig 3: maxillary and mandibular all on four implant placement



Fig 4: maxillary multiunit abutments

Fig 5: bisque try in

Fig 6: final prosthesis

DISCUSSION:

The challenge today is not to prove functionality but, rather, to develop simple, reproducible and cost-effective protocols [3]. The all-on-four concept has been proven to be clinically effective treatment for completely edentulous jaw with affordable time and cost through immediate rehabilitation, by providing relatively straightforward and predictable treatment, thereby ensuring a high outcome of quality of life [9, 10].

The use of angulated implant in the maxilla has been demonstrated to be an alternative to bone grafting [7, 6]. By tilting the distal implant, the cantilever extension decreases and the length of the prosthesis supported polygon increases achieving a better load distribution [11]. In addition, the tilting may also provide good cortical anchorage and improve primary stability allowing the use of longer implants. The "All-on-Four" procedure also improves cortical anchorage and primary stability, allowing the use of longer implants. In a three-dimensional finite element analysis about load transmission using different implant inclinations and cantilever lengths, Bevilacqua et al. Reported a reduction of stress around anterior implants in a full fixed prosthesis design, when tilted implants were compared to straight implants.

A distal position of the posterior implant, reducing the cantilever length, achieves a more favorable stress distribution in the bone. Bevilacqua et al. demonstrated that tilting of the distal implant decreased the level of stress bone, when compared to axial implants supporting a higher number of cantilevered teeth [5]. Furthermore, implant tilting does not induce any biological disadvantage. Strain gauge measurements performed by Krekmanov showed no significant differences in forces and bending moment between tilted and non tilted implants. From a biomechanics point of view, the position of the coronal end of the implant is more important than its inclination, whether the implant is tilted or not [6]. Immediate loading of implant supported full-arch prostheses is considered a predictable technique [11]. Maló et al., in a 1-year retrospective study, reported a high survival rate of 97,6% [4]. The immediate loading procedure improved patient acceptance of implant treatment [8]. It has shown a high level of satisfaction for patients in terms of esthetics, phonetics, and functionality [14]. According to Callandriello R et al., the use of tilted implants and immediate load decreases treatment time, reduces surgical invasion and represents a lower cost [10]. In particular, it may avoid months of complete edentulism or the wearing of an uncomfortable denture. These factors benefit the patient as well as the clinician [8, 16]. To accomplish immediate function, all-acrylic prosthesis is usually placed within a few hours after surgery [7]. However, the most frequent prosthetic complication is the acrylic provisional prosthesis fracture, which usually occurs close to the coping, a relatively weak region [9, 8]. Maló et al., observed acrylic fracture as 27% [3].

Strengthening prosthesis with metal wire and compliance with the soft diet is suggested to prevent them [8]. Furthermore, tooth fractures in definitive acrylic resin prostheses are a frequent complication, especially when the opposing teeth are natural, and they seem to have a higher incidence comparing to cases with full denture or implant-

supported overdenture, because of the application of greater forces

CONCLUSION:

The all-on-four concept can be an alternative treatment option especially in atrophic edentulous jaws. It represents a less invasive and financially competitive treatment, with reduced time procedure and a higher patient quality of life

REFERENCES:

1. Clavero J, Lundgren S. Ramus or chin grafts for maxillary sinus inlay and local onlay augmentation: comparison of donor site morbidity and complications. *Clin Implant Dent Relat Res.* 2003;5:154-160.
2. Sannino, G. All-on-4 Concept: A 3-Dimensional Finite Element Analysis. *J. Oral Implantol.*, 41(2):163-71, 2015
3. Malo P, Rangert B, Nobre M. „All-on-Four“ immediate-function concept with Branemark System implants for completely edentulous mandibles: a retrospective clinical study. *Clin Implant Dent Relat Res.* 2003; 5(suppl 1):2-9.
4. Malo P, Rangert B, Nobre M. All-on-4 immediate-function concept with Branemark System implants for completely edentulous maxillae: a 1- year retrospective clinical study. *Clin Implant Dent Relat Res.* 2005;7(suppl 1): S88-S94.
5. Bevilacqua M, Tealdo T, Menini M, et al. The influence of cantilever length and implant inclination on stress distribution in maxillary implant supported fixed dentures. *J Prosthet Dent.* 2010;105:5-13.
6. Krekmanov L, Kahn M, Rangert B, Lindstrom H. Tilting of posterior mandibular and maxillary implants for improved prosthesis support. *Int J Oral Maxillofac Implants.* 2000;15:405-414.
7. Maló P, de Araújo Nobre M, Lopes A, Francischone C, Rigolizzo M. "All-on-4" immediate-function concept for completely edentulous maxillae: A clinical report on the medium (3 years) and long-term (5 years) outcomes. *Clin Implant Dent Relat Res* 2012;14(suppl 1): e139-e150.
8. Di, P.; Lin, Y.; Li, J. H.; Luo, J.; Qiu, L. X.; Chen, B. & Cui, H. Y. The All-on-Four implant therapy protocol in the management of edentulous Chinese patients. *Int. J. Prosthodont.*, 26(6):509- 16, 2013.
9. Soto-Peñaloza, D., Zaragoza-Alonso, R., Peñarrocha-Diogo, M., & Peñarrocha-Diogo, M. (2017). The all-on-four treatment concept: Systematic review. *J Clin Exp Dent.* 2017;9(3):e474-88.
10. Ben hadj Hassine M, Bucci P, Gasparro R, Di lauro AE, Sammartino G. Safe approach in "All-on-four" technique: a case report. *Annali di Stomatologia* 2014; 142 V (4): 142-145
11. Falk H, Laurell L, Lundgren D. Occlusal interferences and cantilever joint stress in implant-supported prostheses occluding with complete dentures. *Int J Oral Maxillofac Implants.* 1990; 5(1): 277-283.