



“PULMONARY FUNCTION DERANGEMENT IN PATIENTS OF DIABETES MELLITUS”

General Medicine

**Dr Manjunath
Gavade**

Post graduate in Department of General medicine , GIMS ,Gadag , Karnataka,

**Dr Jagadeesh
Gaddeppannavar***

Associate Professor, Department of General Medicine, GIMS, Gadag, Karnataka
*Corresponding Author

**Dr Irfanahmed M
Makandar**

Senior resident, Department of pulmonology, GIMS, Gadag, Karnataka

**Dr Prakash
Hosamani**

Assistant Professor, Department of medicine, GIMS, Gadag, Karnataka

ABSTRACT

Background: Diabetes mellitus is a metabolic disorder characterized by hyperglycaemia resulting from defects in insulin secretion, action or both. Nongenyzmatic glycosylation induced alteration of lung connective tissue is the most likely mechanism underlying the mechanical pulmonary dysfunction in diabetic subjects. The present study evaluates the impact of diabetes mellitus on pulmonary function in this cross sectional study of population. **Aims & Objective:** To evaluate the impact of diabetes mellitus on pulmonary functions by comparing with control groups. **Materials And Methods:** All patients presenting in Department of Medicine, Gadag institute of medical sciences, Gadag. who were proved cases of type 2 diabetes mellitus were included in this study from December 2020 to December 2021. 50 patients and 50 controls with matched age and sex were included. All patients were interviewed, detailed history was taken with respect to risk factors and detailed physical examination was carried out and appropriate investigations were carried out. **Results:** FVC among diabetics was low with a mean of 2.46 as compared to controls with mean of 2.99 with p value of 0.001 which is statistically significant. FEV1 was reduced in cases with mean of 2.0542 compared to controls with mean of 2.4906 with a p value of 0.001 being statistically significant. FEV1/ FVC was higher in diabetic group with mean value of 0.8304 as compared with controls with mean value of 0.8316. **Conclusion:** Diabetic patients show reduced lung function with a restrictive pattern of lung disease on the Spirometry.

KEYWORDS

Pulmonary function, Diabetes mellitus.

INTRODUCTION:

Diabetes mellitus is a metabolic disorder characterized by hyperglycemia resulting from defects in insulin secretion, action or both. Diabetes is a multisystem disorder that affects many organs of the body¹. This metabolic disorder is a risk factor precipitating micro vascular pathologies leading to autonomic neuropathy, nephropathy, retinopathy and peripheral neuropathy, and macro vascular pathologies leading to coronary artery disease, cerebrovascular accidents and peripheral vascular disease². There are histopathological changes seen in lungs of diabetics such as thickened alveolar epithelial and pulmonary capillary basal lamina leading to reduced pulmonary elastic recoil and lung volume. Nongenyzmatic glycosylation induced alteration of lung connective tissue is the most likely mechanism underlying the mechanical pulmonary dysfunction in diabetic subjects³.

This suggests that lung is also a target organ⁴. Spirometry is a widely used pulmonary function test (PFT), ideally suited to describing the effects of obstruction or restriction of lung function. The spirometric parameters have gained more popularity when it has been reported that impaired Forced Vital Capacity (FVC) and Forced Expiratory Volume in 1 sec (FEV1) are emerging novel risk factors for diabetes mellitus⁵. In spite of this, spirometry is not used routinely as part of a management system in diabetic patients. Its role is neither fully explored, nor fully utilized to achieve quality of life when managing diabetes mellitus. The present study is undertaken to evaluate the impact of diabetes mellitus and pulmonary function in this cross sectional study of population.

MATERIALS AND METHODS:

Inclusion Criteria includes

All patients presenting in Department of Medicine, Gadag institute of medical sciences, Gadag. who were proved cases of type 2 diabetes mellitus were included in this study from September 2017 to December 2018. Selection of participants: 50 cases and 50 controls with matched age and sex were included.

Exclusion Criteria :

1. Previous history of lung disease 2. Signs and symptoms of

respiratory infections at the time of test 3. History of being admitted during past six months with respiratory symptoms 4. History of cardiovascular illness 5. Alcoholic 6. Obesity 7. Smokers

Data were collected by using performa meeting the objectives of the study. Purpose of the study was carefully explained to patients and consent was taken. All patients were interviewed, detailed history was taken with respect to risk factors and detailed physical examination was carried out and appropriate investigations done which included complete blood counts, fasting blood sugar, post-prandial blood sugar, HbA1C, X ray chest (PA view), Spirometry (FVC, FEV1, FEV1/FVC values recorded) Pulmonary function tests(PFT) of diabetic patients and controls were compared by applying student's unpaired t-test while correlations between FVC , FEV1 , HbA1C and duration of illness in diabetic patients were analyzed by applying Pearson's co-efficient.

RESULTS:

A total of 100 subjects were included in the present study 50(29 males and 21 females) of them were diabetics and the other 50(32 males and 18 females) were non diabetic controls. 50% cases belonged to the age group of 40-49 years, 24% to 50-59 years and 26% to 60 and above age group. Similar distribution was assured in controls with 38%, 34% and 28% respectively in the three age groups. The mean age in both groups being 51.43 The glycemic parameters which included the FBS, PPBS and HbA1c are 204.86mg/dl, 254.30mg/dl and 8.1474% respectively for cases and 96.58, 127.04 and 4.802 respectively for controls.(Table I) Body mass index (BMI) of cases and controls were comparable with mean among study group being 24.407 and in controls its value was 25.18. The difference was not statistically significant with a p value of 0.209. The spirometric values in both groups can be seen summarized in Table II. The FVC was 2.4616+0.42553 and 2.9998+0.43302 in cases and controls respectively while value of FEV1 was 2.0542+0.3999 and 2.4906+0.36564 in the same manner. Thus making the ratio of FEV1/FVC almost similar. The study also tried to correlate the severity of impairment of the lung function test with duration of diabetes mellitus. Two groups were taken- one with disease duration 5 years. The glycemic parameters of the two groups are shown in Table III. It was observed that glycemic control was better in long term diabetics(>5years duration) than the newly diagnosed ones(<5 years).

Table I: Comparison of glycemic parameters among cases and controls

Parameter	Group	N	Mean	S.D.
FBS	Cases	30	204.86	57.59
	Controls	30	84.48	15.52
PPBS	Cases	30	204.30	57.59
	Controls	30	127.04	17.70
HbA1c	Cases	30	8.5476	1.23736
	Controls	30	5.6222	0.59687

FBS - Fasting Blood Sugar, PPBS - Post Prandial Blood Sugar, HbA1c-

Hemoglobin A1cN = sample size S.D. = Standard Deviation

Table III: Comparison of duration (<5 yrs and >5 yrs) with FBS and PPBS scores in cases by t test

Parameter	Duration	N	Mean	S.D.	P value
FBS	<5 years	22	214.9915	56.2289	0.012
	>5 years	28	198.0000	40.8176	
PPBS	<5 years	22	217.8231	54.7559	0.007
	>5 years	28	198.0000	40.8176	

Table II: Comparison of cases and controls with FVC, FEV1 and FEV1/FVC

Parameter	Group	N	Mean	S.D.	P value
FVC	Cases	30	2.4618	0.40353	0.000
	Controls	30	2.8988	0.43302	
FEV1	Cases	30	2.0942	0.3988	0.000
	Controls	30	2.4888	0.3884	
FEV1/FVC	Cases	30	0.8504	0.0509	0.000
	Controls	30	0.8504	0.0509	

*P<0.05

Table IV: Comparison of duration (<5 yrs and >5 yrs) with FVC, FEV1 and FEV1/FVC

Parameter	Duration	N	Mean	S.D.	P value
FVC	<5 years	22	2.3077	0.4178	0.000
	>5 years	28	2.7037	0.3302	
FEV1	<5 years	22	1.6823	0.3704	0.000
	>5 years	28	2.0538	0.3884	
FEV1/FVC	<5 years	22	0.8015	0.0584	0.000
	>5 years	28	0.8015	0.0584	

*P<0.05

DISCUSSION:

Present study was undertaken to assess ventilatory function of subjects with diabetes mellitus in comparison to subjects without diabetes. Larger population-based studies have been more consistent, demonstrating reduced pulmonary functions in patients with an elevated plasma glucose level and a diagnosis of DM. The present study is in agreement with the previous study⁶⁻¹⁰ (Table V) and the inference drawn from this study is similar to those studies. The study proves the hypothesis of reduced FVC in diabetics compared to their age, sex and BMI matched controls. The study also shows coherence the study of Walter E Robert et al¹¹ who studied the relationship between diabetes mellitus and pulmonary function and showed a decrease in FVC by 109 ml and FEV1 by 27ml in diabetes mellitus. Davis M.E. Timothy¹² studied the pulmonary function and its association with Type-2 diabetes mellitus and showed an average decrease of 9.5% in both FVC and FEV1 of diabetics. Different studies⁶⁻¹⁰ noting the same results are compared in table VI. Due to the combined reduction in FEV1 and FVC the FEV1/FVC ratio yielded the value of 0.83 which is suggestive of a restrictive pattern. Studies by Muhammad Irfan et al⁸ (0.819), Sultan⁷ (0.851) and Sanjeev Verma et al⁹ (0.90) had similar results. The present study, thus, is in agreement with previous studies which have shown almost same values of FEV1/FVC in diabetic subjects when compared to age, sex and BMI matched controls. The key aspect of the present study is the observation that although diabetic subjects had low FVC compared to controls the group of diabetics with diabetes mellitus less than 5 years had lower FVC than in those with duration > 5 years. FVC in diabetics with 5 years (28 study subjects) it was 2.70 only marginally (but significantly) less than the controls. These results were contrary to the study done by Sultan⁷ where there was no significant reduction in FVC in diabetics of 12 years duration (FVC=3.11) compared to controls (FVC=3.74). (Figure 1) When the natural history of disease was compared on the time line with regard to FEV1 we still observed the same improving pattern with diabetics >5 years (comparison shown in table VII and figure II). While unlike FVC, FEV1 in Sultan's⁷ study showed similar pattern to that of ours with a decline in diabetics Masmoudi and Zouari¹³ concluded that pulmonary volumes impairments are slightly more marked with diabetes mellitus duration especially after 10 years. Barret-Connor E et al¹⁴ also found that pulmonary function in older adults is altered in subjects with diabetes mellitus with duration more than 10 years. The improvement in respiratory function can be hypothesized to a better metabolic control. Our results are similar to those of Lange P¹⁵ who found that the decrease of ventilator capacity is more marked at the beginning of diabetes mellitus. Davis A Wendy et al¹⁶ who studied the glycemic exposure and associated reduced pulmonary function in type 2 diabetes found a decrease in FVC at an annual rate of 68ml/year and FEV1 at an annual rate of 71 ml/year. This linear decline was not noticed in our study which may be the reflection of very high sugar levels at the diagnosis and eventual better control which is also reflected in a small scale six years study by Ramirez L.C. et al¹⁷ (1991) demonstrating that intensive treatment by subcutaneous insulin infusion improved both FVC & FEV1 percentage predicted values. Recently Nakajima et al¹⁸ reported reduced FVC and normal FEV1/FVC and concluded that impaired restrictive pulmonary function but not the obstructive pattern might be associated with metabolic disorders and metabolic syndrome in a severity dependant manner.

CONCLUSION:

Lungs also bear the brunt of uncontrolled diabetes mellitus as evident by this study where mean reduction in FVC was 500ml and in FEV1 was 360 ml with the parameters showing a restrictive pattern. Thus lung is also a target organ for diabetes mellitus.

Conflicts Of Interest:

There are no conflicts of interest.

REFERENCES:

- Shlomo Melmed Kenneth Polonsky P. Reed Larsen Henry Kronenberg. Williams Textbook of Endocrinology 13th Edition 2015
- Benbassat CA, Stern E, Kramer M, Lebzelter J, Blum I, Fink G. Pulmonary functions in patients with diabetes mellitus. Am J Med Sci 2001;322:127-32
- Dalquen P. The lung in diabetes mellitus. Respiration 1999;66:12-3
- Sandler Malcom. Is the lung a 'target organ' in diabetes mellitus? Arch Intern Med. 1990 Jul;150(7):1385-8
- SA, Meo. (2010). Significance of spirometry in diabetic patients. International Journal of Diabetes Mellitus. 2. 47-50. 10.1016/j.ijdm.2009.12.003.
- Klein, O. L., Kalhan, R., Williams, M. V., Tipping, M., Lee, J., Peng, J. and Smith, L. J. (2012). Lung spirometry parameters and diffusion capacity are decreased in patients with Type 2 diabetes. Diabetic Medicine, 29: 212-219
- Sultan Ayoub Meo. Significance of spirometry in diabetic patients International Journal of Diabetes Mellitus 2 (2010) 47-50.
- Muhammad Irfan, Abdul Jabbar, Ahmed Suleman Haque, Safia Awan, and Syed Fayyaz Hussain. Pulmonary functions in patients with diabetes mellitus. Lung India. 2011 Apr-Jun; 28(2): 89-92
- Sanjeev Verma, Mumtaz Goni, Kudyar P. Assessment of pulmonary functions in patients with Diabetes Mellitus. JK science April-June 2009; vol. 11, No.2
- Anand R. Dharwadkar, Asha A. Dharwadkar, Gouher Banu, Shrilaxmi Bagali reduction in lung functions in type-2 diabetes in Indian population : correlation with glycemic status Indian J Physiol Pharmacol 2011; 55 (2): 170-175.
- Walter ERobert, Alexa Beiser, Rachel J, Givelber, George T, O'Connor, et al. Association between glycemic state and lung function. Am J Respir Crit Care Med 2003; 167:911-916
- Davis Timothy M E, Matthew Knuiman, Peter Kendall, Hien Vu, Wendy A Davis. Reduced pulmonary function and its associations in type 2 diabetes: Diabetes research and Clinical practice 2000; 50:152-159. 13. K. Masmoudi, N. Zouari, respiratory function evolution in a duration of four years in patients with diabetes mellitus J.I. M. Sfax, N°13/14; Juin07/Déc07: 27-29.
- Barrett Connor E, Frette C. NIDDM, impaired glucose tolerance, and pulmonary function in older adults. The Rancho Bernardo Study. Diabetes Care 1996;19(12):1441-4
- Lange P. Diabetes mellitus, plasma glucose and lung function in a cross sectional population study. Eur Respir J 1989;2:14-19.
- Wendy A. Davis, Matthew Knuiman, Peter Kendall, Valerie Grange, Timothy M.E. Davis, Glycemic Exposure Is Associated With Reduced Pulmonary Function in Type 2 Diabetes The Fremantle Diabetes Study 27:752-757, 2004.
- Ramirez LC, Dal Nogare A, Hsia C, Aruaz C, Butt I, Trowig SM, Schnurr-Breen L, Raskin P. Relationship between diabetes control and pulmonary function in insulin-dependent diabetes mellitus. Am J Med 91:371-376, 1991
- Nakajima K, Kubouchi Y, Muneyuki T, et al. A possible association between suspected restrictive pattern as assessed by ordinary pulmonary function test and the metabolic syndrome. Chest 2008; 134 (4): 712-18