



A STUDY OF EMERGENCE OF HANSEN'S IN A TERTIARY CARE HOSPITAL IN THE ERA OF ERADICATION

Microbiology

Dr. S. Sindhuja	Post graduate, Department of Microbiology, Andhra Medical College, Visakhapatnam, Andhra Pradesh, India.
Dr. G. A. Praveen	Post graduate, Department of Microbiology, Andhra Medical College, Visakhapatnam, Andhra Pradesh, India.
Dr. B. Aruna Sree	MD, Associate Professor, Department of Microbiology Andhra Medical College, Visakhapatnam, Andhra Pradesh, India
Dr. P. Venkata Ramana*	MD, Assistant Professor, Department of Microbiology Rangaraya Medical College, Kakinada, Andhra Pradesh, India *Corresponding Author
Dr. P. Ratna Kumari	MD, Professor, Department of Microbiology Andhra Medical College, Visakhapatnam, Andhra Pradesh, India
Dr. P Kamala	MD, Professor and HOD, Andhra Medical College, Department of Microbiology, Visakhapatnam, Andhra Pradesh, India

ABSTRACT

Introduction: Hansen's disease is a chronic infectious disease caused by *Mycobacterium Leprae*. India declared elimination of leprosy in December 2005, but a slow resurgence of the disease still continues. **Aims And Objectives:** To analyse the clinical and bacteriological findings of new and recurrent cases of Hansen's disease in the post elimination era. **Material And Methods:** Study includes a total of 292 slit-skin smears received from Department of Dermatology, Venereology and Leprosy to the Department of Microbiology for a period of 18 months (Jan 2020 to June 2022) from patients. All samples are processed according to standard microbiological procedures. **Results:** A total of 276 patients for a period of 2 years 6 months were included. The most common clinical spectrum was Lepromatous Leprosy followed by Borderline Lepromatous Leprosy. Out of 292 SSS, 181 were positive. **Conclusion:** The study brings forth evidence on the slow re-emergence of Leprosy in India.

KEYWORDS

Hansen's disease, leprosy, slit-skin smear, infection, resurgence

INTRODUCTION:

Mycobacterium leprae is the cause of the chronic infectious disease Hansen's disease. It mainly affects the skin, nasal mucosa and peripheral nerves. *Mycobacterium leprae*, is an acid-fast bacillus which is transmitted via droplets from the nose and mouth during close contacts with untreated cases. It eventually lead to disability, disfiguration and social stigma for the rest of patient's life if untreated. Therefore, early detection of *M. leprae* is the key element to timely identification and treatment of patients before nerve involvement occurs. Leprosy was declared eradicated in India in December 2005, however the illness has been slowly resurgent ever since. Leprosy is primarily diagnosed clinically, however straightforward slit-skin smear microscopy helps with an accurate diagnosis. Slit-skin smears, which have an almost 100% specificity when done correctly, continue to be the most basic diagnostic method until new state-of-the-art diagnostic technologies are regularly made accessible for use at the bedside.

AIMS AND OBJECTIVES:

To analyse the clinical and bacteriological findings of new and recurrent cases of Hansen's disease in the post elimination era

MATERIAL AND METHODS:


This is a descriptive, hospital-based, retrospective study of newly diagnosed patients with Hansen's disease enrolled in the Hansen's disease clinic in Department of Dermatology, Venereology and Leprosy outpatient department (OPD) and Department of Microbiology, Andhra Medical College-King George Hospital, Visakhapatnam for a period of 2 years 6 months (January 2020 to June 2022) from 276 patients.

The current study sought to examine the clinical and bacteriological results of newly diagnosed cases of Hansen's disease in the post-elimination period. This research enrolled treatment-naïve, fresh instances of Hansen's disease identified using WHO criteria. Sample size of Study includes a total of 292 slit-skin smears. All samples are processed according to standard microbiological procedures for SSS by Acid Fast Staining and Microscopy. A retrospective chart review of the patients was performed to acquire information regarding clinical features and SSS findings.

RESULTS:

In this study, a total of 276 patients met inclusion criteria (newly diagnosed cases, old cases, all ages and sex) were included, 151 (54.72%) were found to be male and 125 (45.28%) female. The age range between 46 and 74 years is more represented in the current study, with a total of 120 (43.47%) cases.

The most common clinical spectrum was Lepromatous Leprosy followed by Borderline Lepromatous Leprosy. Out of 292 SSS, 181 (61.9%) were positive and 111 (38.1%) were negative.



DISCUSSION:

STUDY	SLIT SKIN SMEAR POSITIVE
Neirita Hazarika et al	39.6%
Kakkad et al	70%
Bhushan. P. et al	56.57%
Gopala Rao B et al	59.31%
Present study	61.9%

Distribution Based On Gender


STUDY	MALE	FEMALE	TOTAL
N.Hazarika et.al	79	37	116
Shraddha et.al	27	23	50

P.Bhushan et.al	102	39	141
P.Premalatha et.al	26	4	30
Present Study	151	125	276

Generally, leprosy is believed to be more common in males. The present study also showed same. Male predominance may be because of many factors such as industrialization, urbanization and more opportunities for contact in males.



Classical case of Lepromatous Leprosy (LEONINE FACIES)



Age Wise Distribution (n=276)

1-20 yrs, 57(in no.), 20.65%

21-40yrs, 120(in no.), 43.47%

41-60yrs, 71(in no.), 29.58%

61-80yrs, 28(in no.), 11.66%

CONCLUSION:

The study brings forth evidence on the slow re-emergence of Leprosy in India. In this study, Multibacillary cases outnumber the Paucibacillary cases. There is a need to take steps to curb the further spread of Hansen's disease in the community.

REFERENCES:

1. Mahajan VK. Slit-skin smear in leprosy: lest we forget it! *Indian J Lepr.* 2013 Oct-Dec;85(4):177-83. PMID: 24834639.
2. Hazarika N, Gupta PK, Dhanta A, Singh A, Mohanty A, Gupta P. Renaissance of Hansen's Disease in Post-Elimination Era in North India: A Retrospective Clinico-Bacteriological Study. *Cureus.* 2021 Aug 28;13(8):e17514. doi: 10.7759/cureus.17514. PMID: 34603886; PMCID: PMC8476198.
3. Premalatha P, Renuka IV, Meghana A, Devi SI, Charyulu P, Sampooran G. Utility of Bacillary Index in Slit Skin Smears in Correlation with Clinical and Histopathological Alterations in Hansen's Disease: An Attempt to Revive a Simple Useful Procedure. *Ann Med Health Sci Res.* 2016 May-Jun;6(3):181-4. doi: 10.4103/2141-9248.183936. PMID: 27398251; PMCID: PMC4924493.
4. Bhushan P, Sardana K, Koranne R V, Choudhary M, Manjul P. Diagnosing multibacillary leprosy: A comparative evaluation of diagnostic accuracy of slit-skin smear, bacterial index of granuloma and WHO operational classification. *Indian J Dermatol Venereol Leprol* 2008;74:322-326
5. Kumaran, S. M., Bhat, I. P., Madhukara, J., Rout, P., & Elizabeth, J. (2015). Comparison of Bacillary Index on Slit Skin Smear with Bacillary Index of Granuloma in Leprosy and its Relevance to Present Therapeutic Regimens. *Indian Journal of Dermatology*, 60(1), 51-54. <https://doi.org/10.4103/0019-5154.147791>
6. Kamal R, Natrajan M, Katoch K, Katoch VM. Evaluation of diagnostic role of in situ PCR on slit-skin smears in pediatric leprosy. *Indian J Lepr.* 2010 Oct-Dec;82(4):195-200. PMID: 21434596.
7. Banerjee S, Biswas N, Kanti Das N, Sil A, Ghosh P, Hasanooor Raja AH, Dasgupta S, Kanti Datta P, Bhattacharya B. Diagnosing leprosy: revisiting the role of the slit-skin smear with critical analysis of the applicability of polymerase chain reaction in diagnosis. *Int J Dermatol.* 2011 Dec;50(12):1522-7. doi: 10.1111/j.1365-4632.2011.04994.x. PMID: 22097999.
8. Siwakoti S, Rai K, Bhattarai NR, Agarwal S, Khanal B. Evaluation of Polymerase Chain Reaction (PCR) with Slit Skin Smear Examination (SSS) to Confirm Clinical Diagnosis of Leprosy in Eastern Nepal. *PLoS Negl Trop Dis.* 2016 Dec 27;10(12):e0005220. doi: 10.1371/journal.pntd.0005220. PMID: 28027305; PMCID: PMC5189933.
9. Lima FR, de Paula NA, Simões MMR, Manso GMD, Albertino GS, Felisbino GC, Antunes VMG, Perecin FAMC, Westin AT, Luga HB, Frade MAC. Bacilloscopy and polymerase chain reaction of slit-skin smears and anti-phenolic glycolipid-Iserology for Hansen's disease diagnosis. *Front Med (Lausanne).* 2022 Aug 10;9:972244. doi: 10.3389/fmed.2022.972244. PMID: 36035405; PMCID: PMC9399463.
10. Demiss W, Van Henten S, Takarinda KC, Kamau EM, Abdela SG. Slit-skin smear for the classification of leprosy: are we wasting time and resource? *J Infect Dev Ctries.* 2022 Aug 31;16(8.1):3S-7S. doi: 10.3855/jidc.15992. PMID: 36156495.