



EFFECTS OF ESMOLOL ON THE PREVENTION OF HEMODYNAMIC RESPONSE TO TRACHEAL INTUBATION

Medical Science

**Dr Hemant
Chourey**

Pg3rd yr ssmc rewa

Dr Ajay Singh

Assistant Professor BKMC sagar

**Dr Arsha P
Santhosh**

PG 3rd yr ssmc rewa

ABSTRACT

Introduction: Endotracheal extubation is almost always associated with haemodynamic changes. This increase in sympatho-adrenal activity may result in hypertension, tachycardia and arrhythmias and is more hazardous to the patients with hypertension, myocardial insufficiency or cerebrovascular disease. **Aims:** To compare the effects of esmolol in attenuating the cardiovascular responses to tracheal extubation. **Material And Method:** After ethical committee's approval, this observational study was carried out on ASA class I-II patients. After pre-anaesthetic checkup, 60 patients were randomly allocated to 2 equal groups by using chit method. Patients were given 10 ml saline in Group – S, and 1mg/kg of esmolol diluted in 10 ml saline in Group – E. These drugs were given 3 minutes after reversal and 2 minutes before extubation. **Statistical Analysis:** The results were analyzed using ANOVA, student's paired t-test and chi-square test. 'p' value of <0.05 was statistically significant and <0.001 was highly significant. **Results:** Esmolol had greater attenuating effects on haemodynamic changes than control group. Attenuation on changes of heart rate and blood pressure was greater with esmolol. **Conclusion:** esmolol 1mg/kg i.v. given at 2 min before extubation was of value in attenuating the cardiovascular changes during tracheal extubation & emergence from anaesthesia. Esmolol is more effective in attenuating the heart rate and systolic blood pressure.

KEYWORDS

INTRODUCTION

Haemodynamic changes due to reflex sympathetic discharge caused by epipharyngeal and laryngopharyngeal stimulation are almost always associated with endotracheal extubation.

This increase in sympatho adrenal activity may result in hypertension, tachycardia and arrhythmias 1,2. This increase in blood pressure and heart rate are usually transitory, variable, unpredictable and more hazardous to the patient with hypertension, myocardial insufficiency or cerebrovascular diseases 3.

Many pharmacological agents have been used to reduce the extent of haemodynamic events accompanying extubation e.g esmolol, alfentanil, fentanyl, diltiazem 1,2,7. Esmolol is an ultrashort-acting, highly cardioselective beta-adrenergic receptor antagonist 8 extensively to blunt the haemodynamic responses associated with laryngoscopy and tracheal extubation.

METHOD

Study type : Observational study

Duration of study: one year and 6 months

Sample size:

Selected 60 patients were allocated into equal Two groups by chit method.

After obtaining approval of the ethical committee & informed consent 60 patients of ASA status I & II were randomly divided into 2 groups using chit method (20 patients per group). Patients were given 10 ml saline in Group – S and 1mg/kg of esmolol diluted in 10 ml saline in Group – E. These drugs were given 3 minutes after reversal and 2 minutes before extubation.

Patients belonging to ASA I&II status, age between 18-60 years of either sex, posted for surgery under general anaesthesia were included whereas patients with cardiovascular or respiratory disease, history of allergy to study drugs, comorbidities like hypertension, diabetes mellitus and anticipated difficult intubation were excluded from the study.

In the operation theatre i.v line was secured with 18G veinflow & standard monitors ECG, NIBP, SPO2 were attached, baseline values noted. All patients were pre-medicated with inj. glycopyrrolate 0.004mg/kg i.v., inj. ondansetron 0.1mg/kg, inj. midazolam 0.02

mg/kg i.v., inj. tramadol 1mg/kg i.v. Standard maintenance i.v. fluids continued prior to induction of anaesthesia as per NPO guidelines. Preoxygenation with 100% O for 3 minutes done. Induction was done 2 with inj. Propofol 1.5-2mg/kg i/v. Tracheal intubation was facilitated with inj. succinylcholine 1.5 to 2 mg/kg i.v. Maintenance of anaesthesia was with O₂, N₂O, Isoflurane (0.2 -1%), inj. vecuronium 2 mg loading dose 0.1 mg/kg i.v and intermittently 0.02 mg/kg i.v. Pulse, NIBP, ECG, SPO₂ were monitored throughout anaesthesia. After 2 completion of surgery isoflurane and N₂O were discontinued and 2 residual muscle relaxation was reversed by inj. neostigmine (0.07 mg/kg) + inj. glycopyrrolate (0.001 mg/kg) i/v. Three minutes after reversal, 1 mg/kg esmolol diluted to 10 ml or saline 10 ml were given. Extubation was done 2minutes after administering these drugs. Immediately before tracheal extubation we confirmed that patients could breathe spontaneously and open the eyes on command. Oropharyngeal secretions were aspirated just prior to extubation. Laryngoscopy was not performed during extubation. After tracheal extubation 100% O₂ was given for 5 minutes.

The systolic blood pressure (SBP), diastolic blood pressure (DBP), and heart rate (HR) measured at the end of surgery served baseline values. Hemodynamic data obtained from 3 minutes after the injection of neostigmine-glycopyrrolate (reversal agents) was analyzed for cardiovascular changes associated with emergence from anesthesia and tracheal extubation. Values of SBP, DBP, HR and Mean arterial pressure (MAP) recorded after completion of reversal, at the time of administration of esmolol or saline (3 minutes after reversal-taken as 0 minute) and at 1, 2, 3, 4, 5, 7, 10 and 15 minutes after administration of drugs and these values were also compared with baseline values within individual study groups. Any haemodynamic effects of i.v injection of drugs were monitored for first 2 hours post surgery .

Study Parameters –

HR, SBP, DBP and MAP

RESULTS:

Table 1: Changes in mean Heart Rate between Esmolol and Saline groups

Conditions	Esmolol (n=30)	Saline (n=30)
Preoperative	92.5 ± 12.18	80.45±10.95
Reversal	96.45 ±15.60	86.65±10.93
During drug administration	97.2 ±14.45	91.65±9.32
After 1 minute	82.3 ±13.04	96.2±10.30

Extubation	74.4 ±11.63	100.05±11.40
E1	70.3 ±11.35	109.9±9.08
E2	68.4 ±10.79	102.1±9.26
E3	69±9.83	98.7±8.65
E5	71±9.48	96.4±8.66
E10	74±9.31	94.45±8.81
E 15	78.4±9.61	82.1±7.72

Table 2: Changes in mean SBP & DBP between Esmolol and Saline group

Conditions	Systolic BP (Each group n=30)		Diastolic BP (Each group n=30)	
	Esmolol	Saline	Saline	Esmolol
Pre operative	130±5.2 7	128.1±7. 07	82.3±4. 95	70.7±4.2 0
Reversal	137.75±5. 67	133.95±6. 21	83.6±6. 66	73.55±4. 91
During drug administration	132.6±6. 49	137.3±5. 64	82.9±7. 85	73.3±3.3 8
After 1 minute	128±5.7 3	140.45±6 .38	79.55±7 .22	76.75±3. 76
Extubation	126.65± 5.33	143.05±4 .90	79.3±6. 19	77.8±4.3 4
E1	127±4.9 2	148.65±6 .40	78.9±6. 88	85.3±6.0 2
E2	126.65± 5.11	144.7±5. 13	78.9±6. 37	84.65±6. 83
E3	128.9±7. 21	141.2±4. 16	80.45±6 .88	82.5±6.5 4
E5	129.9±7. 57	138.85±3 .73	81.25±7 .60	80.05±6. 09
E10	132±7.4 5	136.3±4. 49	81.75±6 .60	78±5.30
E15	133.78± 7.16	136±4.69	82.6±5. 92	76.1±4.7 1

The SBP in the group E was found to be significantly lower than those in group S during extubation and 1,3 and 5 minutes after extubation. The DBP in the group E was significantly lower than those in group S during extubation and at 1, 3 and 5 minutes after extubation

Values of systolic blood pressure, diastolic blood pressure, mean arterial pressure and heart rate were significantly lower in the esmolol than in the control group of patients. Patients from the esmolol group required significantly less opioids and isoflurane and recovered after the anaesthesia significantly faster than the patients in the control group. Esmolol also seemed to reduce the incidence of myocardial ischemia in noncardiac surgery without increasing the episodes of hypotension and bradycardia.

DISCUSSION

Tracheal extubation often provokes hypertension and tachycardia as does tracheal intubation due to reflex sympathetic discharge caused by pharyngeal and laryngeal stimulation. This stimulation is associated with increase in plasma epinephrine concentration. These cardiovascular responses to tracheal extubation are probably of little consequence in healthy individuals, but may be more severe and more hazardous in patients with hypertension, coronary artery disease and cerebrovascular disease 9.

Several methods and drugs have been used to provide haemodynamic stability during extubation and in early post-operative period 4,5,6. A number of pharmacological agents including lidocaine, esmolol, alfentanil, fentanyl and diltiazem have been recommended for the 9 attenuation of haemodynamic changes associated with extubation.

We studied patients in ASA physical status I & II without any cardiovascular disease. This population was chosen to ensure the safety.

In this study we compared the effects of intravenous Esmolol 1mg/kg (Group E) & Saline 10ml (Group S) given 3minutes after inj.neostigmine and glycopyrrolate and 2 minutes before extubation.

All the study groups were assessed for any significant differences between the demographic data such as Age variation, male: female ratio. It was then concluded that there was significant differences in the demographic data between the 2 groups [P>0.05].

The onset of antihypertensive action of esmolol was approximately 30 sec after a single i.v. injection with a peak effect occurring at 1.5- 2 minutes 9. The decision to give the drug 2 minutes prior to tracheal extubation was based on this data. The same also holds true regarding

esmolol. Intravenous administration of the mixture of neostigmine and atropine increases heart rate within 1 minute, the effect peaking 1-2 minutes after injection. The heart returns to basal values 3 minutes after injection 10. This was the rationale for administering esmolol, saline, 3 minutes after reversal for attenuation of hemodynamic response to extubation.

Esmolol was used by Andrew Dyson 8 et al in 1990 to attenuate the cardiovascular responses associated with extubation in a dose of 1mg/kg, 1.5mg/kg, 2mg/kg. They have concluded that the increase in heart rate that occurs during extubation can be successfully attenuated by bolus injection of 1 mg/kg esmolol, although this dose is insufficient to effectively block increases in SBP. A dose of 1.5 mg/kg blocks the maximal increase in HR and controls SBP. Doses of 2 mg/kg produce significant (>20%) decreases in SBP without further attenuation of the pressor or HR responses. The use of esmolol before extubation may be beneficial in patients with ischemic heart disease and good left ventricular function, especially in the presence of borderline hypertension.

Similarly In our study, by using esmolol 1mg/kg during extubation, increase in heart rate was attenuated successfully and this dose did not have much effect on SBP, DBP& MAP.

Dae Hue Nam 11 et al in 1996 used esmolol in the dose of 1.5 mg/kg and diltiazem in the dosage of 0.2mg/kg to attenuate cardiovascular responses to tracheal extubation. They have concluded that at a bolus dose of i.v. esmolol 1.5mg/kg or diltiazem 0.2mg/kg given 2 min before extubation was of value in attenuating the cardiovascular changes occurring in association with tracheal extubation and emergence from anaesthesia. Esmolol is more effective than diltiazem in attenuating the heart rate changes. Diltiazem is more effective than Esmolol in attenuating the systolic blood pressure changes. Thus their findings were comparable to our study also Habib Boston, Ahmed Eroglu 13 in 2012 had used esmolol in the dose of 1 mg/kg in preventing the haemodynamic responses to endotracheal intubation and extubation. They concluded that when administered before induction and emergence from anaesthesia, 1mg/kg of esmolol is effective in controlling the haemodynamic response to laryngoscopy, intubation and extubation. Thus their findings were comparable to ours. 1/2 Esmolol, with its rapid onset and extremely short duration of action (t -9 minutes), would appear to be ideal drug for preventing acute increase in heart rate and systolic blood pressure. However, we would advice caution when using bolus dose of esmolol during extubation unless the patient has received atropine or glycopyrrolate (as part of the reversal of neuromuscular blockade) because tracheal stimulation in the presence of beta blockade may potentially produce profound bradycardia.

CONCLUSION

We concluded that SBP, DBP and heart rate in the control group increased significantly in association with tracheal extubation. Esmolol hydrochloride given in dose of 1 mg/kg i.v. given 3 minutes after reversal and 2 minutes before extubation is more effective in decreasing the blood pressure than the heart rate during tracheal extubation.

REFERENCES

- Kahoru Nishina, Katsuya Mikawa, Nobuhiro Maekawa and Hidefumi Obara. Attenuation of cardiovascular responses to tracheal extubation with diltiazem. *Anesthesia Analgesia* 1995; 80:1217-22.
- Katsuya Mikawa, Kahoru Nishina, Nobuhiro Maekawa and Hidefumi Obara. Attenuation of cardiovascular responses to tracheal extubation: verapamil Vs diltiazem. *Anaesthesia Analgesia* 1996; 82: 1205-10.
- Yoshitaka Fujii, Yuhjisa Koh, Shinji Takahashi. Combined diltiazem and lidocaine reduces cardiovascular responses to tracheal extubation and anaesthesia emergence in hypertensive patients. *Canadian journal of anaesthesia* 1999; 46:952-956.
- Y Unal, O Ozsoylar, D Saryguney, M Arslan, RS Yardim. The efficacy of esmolol to blunt the haemodynamic response to endotracheal extubation in lumbar disc surgery. *Research journal of medical sciences Medwell journals* 2008; 2 (2): 99-104.
- Philippe Grillo, Nicolas Bruder, Pascal Auquier, Daniel Pellissier, and Francois Gouin. Esmolol blunts the cerebral blood flow velocity increase during emergence from anaesthesia in neurosurgical patients. *Anaesthesia Analgesia* 2003; 96:1145-9.
- Murat Alp Alkaya, Kemal Tolga Saraçoğlu, Gökhan Pehlivan, Zeynep Eti, Fevzi Yılmaz Göğüş. Effects of esmolol on the prevention of haemodynamic responses to tracheal extubation after craniotomy operations. *Turkish Journal of Anaesthesia* 2014; 42: 86-90.
- Kyutak Lee, Yoonsuk Lee, Haekyoung Kim, Chookun Chung. Effect of diltiazem on mean arterial pressure and heart rate during tracheal extubation in hypertensive patients. *Korean Journal of Anaesthesiology* 1998; 35:300-305.
- Andrew Dyson, Philip A. Isaac. Esmolol Attenuates Cardiovascular Responses to Extubation. *Anaesthesia Analgesia* 1990; 71:675-8.
- Bidwai AV, Bidwai VA, Rogers CR, Stanley TH. Blood pressure and pulse rate responses to endotracheal extubation with and without prior injection of lidocaine. *Anesthesiology* 1979; 51: 171-173.
- Muzzi, Black S, Losaso TJ, Cucchiara RF. Labetolol and Esmolol in the control of

- hypertension after intracranial surgery. *Anesthesia Analgesia* 1990; 70 68-7.
11. Dae Hue Nam, Byeong Sun Park, Hoon Soo Kang. Cardiovascular response of esmolol and diltiazem to tracheal extubation. *Korean J. Anaesthesiology* 1996; 31: 43-48.
 12. Yuji Morimoto, Yoshiko Morimoto, Osamu Kemmotsu et al. The effect of calcium channel blockers on cerebral oxygenation during tracheal extubation. *Anaesthesia Analgesia* 2000; 91: 347-52.
 13. Habib Bostan, Ahmet Eroglu. Comparison of the clinical efficacies of fentanyl, esmolol and lidocaine in preventing the hemodynamic responses to endotracheal intubation and extubation. *Journal of current surgery* 2012; 2 (1): 24-28.