



RARE OCCURRENCE OF FAHR'S DISEASE IN A YOUNG BOY WITH SEIZURES – A CASE REPORT

General Medicine

Dr. Anjaneya Prasad V*

Professor & HOD, , Department of Internal Medicine, Dr. PSIMS and RF, Chinnoutpally, Gannavaram Mandal, Krishna District, Andhra Pradesh, India *Corresponding Author

Dr. Venkata Lohit Velivela

Senior House surgeon, Department of Internal Medicine, Dr. PSIMS and RF, Chinnoutpally, Gannavaram Mandal, Krishna District, Andhra Pradesh, India

ABSTRACT

Idiopathic calcification of basal ganglia and cerebellar dentate nucleus is synonymous with Fahr's disease, which is often an autosomal dominant, rarely occurring neurodegenerative disorder. Few sporadic autosomal recessive cases have also been described. Symptomatology of the Fahr's disease ranges from movement disorders, Parkinsonism like symptoms to cognitive defects. Majority of the cases reported were in adults with typical presentation. Here we report a rare occurrence of Fahr's disease in a young boy presented only with seizures.

KEYWORDS

Calcification, Basal Ganglia, Seizures

INTRODUCTION

Idiopathic symmetric calcification of basal ganglia, cerebellar dentate nuclei of globus pallidus region and centrum semiovale is synonymous with Fahr's disease [1]. Various types of movement disorders and dementia are its prominent clinical features [2]. Extrapyramidal and pyramidal tract symptoms such as gait disturbances, paresis, spasticity, chorea, tremors, dystonia, myoclonia and parkinsonism like symptoms have been associated with this disease. Cognitive impairment, behavioural disturbances such as apathy, intermittent disinhibition, anxiety, ritualistic and anti-social behaviour, irritability, frequent mood changes and psychosis are the psychological disturbances attributed to Fahr's disease [3, 4]. It has been demonstrated that Fahr's disease is autosomal dominant, most likely linked to chromosome 14q and is frequently familial [5, 6]. Some sporadic autosomal recessive cases have also been reported [6]. There have been studies associating TULIP1 coding in Fahr's disease but these are far from being conclusive [7,8,9]. The pathophysiology of Fahr's disease is less understood but it has been hypothesized that one or the combination of the following factors such as- local disruption of blood- brain barrier, metastatic calcification secondary to disturbances in calcium metabolism resulting from hypoparathyroidism, hyperparathyroidism, pseudohyperparathyroidism, anoxia, irradiation, toxins and systemic disorders [10]. If it is metastatic calcification indeed, particular proclivity for above detailed structures needs also to be elicited.

We report a singular case of idiopathic Fahr's disease in a young male with seizures, which is an uncommon presentation.

Case Report:

An eighteen years old male patient presented to the casualty with recurrent episodes of tonic-clonic convulsions. He was a known case of complex partial seizures with secondary generalization on antiepileptic medications since last five years. His symptoms were insidious in onset, gradually progressed over years. Seizures occurred randomly once or twice in 2-3 days. There was no history of fall, headache, neck rigidity, diplopia, vomiting, post-ictal weakness, behavioural disturbances, movement disorders or gait disturbances. He was reasonably well at his studies. He was born to non-consanguineous parents and his birth was by normal vaginal delivery and there were no neonatal complications. He had attained his mile stones normal for his age. There was no history suggestive of previous illness other than seizures. There was no history of similar complaints or parkinsonism like features and behavioural disturbances in his family members

On examination, his vitals were within normal limits. His BMI was 18.75. General survey was unremarkable, except for hypertrichosis all over the body which might be secondary to the long term use of phenytoin. His IQ was normal for his age. His mini mental status examination was normal. He had attained secondary sexual characters normal for his age. His systemic examination was unremarkable

Routine investigations such as- CBP, Peripheral smear, ESR, Blood

sugar, Serum Creatinine, Electrolytes, Serum Calcium & Phosphate, Urine examination, Stool microscopy, Chest X Ray and ECG were within normal limits. He was screened for HIV, Hepatitis B and Hepatitis C, Tuberculosis, Malaria and other infectious diseases and all of them were negative.

Assays of thyroid and parathyroid hormones were normal. His Vitamin D Assay and Serum ceruloplasmin were normal.

Cranial MRI scan (Figure 1) revealed multiple T1W hyper intense and T2W is to mildly hypo intense areas involving bilateral dentate nuclei, bilateral head of caudate nuclei, bilateral lentiform nuclei, bilateral thalami and bilateral frontal and parietal gyri. Lesions are profoundly hypo intense on gradient sequences and have shown near symmetry. Complimentary plain cranial CT (Figure 2) revealed dense symmetrical calcifications involving the regions of brain parenchyma as described in the MRI findings. Patient was diagnosed to be suffering from Fahr's disease based on the above radiological features.




Figure 1: MRI brain showing T2W hypointense areas involving bilateral head of caudate and lentiform nuclei




Figure 2: Plain CT Brain showing dense symmetrical calcification of bilateral caudate and lentiform nuclei

DISCUSSION:

Clinical diagnosis of Fahr's disease is based on clinical, radiological features and by the exclusion of other causes. None of the symptoms classically described in literature detailing Fahr's disease were present in this patient [11]. Diagnosis was arrived at only after reviewing the radiological reports. Absence of family history further confirms that this is a sporadic, probably autosomal recessive case of Fahr's disease [6]. Age of onset of the disease in this case is much earlier than that of most of the cases described in the literature [12]. The differential diagnosis of Fahr's disease includes- hypoparathyroidism, pseudohypoparathyroidism, hyperparathyroidism, other causes resulting in metastatic calcification such as- neoplasms, cerebrovascular lesions, infectious diseases such as- toxoplasmosis, syphilis and inflammatory conditions such as- SLE, which have been ruled out in this case [13]. Prenatal or genetic testing is not available yet for counselling for Fahr's disease. There is neither a cure nor a definitive treatment strategy. This patient was treated symptomatically and antiepileptics have been stepped up to the maximum tolerable doses. It's hard to predict the prognosis in this case at this stage.

CONCLUSIONS:

The age of onset of the Fahr's disease can be much earlier i.e. in teens and behavioural disturbances and movement disorders may not be associated always with this disease. The extent of calcium deposition in the brain need not correlate with the symptomatology. The prognosis of this disease is variable and is very difficult to predict. Secondary causes of abnormal calcium deposition in brain need to be searched for before establishing the diagnosis. Fahr's disease may be added to the differential diagnosis of seizures.

REFERENCES:

1. Smits MG, Gabree'ls FJ, Thijssen HO et al. Progressive idiopathic strio-pallido-dentate calcinosis (Fahr's disease) with autosomal recessive inheritance. Report of three siblings. *Eur Neurol* 1983;22:58-64.
2. Shakibai SV, Johnson JP, Bourgeois JA. Paranoid delusions and cognitive impairment suggesting Fahr's disease. *Psychosomatics* 2005;46:569-572.
3. Manyam BV, Walters AS, Narla KR. Bilateral striopallidodentate calcinosis: clinical characteristics of patients seen in a registry. *Mov Disord*
4. Lam JS, Fong SY, Yiu GC, Wing YK. Fahr's disease: a differential diagnosis of frontal lobe syndrome. *Hong Kong Med J* 2007;13(1):75-7. Victor M, Ropper AH (2001)
5. Calcification of vessels in basal ganglia and cerebellum (hypoparathyroidism and Fahr syndrome). In: Victor M, Ropper AH (eds) *Principles of neurology*. McGraw-Hill, New York, pp 1032-1033.
6. Geschwind DH, Loginov M, Stern JM. Identification of a locus on chromosome 14q for idiopathic basal ganglia calcification (Fahr disease). *Am. J. Hum. Genet.* 65: 764-772, 1999.
7. Brodaty H, Mitchell P, Luscombe G et al. Familial idiopathic basal ganglia calcification (Fahr's disease) without neurological, cognitive and psychiatric symptoms is not linked to the IBGC1 locus on chromosome 14q. *Hum Genet* 2002;110:8-14
8. Oliveira JR, Lemos RR. Linkage studies in familial idiopathic basal ganglia calcification: separating the wheat from the chaff. *Am J Med Genet B Neuropsychiatr Genet* 2009;150B:594-595
9. Oliveira JR, Sobrido MJ, Spiteri E et al. Analysis of candidate genes at the IBGC1 locus associated with idiopathic basal ganglia calcification ("Fahr's disease"). *J Mol Neurosci* 2007;33:151-154
10. Malik R, Panday VK, Naik D. Fahr's disease. A rare neurodegenerative disorder. *Int J Radiol Image* 2004;14:383-4.
11. Sobrido MJ, Hopfer S, Geschwind DH. Familial idiopathic basal calcification. *GENE Reviews* 2007; 1-15
12. Yasuhiko B, Daniel FB, Rayan JU, Michael LH, Zbigniew KW: Hereditary familial Brain Calcinosis Syndrome; *Mayo Clin Proc.* 2005;80(5):641-651.
13. Goodwin RH. Computed tomographic image of Fahr disease mistaken for acute hemorrhagic cerebrovascular accident. *Am J Emerg Med* 2006;24(3):378