



ACUTE FATTY LIVER OF PREGNANCY- CASE STUDY AT TERTIARY CARE HOSPITAL

Obstetrics & Gynaecology

Dr. Devineni Sindhu

Postgraduate, Department of Obstetrics and Gynaecology, Alluri Sitarama Raju Academy of Medical Sciences Eluru, West Godavari District Andhra Pradesh 534005 India

Dr. D. Murmu

Professor, Department of Obstetrics and Gynaecology, Alluri Sitarama Raju Academy of Medical Sciences Eluru, West Godavari District Andhra Pradesh 534005 India

ABSTRACT

Introduction Acute fatty liver of pregnancy (AFLP) is life threatening complication which is rare, usually presents in second and third trimesters of pregnancy. [1] Underlying pathology of disease thought to be linked to an abnormality in free fatty acid metabolism and exact pathology is not known. [2] Early identification, supportive care and expeditious delivery are the main stay of treatment of AFLP. **Material and methods** Observational prospective study of cases of Acute Fatty Liver of Pregnancy with various presentations and their outcomes are studied over a period of 1 year (December 2022- November 2023) at obstetrics and gynaecology department at ASRAM medical college, Eluru, Andhra Pradesh after getting approval from Institutional ethics committee and consent from patients. Total 7 patients were diagnosed using Swansea criteria, laboratory investigations specific to AFLP and subjected to imaging. **Results** Out of 7 patients with AFLP, 5 were primi gravida and 2 were second gravida. Period of gestation at which they presented was between 32-36 weeks. 6 required preterm termination of pregnancy. 4 underwent LSCS (Lower Segment Cesarean Section). Average hospital stay was 12 days. Complications seen were DIC (57%), Liver Failure (57%), MODS (57%), Hypoglycaemia (57%) and renal insufficiency (100%) and 1 maternal death. **Conclusion** Early diagnosis, prompt delivery, adequate supportive care and a multidisciplinary approach are the key to good outcome.

KEYWORDS

Acute Fatty Liver of Pregnancy (AFLP), LSCS, Liver failure, Swansea criteria.

INTRODUCTION:

Acute fatty liver of pregnancy (AFLP) is a life threatening complication which is rare, usually presents in the second and third trimesters of pregnancy [1] sometimes in the early postpartum period also. It was first described by Sheehan as acute yellow atrophy of liver in 1940 [2]. Underlying pathology of disease thought to be linked to an abnormality in foetal free fatty acid metabolism and exact pathology is unknown. [3]. Acute fatty liver of pregnancy (AFLP) is characterised by microvesicular fatty infiltration of hepatocytes without any inflammation or necrosis.

It has an incidence of approximately 1 in 10,000 to 15,000 pregnancies [4]. In past maternal and perinatal mortality were reported to be as high as 75% and 85% respectively [5]. More current data showed that with prompt diagnosis and treatment the maternal and perinatal mortality have greatly decreased to approximately 18% and 23% respectively. [6,7]

Early diagnosis of Acute Fatty Liver of Pregnancy (AFLP) sometimes can be difficult as it shares features with other conditions such as pre-eclampsia, viral hepatitis and cholestasis of pregnancy. Careful history and physical examination, along with appropriate laboratory and imaging results, are often sufficient to make the diagnosis, and biopsy of liver is rarely required. Early identification, supportive care and expeditious delivery are the mainstay treatment for Acute fatty liver of pregnancy (AFLP) for maternal and neonatal outcome.

Vaginal delivery is probably the best approach if tolerated, however caesarean delivery is often performed because of rapidly deteriorating maternal and foetal status. During postpartum recovery period haemodynamic monitoring is necessary because patients with acute fatty liver of pregnancy (AFLP) are at high risk of bleeding as a result of coagulopathy. Transfusion of fluids and blood products may be needed. Besides risk of bleeding patients are also at risk of hypoglycaemia and glucose infusion may be needed.

MATERIAL AND METHODS:

Observational prospective study of cases of Acute Fatty Liver of Pregnancy with various presentations and their outcomes are studied over a period of 1 year from December 2022 to November 2023 at obstetrics and gynaecology department at ASRAM medical college, Eluru, Andhra Pradesh. Approval from institutional ethics committee taken. Consent of the patients participated in study taken. During our study total 7 cases were diagnosed as Acute Fatty Liver of Pregnancy.

All the patients were conscious and coherent at the time of presentation. Detailed history of the patients was taken and thorough

clinical examination done. Laboratory investigations like Complete blood picture, liver function tests, renal function tests, coagulation profile, urine microscopy and imaging with ultra sonogram were done. Patients who met the Swansea criteria were included in the study. Patients were subjected to daily liver function tests and coagulation profile to assess prognosis. Medical gastroenterologist opinion was taken during evaluation.

Swansea criteria includes:

1. Vomiting
2. Encephalopathy
3. Polydipsia/polyuria
4. Abdominal pain
5. Elevated bilirubin (over 0.8 mg/dL or 14 micromol/L)
6. Hypoglycemia (less than 72 mg/dL or 4 mmol/L)
7. Leukocytosis (over 11000 cells/microL)
8. Elevated transaminases (AST or ALT) (greater than 42 international unit/L)
9. Elevated ammonia (over 47 micromol/L)
10. Elevated uric acid (above 5.7 mg/dL or 340 micromol/L)
11. Acute kidney injury, or creatinine over 1.7 mg/dL or 150 micromol/L
12. Coagulopathy or prothrombin time greater than 14 seconds
13. Ascites or bright liver on ultrasound scan
14. Microvesicular steatosis on liver biopsy

6 or more of the following criteria are needed in the absence of another known cause of liver dysfunction to establish a diagnosis.

RESULTS:

During our study total 7 cases were diagnosed as Acute Fatty Liver of Pregnancy. 100% had complaints of vomiting and examination revealed jaundice. 86% had epigastric pain, 57% had polydipsia, 43% had pruritus and 43% had Encephalopathy.

Table 1: Clinical Presentations:

SYMPTOM	PATIENT 1	PATIENT 2	PATIENT 3	PATIENT 4	PATIENT 5	PATIENT 6	PATIENT 7
MS	NT 1	NT 2	NT 3	NT 4	NT 5	NT 6	NT 7
Vomiting	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Epigastric pain	Yes	Yes	No	Yes	Yes	Yes	Yes
Jaundice	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Polydipsia	Yes	Yes	No	Yes	No	No	Yes
Pruritus	Yes	No	No	Yes	No	Yes	No
Encephalopathy grade (0-4)	2	0	0	0	3	0	2

Pregnancy outcome:

Age of presentation - <20 years (0%), 21-30 years 7 (100%), >30 years 0 (0%). Primi gravida were 5 (71%), second gravida 2 (29%). Out of 5 primi gravida 1 was twin pregnancy. Period of gestation - <32 weeks 0%, 32-36 weeks - 6 (86%), 37 weeks and above -1(14%). 3 (43%) delivered by Vaginal route and 4 (57%) by LSCS. 1 Intrauterine death was noted. 5 (63%) were male and 3 (27%) were female. Weight of the newborn - < 2 kg -1(13%), 2 - 2.5 kg- 7 (77%), >2.5kg - 0. APGAR Score - 6 (75%) had 7-9 score, 1 (12.5%) had score <7, and 1 (12.5%) had score 0.

Table 2: Pregnancy Outcome:

Pt No	Age	Parity	Gestation al age (weeks)	Days b/w symptoms and delivery	Mode of delive ry	Sex of baby	Fetal weight (kg)	APGAR
1	26	G2P1L1	37+5	3	VD	Male	2.22	7-9
2	24	Primi	36+6	2	VD	Male	2.07	7-9
3	22	Primi with MCDA	36	4	LSCS	Female	2.34 2.15	7-9 7-9
4	28	Primi	34+1with IUD	2	VD	Male	2	0
5	21	G2P1L1	35+6	5	LSCS	Female	1.64	5-7
6	24	Primi	32	3	LSCS	Male	2.01	7-9
7	28	Primi	36	2	LSCS	Male	2.4	7-9

Table 3: Complications:

Findings	Number	Percentage (%)
Disseminated Intravascular Coagulation(DIC)	4	57.1
Ascites	2	28.5
Liver failure	4	57.1
Hepatic encephalopathy	3	42.8
Acute Respiratory Distress Syndrome(ARDS)	3	42.8
Hemorrhagic shock	3	42.8
Pulmonary edema	1	14.2
Sepsis	1	14.2
Metabolic acidosis	2	28.5
Multi organ dysfunction syndrome (MODS)	4	57.1
Impaired glucose tolerance	1	14.2
Hypoglycemia	4	57.1
Maternal death	1	14.2
Intra uterine death (IUD)	1	14.2
Post partum haemorrhage (PPH)	1	14.2
Acute renal insufficiency	7	100

Complications:

Various complications were observed in these patients. 4 (57%) had liver failure with elevated enzymes. Hepatic encephalopathy was seen in 3 (43%), patients. 3 (43%) had metabolic acidosis, 4 (57%) had Multi Organ Dysfunction, 1 (14%) had impaired glucose tolerance, 4 (57%) had hypoglycaemia. 3 (43%) had haemorrhaging shock, 3 (43%) had ARDS, 1 (14%) had pulmonary edema, 1 (14%) landed in sepsis, 4 (57%) had Disseminated Intravascular Coagulation (DIC), 2 (29%) had Ascites, 1 (14%) presented with intrauterine death of foetus, 1 (14%) had postpartum hemorrhage (PPH), 7 (100%) had Acute Renal insufficiency and 1 (14%) maternal death happened in postpartum period due to MODS along with COVID-19 infection on POD-6.

DISCUSSION:

The diagnosis of Acute fatty liver of pregnancy (AFLP) was made on the basis of clinical, laboratory criteria and imaging. All cases exhibited six or more of the Swansea criteria to objectively confirm the diagnosis of AFLP. AFLP is rare, late-gestational, potentially life-threatening complication which occurs at mean gestational age of 36 weeks (range 32-38 weeks).

Zhu et al. [8] showed the 34th gestation week might be important time

for screening Acute fatty liver of pregnancy (AFLP) outpatients. In our study mean gestational age of presentation was 35+/-3 weeks.

Lee et al. and Kilpatrick et al. agree that signs and symptoms commonly associated with Acute fatty liver of pregnancy (AFLP) include nausea/vomiting (75% of patients), epigastric abdominal pain (50% of patients), malaise, anorexia, and jaundice [9]. In our study nausea and jaundice was seen in 100% of patients and 86% had epigastric pain.

According to Sibai et al. early termination of pregnancy in case of associated HELLP syndrome instead of undue prolongation of labor.

In our study 43% delivered by vaginal route and 57% underwent LSCS.

In our study average age of presentation is 24.71+/-5 years. In our study average baby weight is 2.1kg+/-0.46kg. In our study complications seen were Hepatic encephalopathy 3 (43%), 3 (43%) had metabolic acidosis, 4 (57%) had Multi Organ Dysfunction (MODS), 1 (14%) had impaired glucose tolerance, 4 (57%) had hypoglycaemia. 3 (43%) had haemorrhaging shock, 3 (43%) had Acute respiratory distress syndrome (ARDS), 1 (14%) had pulmonary edema, 1 (14%) landed in sepsis, 4 (57%) had Disseminated Intravascular Coagulation (DIC), 2 (29%) had Ascites, 1 (14%) presented with intrauterine death of foetus, 1 (14%) had postpartum hemorrhage (PPH), 7(100%) had Acute Renal insufficiency and 1 (14%) maternal death.

In a study conducted by Nelson et al. Maternal mortality, which was as high as 80% to 90% in the 1980s, has now decreased to less than 10%. Perinatal mortality remains substantial and is now recorded in approximately 20% of AFLP cases. In our study maternal mortality was 1(14.2%) and there was 1(14.2%) Intrauterine death.[10]

Medical management of Acute fatty liver of pregnancy (AFLP) is supportive. Blood sugar levels should be monitored and treated. Initial treatment involves supportive management with intravenous infusion, intravenous glucose, fresh frozen plasma, and packed red blood cells with coagulation dysfunction to reduce blood loss and cryoprecipitate to correct DIC. Once the patient is stabilized, further course of management includes the appropriate mode for delivery.

The rare patient who progresses to fulminant hepatic failure can be treated by liver transplantation. Further pregnancies are often uncomplicated but remain at risk for recurrent AFLP. Other systemic complicating effects include acute respiratory distress syndrome sometimes requiring assisted ventilation, ascites, and upper gastrointestinal bleeding from gastric ulceration, and Mallory-Weiss syndrome. Maternal deaths occurs due to hemorrhage, gastrointestinal bleeding, sepsis, aspiration, pancreatitis, renal failure.

Acute fatty liver of pregnancy also has a detrimental effect on the foetus. One of the complications of acute fatty liver of pregnancy (AFLP) is maternal metabolic acidosis secondary to impaired clearance of serum lactate by damaged hepatocytes.[11] Maternal metabolic acidosis directly affects foetal acid base status. Therefore prompt correction of maternal metabolic acidosis is essential to foetal well being.[12]

CONCLUSION:

Acute fatty liver of pregnancy (AFLP) is an rare, life-threatening complication of third trimester with variable presentation. Rapid progression is unpredictable.

The patients, who are critically ill at the time of clinical presentation, develop complications, or continue to deteriorate despite emergency delivery, require management in the intensive care unit (ICU).

Early diagnosis, prompt delivery, adequate supportive care, and a multidisciplinary approach are the key to a good outcome.

REFERENCES:

1. Knight M, Nelson-Piercy C, Kurinczuk JJ, Spark P, Brocklehurst P, UK Obstetric Surveillance System. A prospective national study of acute fatty liver of pregnancy in the UK. Gut. 2008 Jul;57(7):951-6. [PubMed]
2. Sheehan HL. The pathology of acute yellow atrophy and delayed chloroform poisoning. J Obstet Gynaecol Br Emp. 1940;47:49-62
3. Ibdah JA, Bennett MJ, Rinaldo P, Zhao Y, Gibson B, Sims HF, Strauss AW. A fetal fatty-acid oxidation disorder as a cause of liver disease in pregnant women. N Engl J Med.

- 1999 Jun 03;340(22):1723-31. [PubMed]
4. Watson WJ, Seeds JW. Acute fatty liver of pregnancy. *Obstet Gynecol Surv.* 1990;45:585
 5. Varner M, Rinderknecht NK. Acute fatty metamorphosis of pregnancy. A maternal mortality and literature review. *J Reprod Med.* 1980;24:177-80
 6. Kaplan MM. Acute fatty liver of pregnancy. *N Engl J Med.* 1985;313:367-70
 7. Knox TA, Olans LB. Liver disease in pregnancy. *N Engl J Med.* 1996;335:569-76
 8. Wei Q, Zhang L, Liu X. Clinical diagnosis and treatment of acute fatty liver of pregnancy: a literature review and 11 new cases. *J ObstetGynaecol Res.* 2010;36(4):751-6.
 9. <https://radiopaedia.org/articles/acute-fatty-liver-of-pregnancy>
 10. Nelson DB, Byrne JJ, Cunningham FG. Acute fatty liver of pregnancy. *Obstet Gynecol.* 2021;137(3):535-546. doi:10.1097/AOG.0000000000004289
 11. Woll PJ, Record CO. Lactate elimination in man: Effects of lactate concentration and hepatic dysfunction. *Eur J Clin Invest.* 1979;9:397-404
 12. Werner CJ, Zoller DP, Baskin WN, Eichmann RE. Acute fatty liver of pregnancy associated with maternal and fetal metabolic acidosis. *J Fam Pract.* 1988;26:198-9