



HYPOKALEMIC PERIODIC PARALYSIS AS INITIAL MANIFESTATION OF PRIMARY SJOGRENS SYNDROME : A CASE REPORT

Medicine

Dr. Jagdish K. Mohale	Junior Resident Dept Of General Medicine Dr. Vaishampayan Memorial Government Medical College Solapur
Dr. Shubhangi V. Dhadke	Associate Professor Dept Of General Medicine Dr. Vaishampayan Memorial Government Medical College Solapur
Dr. Vithal N. Dhadke	Professor And Head Of Dept Of General Medicine Dr. Vaishampayan Memorial Government Medical College Solapur
Dr. Kalyani S. Chavan	Junior Resident Dept Of General Medicine Dr. Vaishampayan Memorial Government Medical College Solapur
Dr. Pranali S. Pol	Junior Resident Dept Of General Medicine Dr. Vaishampayan Memorial Government Medical College Solapur

ABSTRACT

Sjogrens syndrome (SS) is common autoimmune diseases manifesting with various glandular and extra glandular features which are difficult to evaluate and manage. Hypokalemic flaccid paralysis is the rare extra glandular clinical manifestation of SS. We present a case of acute flaccid paralysis which on evaluation turned out to be primary sjogrens syndrome. We present a case of 34 year old young female with no comorbidities presented with acute flaccid paralysis. Her evaluation reveals severe hypokalemia with K^+ - 1.7, metabolic acidosis with pH - 7.23, urinary pH - > 6.5. An initial diagnosis of hypokalemic periodic paralysis secondary to distal renal tubular acidosis (dRTA) is made. She was later found to have sjogrens syndrome on the basis of positive ANA and Anti Ro (SS-A) antibody. Also Ultrasound abdomen reveals nephrocalcinosis which is commonly associated with dRTA. She was managed with potassium supplements, sodium bicarbonate and immunomodulators. Primary sjogrens syndrome usually associated with distal renal tubular acidosis which may cause hypokalemia leading to HPP, rarely which may be the initial presenting symptom in young women as in our case. Early diagnosis and treatment of these patients would lead to less complications and better quality of life.

KEYWORDS

Sjogrens syndrome, hypokalemic paralysis, distal renal tubular acidosis, autoimmune disease, young female

INTRODUCTION :-

Metabolic and electrolyte abnormalities leads to several types of periodic paralysis with first ever description in 1727 by Musgrave.^{1,2} Hypokalemic periodic paralysis (HPP) is commonest of all metabolic myopathies with prevalence of 1 in 100000.³ It is characterized by acute flaccid paralysis ranging from transient muscular weakness to potentially fatal severe respiratory muscle paralysis and cardiac arrhythmias. HPP is manifested by varying group of disorders mainly thyroid disorders such as thyrotoxicosis, renal tubular dysfunctions such as distal renal tubular acidosis (dRTA), Gittelman's syndrome and non-renal loss of potassium in the form of diarrhoea, recurrent vomiting and excessive sweating. Distal RTA may be inherited or acquired with autoimmune disorders such as Sjogren's syndrome, Systemic lupus erythematosus and rheumatoid arthritis were main acquired disorders.

Sjogren's syndrome is chronic autoimmune connective tissue disorder characterized by lymphocytic infiltration of the secretory exocrine glands mainly salivary and lacrimal glands.³ It may present with extra glandular manifestations in form of renal, respiratory, neurologic, hepatic, cutaneous and vascular involvement. Sicca syndrome which is combination of dryness of oral cavity, pharynx, larynx, eyes and vagina without any association with other autoimmune connective tissue disorder is called as primary Sjogren's syndrome (pSS). Association of sicca syndrome with various rheumatic disorders such as rheumatoid arthritis, systemic lupus erythematosus and systemic sclerosis is called as secondary Sjogren's syndrome (sSS).⁴ Rheumatoid arthritis is most commonly associated with sSS with prevalence of 20%.^{5,6} SS presents as hypokalemic periodic paralysis as initial presentation which was first described in 1981.⁷ We report a case of pSS presented with hypokalemic periodic paralysis.

Case History :-

A 34 year old female was brought to emergency with complaints of acute onset, progressive quadriparesis of 2 days duration with involvement of neck and trunk. Muscle cramps, dysphagia and difficulty in breathing were also present. She had no significant past history. On evaluation she was afebrile, blood pressure 130/80 mmhg, pulse rate 90/min and respiratory rate 19/min. Her single breath count

was 16 sec. On nervous system examination she had decreased muscle tone, proximal and distal muscle strength 2/5 in all four limbs with diminished reflexes. A provisional diagnosis of acute flaccid paralysis was made and further laboratory investigations reveals severe hypokalemia of 1.7 meq/L, metabolic acidosis with pH- 7.23, pCO_2 - 18.1 mmhg, HCO_3^- - 11.4 mmol/L and creatine phosphokinase with in normal range. ECG shows prominent U waves, Hemogram reveals Hb- 15.5 gm%, total leukocyte count 7800/mm³, platelet count 202000/mm³, while liver and renal functions were in normal range. Urine analysis reveals pH- 6.5, Na^+ - 80 meq/l and K^+ - 23.2 meq/l and thyroid functions reveals hypothyroidism with t_3 - 0.528 ng/ml (1.3-3.1), t_4 - 1.94 ug/dl (5.13 - 14.03) tsh - 69.14 uIU/ml (0.27-4.2). Imaging studies with MRI brain and cervical spine were normal. Ultrasound abdomen reveals nephrocalcinosis in both kidneys. The initial evaluation was suggestive of distal renal tubular acidosis. She was managed with intravenous potassium chloride, sodium bicarbonate and supportive care.

Her condition improved with treatment and further evaluation revealed significant positive anti-nuclear antibody anti Ro 60 (SSA) 175 RU/ml (>25 positive), anti Ro 52KD 200 RU/ml (>25 positive), normal C3 and C4 and negative anti La (SS-B) antibody. Based on Positive ANA and anti Ro (SS-A) antibody with distal RTA diagnosis of primary sjogrens syndrome was made. She was started on Deflazacort and Hydroxychloroquin (HCQ). Her condition improved with treatment and she was discharged on potassium citrate solution, sodium bicarbonate and HCQ. She was on regular follow up and remains asymptomatic.

DISCUSSION :-

Sjogren's syndrome presenting as HPP was first reported in 1981;⁷ after that till now only few cases were reported and in India first case was reported in 1996.⁸ Kulkarni et al had done cross sectional retrospective study from database maintained at Centre for Rheumatic Diseases, Pune since 1996 with records of over 50000 patients and found a total 16 (0.032%) patient of pSS with initial presentation with HPP.⁹ They found younger age of onset, female dominance, ANA and SSA positivity in these patients of pSS with HPP which is similar to other case series studies as well as in our case.¹⁰ Hence early diagnosis

of this small subset of patient is required for better management.

Our patient presented with acute progressive flaccid paralysis. She was having no previous comorbidities and normal sensory examination therefore AIDP and HPP were considered after ruling out infective and traumatic etiologies. The diagnosis of HPP was made due to severe hypokalemia and prompt response to potassium supplements. Diagnosis of dRTA was considered due to metabolic acidosis with urinary $\text{pH} \geq 5.5$. Distal RTA is mainly of primary (Inherited) and secondary (acquired) types. Primary dRTA may be inherited as autosomal dominant or autosomal recessive forms and involves SLC4A1, ATP6V0A4 or ATP6V1B1 genes which are responsible for managing levels of acid base in the body.¹¹ Acquired dRTA mainly caused by autoimmune disorders (Sjogren's syndrome, Systemic lupus erythematosus or Rheumatoid arthritis), calcium disorders (primary hyperparathyroidism, idiopathic hypercalciuria with nephrocalcinosis), dysproteinemic syndromes, drugs and toxins.¹¹

Patient with dRTA usually present with normo-or hypokalemia. ACR/EULAR developed classification criteria for pSS in 2016 which applies to the individuals that have a score of ≥ 4 .¹² Our patient was diagnosed the pSS with total score of 4 with Schirmer's test $< 5 \text{ mm} / 5$ minutes in both eyes and positive Anti-Ro (SS-A) Antibody. In pSS renal involvement is well documented and two main forms are glomerulonephritis and interstitial nephritis. About one third of the patient are associated with interstitial nephritis which manifest mainly as complete or latent distal renal tubular acidosis and proximal RTA rarely.¹³ Distal RTA is associated with hypokalemia which is caused by decreased distal tubular Na delivery, defective H-K ATPase, bicarbonaturia and secondary hyperaldosteronism.^{11,14}

In primary Sjogren's syndrome HPP may occur three months to four years prior to development of sicca syndrome.^{15,16} Systemic (extraglandular) manifestations such as HPP with dRTA of Sjogren's syndrome are managed by immunomodulatory therapy such as hydroxyl chloroquine (HCQ), mycophenolate mofetil, corticosteroids, cyclophosphamide and biologic therapies such as rituximab. Our patient was initially managed with tab Deflazacort 6 mg and tab HCQ 400 mg once a day. Later steroid was tapered off while HCQ continued and she remains asymptomatic till date.

CONCLUSION

We report this case to emphasize that young women presenting with HPP should be screened for dRTA and other extra glandular features with clinical suspicion for pSS for early diagnosis and initiation of appropriate treatment.

REFERENCES

- Kayal A, Goswami M, Jain R, Das M. Clinical and biochemical spectrum of hypokalemic paralysis in North: East India. *Ann Indian Acad Neurol*. 2013;16(2):211–7. doi:10.4103/0972-2327.112469.
- Fontaine B, Vale-Santos J, Jurkat-Rott K, Reboul J, Plassart E, Rime CS, et al. Mapping of the hypokalaemic periodic paralysis (HypoPP) locus to chromosome 1q31–32 in three European families. *Nat Genet*. 1994;6(3):267–72. doi:10.1038/ng0394-267.
- Pillemer SR, Matteson EL, Jacobsson L, Martens PB, Fox P, Michael W, et al. Incidence of physician-diagnosed primary Sjögren syndrome in residents of Olmsted County. *Mayo Clin Proc*. 2001;76(6):593–9.
- Morrow J, Nelson JL, Watts RA, Isenberg DA. Sjogren's syndrome in Autoimmune Rheumatic Diseases. In: 2nd Edn.. vol. 6. Oxford Medical Publication; 1999. p. 147–69.
- Carmona L, Gonzalez AI, Balsa A, Angel BM, Tena X, Sanmarti R, et al. Rheumatoid arthritis in Spain: occurrence of extra-articular manifestations and estimates of disease severity. *Ann Rheum Dis*. 2003;62:897–900.
- Skoumal M, Wottawa A. Long-term Observation Study of Austrian Patients with Rheumatoid Arthritis. *Acta Med Austriaca*. 2002;29(2):52–6. doi:10.1046/j.1563-2571.2002.01043.x.
- Raskin RJ, Tesar JT, Lawless OJ. Hypokalemic periodic paralysis in Sjögren's syndrome. *Arch Intern Med*. 1981;141:1671–3.
- Thomas N, Ramakrishna B, Seshadri MS. Hypokalemic periodic paralysis: an unusual cause. *J Assoc Physicians India*. 1996;44:207–8.
- Kulkarni N, Chopra A. Clinical Profile of Primary Sjogren's syndrome with Hypokalemic Periodic Paralysis. *J Assoc Physicians India*. 2018;66:69–70.
- Jain VK, Antony PT, Mithun CB, Singh BK, Mehra S, Usdadiya J, et al. AB0559Hypokalemic Paralysis in Primary Sjogren Syndrome: is IT A Distinct Clinical Subset Which Requires Lifelong Potassium Supplementation Rather than Immunosuppressants? *Ann Rheum Dis*. 2014;73(2):991. doi:10.1136/annrheumdis-2014-eular.1739.
- Soriano JR. Renal Tubular Acidosis: The Clinical Entity. *J Am Soc Nephrol*. 2002;13(8):2160–70. doi:10.1097/01.asn.0000023430.92674.e5.
- Shiboski CH, Shiboski SC, Seror RF, Criswell LA, Labetoulle M. American College of Rheumatology/European League Against Rheumatism classification criteria for primary Sjogren's syndrome A consensus and data-driven methodology involving three international patient cohorts. *Ann Rheum Dis*. 2016;76(1):9–16.
- Goules AV, Tatouli IP, Moutsopoulos HM, Tzioufas AG. Clinically Significant Renal Involvement in Primary Sjögren's Syndrome: Clinical Presentation and Outcome. *Arthritis Rheumatism*. 2013;65(11):2945–53. doi:10.1002/art.38100.
- Lin SH, Cheema-Dhadli S, Gowrishankar M, Marliiss EB, Kamel KS, Halperin ML, et al. Control of excretion of potassium: lessons from studies during prolonged total fasting in human subjects. *Am J Physiology-Renal Physiol*. 1997;273(5):F796–F800. doi:10.1152/ajprenal.1997.273.5.f796.

- Shioji R, Furuyama T, Onodera S, Saito H, Ito H, Sasaki Y, et al. Sjögren's syndrome and renal tubular acidosis. *Am J Med*. 1970;48(4):456–63. doi:10.1016/0002-9343(70)90045-8.
- Pun KK, Wong CK, Tsui EY, Tam SC, Kung AW, Wang CC, et al. Hypokalemic periodic paralysis due to the Sjogren syndrome in Chinese patients. *Ann Intern Med*. 1989;110(5):405–6.