



ROLE OF HIGH RESOLUTION ULTRASOUND IN TRAUMATIC PERIPHERAL NEUROPATHY IN UPPER LIMBS

Radiology

Dr. Kamireddy Dheeraj Reddy*	Junior Resident Department Of Radiodiagnosis Vydehi Institute Of Medical Sciences & Research Centre, Bengaluru, Karnataka, 560066 *Corresponding Author
Dr. Madan Mohan Babu	Professor Department Of Radiodiagnosis Vydehi Institute Of Medical Sciences & Research Centre, Bengaluru, Karnataka, 560066
Dr. Suresh A	Professor & Hod Department Of Radiodiagnosis Vydehi Institute Of Medical Sciences & Research Centre, Bengaluru, Karnataka, 560066
Dr. Thummala Srimadhav	Junior Resident Department Of Radiodiagnosis Vydehi Institute Of Medical Sciences & Research Centre, Bengaluru, Karnataka, 560066

ABSTRACT

Aim of work: The type of traumatic peripheral nerve injury is a key factor for determining optimal treatment. Proper assessment of peripheral nerve injury facilitates appropriate treatment, significantly affects prognosis, and reduces disabilities. This study evaluated ultrasonography (US) to assess upper limb traumatic nerve injuries and compared US with electrodiagnostic studies as the gold standard. **Materials and Methods:** Participants were 30 adults (24[80%] men, 06[20%] women; mean age 36.3 ± 13.5 years) with a total of 42 peripheral nerve injuries (duration of 1 month–3 years). High-frequency US examinations and electro-physiologic studies confirmed upper limb peripheral nerve injury. **Results:** Nerve discontinuation was diagnosed in 07 (16.6%) nerves; the cross-sectional area was increased in 14 (33.3%) nerves. Of the 42 injuries, 24 (57.1%) were median, 10 (23.8%) were ulnar, and 8 (19.1%) were radial nerves. No statistically significant difference was found between US and electro-physiologic studies for nerve injury diagnosis ($p = 0.054$). **Conclusion:** No significant differences were found between US and electro-physiologic studies for diagnosis of nerve injuries; however, US was valuable to assess surrounding tissue and supplied muscles. The capabilities to detect nerve injury and associated distal muscular, vascular, and other regional structures position the US as a complementary diagnostic tool.

KEYWORDS

Ultrasound, Peripheral nerve, Trauma

INTRODUCTION

Peripheral nervous system trauma is rare, but this injury can have a significant negative influence on a patient's health and functional capabilities. The overall incidence is 2–3% of patients who visit major trauma centers worldwide, or 13–23 per 100,000 people annually. The most frequent cause of peripheral nerve injuries (PNIs) during calm times is motor vehicle collisions, whereas less prevalent causes are penetrating trauma, falls and industrial accidents^[1]. The risk of PNI increases to about 5% when plexus and root injuries are taken into account. Iatrogenic injury is also common. For example, cervical lymph node excisions have a 3–6% risk of PNI to the axillary nerve, and radial nerve injury from humeral shaft osteosynthesis for fracture stabilization has a 5–15% risk of PNI, in addition to the inherent risk of the fracture^[1].

Specific mechanisms of trauma increase the risk of specific type of nerve injury. To determine which nerves may be impacted, knowledge of the precise trauma mechanism is necessary. This knowledge should take precedence over the conventional neurologic methodology when localizing the peripheral nervous system disease during workup^[1]. Most PNIs in the upper limb are the ulnar, median, and radial nerves, whereas PNIs in the lower limb are most often the sciatic nerve, followed by the peroneal nerve, and, less frequently, the tibial and femoral nerves^[2].

Despite the fact that localization of PNIs is clinically possible accurate localization of the injury is challenging, especially when many nerves are involved. Electrophysiologic studies are useful to detect the injury severity and location. However, because the most accurate diagnostic information from these studies is only accessible after 2 weeks when wallerian degeneration is fully developed; electro-diagnostic techniques are not well suited for the acute phase and are inherently painful. In addition, in individuals with a total loss of nerve conduction, electro-diagnostic testing cannot reveal the precise location of the lesion site. Furthermore, these studies cannot display the morphologic changes of the injured nerve^[3,4].

Compared with electro-diagnostic tests, ultrasonography (US) can accurately and painlessly provide this information. Within the whole clinical workup for PNI, ultrasound has a complementary and expanding role. It not only offers a quick and affordable method to probe extensive stretches of peripheral nerves, but it also has unique

benefits based on its dynamic, real-time nature and its small number of clinical limitations and contraindications. Thanks to developments in high-frequency transducers, the function of the US is expanding^[5].

When using the US, it is essential to determine the following factors: whether the nerve is still continuous; the size and position of any gaps in the nerve course; the presence of any focal neuromas or extra nerve damage sites, such as tandem lesions; the presence of any foreign bodies; the degree of neighboring scar tissue; and the condition of surrounding tissues and structures, such as tendons, arteries, and bones^[4]. As an advantageous procedure in the evaluation of peripheral entrapment syndromes, US can determine if thenerves are compressed, tethered, or hypermobile with respect to adjacent structures. Shifting transducer pressure may have an impact on sonopalpation, which can be used to demonstrate that the patient's condition is neuropathic pain, find the location of the peripheral nerves responsible for the symptoms, and identify the exact site that consistently triggers the patient's symptoms, often identifying the responsible basic pathology. Because the whole longitudinal course nerve is typically evaluated in a single examination, it is possible to detect multi-segmental disease with cost-effectiveness^[5]. The current study will evaluate the value of the US in the assessment of traumatic nerve injuries of the upper limb compared with the role of electro-diagnostic studies as the current gold standard diagnostic modality.

MATERIALS AND METHODS

From December 2022 to February 2024, adult patients with a history of PNIs were recruited from the clinical neurology and orthopedic units in the hospital after undergoing physical examination by an orthopedic surgeon sub-specialized with 10 years of experience. The study participants were 30 patients with 42 nerve injuries, which included 24 (80%) men and 06 (20%) women with a mean age of 36.3 ± 13.5 years. All of the study participants gave informed written consent to the study.

Inclusion criteria were a clinical picture of PNI in the upper limb, with a duration of 1 month up to 3 years; and surgical exploration of the PNI. -Exclusion criteria were diabetes mellitus, polyneuropathies, renal or hepatic disorders, or radiculopathy (clinical or electrophysiologic) in the segment supplying the injured nerve; and refusal to participate in the study. All patients underwent thorough history-taking and clinical examination. These patient history data were collected: date of injury

date; onset of neurologic manifestations, such as weakness, numbness, and loss of sensation; and history of surgical intervention for the injury. Clinical examinations

The following clinical examinations were performed for all patients.

Electrophysiological studies

The electrophysiological research was performed using the Neuropak MEB-9200G/K EP/EMG measurement device (Neuropak M1, version 08.1, Nihon Kohden, Tokyo, Japan). Standard motor nerve conduction studies (NCS) were performed to record the nerve damage for these sites and data: abductor pollicis brevis; abductor digiti minimi; wrist; below and above the elbow stimulation for the ulnar nerve, extensor indices properties; and forearm and elbow stimulation for the radial nerve. The second digit (antidromic stimulation at the wrist for the median nerve), the fifth digit (antidromic stimulation at the wrist for the ulnar nerve), and the anatomical snuff box were recorded as part of routine sensory NCS (antidromic stimulation at the wrist for the superficial radial nerve). Sonography

Conventional high-resolution US in the B- Mode was performed for all patients using Philips Affinity 30 with a linear probe of 5-10 MHz frequency. The US examinations were performed by a radiologist and who was blinded to the results of the electrodiagnostic tests and clinical Examinations. Depending on the patient's overall condition, patients were either examined while lying flat on their back or comfortably sitting facing the radiologist. The elbow was supported and flexed, and the forearm was extended when the affected nerve was examined at the forearm. To examine the hand, the wrist was held steady and positioned slightly hyper extended. A generous amount of gel was used to improve contact between the probe and the skin's surface; in cases of painful neuromas, little pressure was used. Using sterile gel and taking proper antiseptic precautions, such as using disposable non-sterile gloves and cleaning the US machine and transducer before and after scanning is advised for susceptible patients or while scanning in wound areas.

To compare the healthy and injured nerves, a US examination was also performed on the healthy unaffected side. Using a recognized anatomic landmark, the nerves were identified sonographically. Once identified, the nerve was tracked proximally and distally while remaining in the center of the US image along its short axis. The area of interest was then positioned properly focused during US inspection using transverse and longitudinal scanning planes.

Image interpretation

To interpret the images, the following steps were taken and factors were assessed:

1. Identification of the injured nerve lesion site by elicitation of the Tinel sign during US examination because it can help identify the nerve lesion site while using sonopalpation, or the compression of the US probe.
2. Testing for nerve continuity, which includes determining whether the nerve's fascicle is swollen and hypoechoic, whether it lacks a typical fascicular pattern, and whether a neuronal stump is present.
3. Degree of nerve injury, determined to be partial or complete based on the Sunderland classification of PNI as follows [6]:
 - **Sunderland grade 1:** No pathologic findings on the US are visualized or only mild swelling of the nerve with an intact fascicular pattern.
 - **Sunderland grade 2:** Enlarged cross-sectional area (CSA) of affected fascicles and nerves at the lesion site because of axonal swelling and edema are evident.
 - **Sunderland grades 3 and 4:** Lesion shows a loss of the normal nerve architecture and echogenicity with a disruption of the fascicular pattern and often sizable hypoechoic enlargement of the nerve.
 - **Sunderland grade 5:** Lesion shows transection of the nerve with a loss of nerve continuity.
4. Amount of nerve retraction and the gap after complete nerve transection.
5. Nerve thickening, by measuring the CSA of the injured nerve within the hyperechoic perineurial rim of the nerve and the transducer perpendicular to the nerve.
6. Bony fragments pressing against the nerve near the site of the bone fracture.
7. Proximal terminal neuroma, characterized by uniform texture,

hypoechoic echogenicity, and concentric swelling at the terminal end of a transected nerve, with measurement of its size by placing several diameter markers proximal to, at, and caudal to the lesion site.

8. Neuroma in-continuity, by determining if it is intact and manifests as a broadening of the nerve's contour in a nodular shape.
9. Perineural fibrosis, vascular injury, and retained foreign bodies, by evaluating for their presence.
10. Muscular injury, recognized in this study as an increase in echogenicity of the supplied muscle and/or reduced girth (atrophic changes).
11. Doppler brightness mode with color or power Doppler to quantify blood flow for both intraneural and epineurial compartments.

Statistical analysis

Statistical analysis was conducted using SPSS (version 22, IBM, Armonk, NY, USA). Quantitative variables were presented as mean $\pm$ standard deviation and range, and qualitative variables were presented as frequency and percentages. Paired comparison of diagnostic tests was conducted using the McNamara test, comparison between proportions was conducted using the chi-squared test. Sensitivity, specificity, positive predictive value (PPV), negative predictive value (NPP), and overall diagnostic accuracy were calculated using a 2 $\times$ 2 contingency table. Any value of $p < 0.05$ was considered significant.

RESULTS

A total of 30 patients with 42 nerve injuries were eligible for inclusion in the final analysis, with 24 (80%) men and 06 (20%) women. The mean age was 36.3 ± 13.5 years. Some patients had more than one injured nerve.

The mode of trauma was crush injury (13 cases), cut wounds (13 cases), compression (3 cases), and traction (1 case). Of the 42 injured nerves, 13 (30.96%) were on the right side and 29 (69.04%) were on the left side. The injury level was at the arm in 16 nerves, followed by the forearm (14 nerves), and 10 injuries were at the hand. Injured nerves were mainly median nerve in 08 (57.1%) cases, ulnar in 10 (23.8%) cases, and radial nerve in 8 (19.1%) cases. Table 1 presents these data. As shown in Table 2, US findings showed 12 (28.5%) intact nerves, neuroma in 09 (21.4%) nerves (Figs. A, G, D) nerve discontinuation in 7 (16.6%) (Fig. B) nerves and an increase of CSA of the injured nerve in 14 nerves (33.3%) (Figs. C and E).



Fig A

A) 40-year-old male with a history of left humeral fracture with internal fixation with plate and screws since July 2021 with a weak extension of the elbow. A Digital radiography image of the left arm showing internal fixation with plate and screws. B) B-mode image demonstrating, hypo-echogenicity along the course of the radial nerve within the spiral groove intimately related to applied screws. C) B-mode image, transverse axis showing, thickened left radial nerve in the spiral groove of the humerus, suggested axonal injury proved by NCS. D) B-mode image, transverse axis of the right normal radial nerve within the spiral groove of the humerus. E) B-mode image, turbid collection noted the lateral aspect of the axilla.






Fig B


A 33-year-old female patient presenting with history of right medial epicondyle fracture which was treated by applying a cast in June 2021. After removal of the cast, anterior transposition and neurolysis of the ulnar nerve was done. A) B-mode image, transverse axis, shows thickened right ulnar nerve, CSA 19mm², just proximal to the medial epicondyle with intimately related hypoechoic scar tissue and basilic vein. B) B-mode image, longitudinal axis, shows focal interruption of the epineurium of the right ulnar nerve (arrows) with related hypoechoic scar tissue. C) B-mode image, transverse axis, shows thickened right ulnar nerve (superior arrow), at the level of the medial epicondyle with related hypoechoic synovial thickening containing echogenic foci of bone fragments (inferior arrow). D) B-mode image, longitudinal axis, shows indented right ulnar nerve at the level of the medial epicondyle with related hypoechoic synovial thickening containing echogenic foci of bone fragments.

**Fig C**


A 40-year-old male patient presenting with history of cut wound of the right arm just proximal to the elbow crease. He gave history of operative repair of brachial artery injury. A) B-mode image, transverse axis, shows abnormal architecture of the right median nerve at the distal arm with loss of the normal honeycomb appearance and related hypoechoic scar tissue formation. B) B-mode image, longitudinal axis, shows discontinuous right median nerve at the distal arm with related hypoechoic scar tissue formation. C) B-mode image, transverse axis, shows abnormal architecture of the right median nerve (N) at the distal arm with loss of the normal honeycomb appearance and related hypoechoic scar tissue formation, aneurysmal dilatation of the brachial artery (A) and distorted biceps muscle (BICEPS).

**Fig D**

A 26-year-old female with a history of right distal radial fracture in July 2023 treated. After the cast removal, numbness and tingling along the radial aspect of the palm with loss of sensation of the middle finger and newly developed progressive increased thenar swelling. A) Patient's hand with prominent swelling of the thenar eminence and mid-palmar incision scar. B) B-mode image, transverse axis, reveals loss of normal sonographic appearance of the left median nerve at the carpal tunnel with related hypoechoic superficial scar tissue, suggesting partial nerve injury. C) Color Doppler mode image reveals pseudo-aneurysm; Pepsi sign with an area of luminal thrombosis. D) The B-mode image of the thenar muscles, increased echogenicity of the left thenar muscles as compared to the right side with rather preserved girth. E) Intraoperative image revealed complete injury of two of the sensory branches of the median nerve.

**Fig E**

A 25-year-old male patient presenting with history of left wrist compression/degloving injury in 2020. He underwent tendon repair surgery. A) Patient's hand. B) B-mode image, transverse axis, reveals thick hypoechoic swelling of the left median nerve with loss of fascicular architecture. C) B-mode image, longitudinal axis, thickened, hypoechoic focal area with loss of fascicular architecture of the median nerve at the site of injury. D) B-mode image of the thenar muscles, reveal loss of normal sonographic appearance of the left side muscles with increased echogenicity as compared to the right side. E) B-mode image, longitudinal axis, shows discontinuous, disorganized left ulnar nerve at the distal forearm with related hypoechoic scar tissue formation. F) B-mode image of the hypothenar muscles, reveal loss of normal sonographic appearance of the left side muscles with increased echogenicity as compared to the right side.

**Fig F**

An 18-year-old male with a history of a cut wound of the right forearm since September 2021 and tendon repair, sensory loss of the skin on the radial aspect of the palm with skin ulceration. A) B-mode image shows a discontinuous right median nerve at the mid-forearm with related hypoechoic tissue suggesting complete nerve injury proved by NCS.

B) B-mode image shows a transverse axis view at the site of the lesion showing a hypoechoic area with non-visualized nerve tissue. C) B-mode image shows a comparison between the right and left mid-forearm muscles. The right side shows an abnormal architecture of the FDS and FDP muscles with increased echotexture as compared to the left normal side. D) B-mode image shows a comparison between the right (left image) and left (right image) thenar muscles. The right side mildly increased echotexture as compared to the left normal side with rather preserved girth.




Fig G

A 28-year-old male patient presenting with history gunshot injury to the left forearm since June 2021. He underwent flexor muscles and tendons repair in addition to median nerve repair.

A) Patient's forearm showing scar tissue. B) X-ray image of the left forearm multiple retained bullets. C, D) B-mode image, transverse axis, shows thickened median nerve (arrow) at proximal forearm with circumferential hypoechoic lesion. E) B-mode image, longitudinal axis, shows intact epineurium of the median nerve (arrows) at proximal forearm with superficially related hypoechoic lesion (oblique arrow). F) B-mode image, transverse axis, shows thickened hypoechoic ulnar nerve at proximal forearm.

Table 1

MODE	COUNT	PERCENTAGE
CUT WOUND	13	30.95%
CRUSH	13	30.95%
TRACTION	1	2.38
COMPRESSION	3	7.14%
SIDE		
LEFT	29	69.04%
RIGHT	13	30.06%
SITE		
ARM	16	38.09%
FOREARM	14	33.3%
HAND	12	28.5%
NERVE		
ULNAR	10	23.8%
MEDIAN	24	57.1%
RADIAL	8	19.04%

Table 2

US FINDINGS	COUNT	PERCENTAGE
Intact	12	28.57%
Neuroma	9	21.42%
Nerve discontinuation	7	16.66%
Increased CSA	14	33.33%

Table 3 - Usg Features In Muscles

ECHOGENICITY	COUNT	PERCENTAGE
Normal	7	16.66
Increased	35	83.44
MUSCLE GIRTH		
Normal	12	28.5%
Reduced	30	71.5%

Table 4 - Associated Usg Findings

	COUNT	PERCENTAGE
FIBROSIS		
Absent	21	50%
Present	21	50%
VASCULAR		
Absent	38	90.4%
Present	4	9.6%
ORTHOPAEDIC		
Absent	26	61.9%
Present	16	38.1%
OTHERS		
Absent	36	85.71%
Present	6	14.29%
MUSCLE		
Absent	25	59.5%
Present	17	40.5%

Table 5 - Eps Among Included Patients

SEVERITY	COUNT	PERCENTAGE
Intact	2	4.7%
Mild	7	16.6%
Moderate	20	47.6%
Severe	8	19.04%
Complete injury	5	11.9%
Axonal injury	42	100%

DISCUSSION

Injury to the peripheral nerves causes significant disability to patients around the world^[1]. The common causes are motor vehicle collisions, falls, industrial accidents, domestic accidents and penetrating trauma^[7]. Electrodiagnostic investigations are the gold standard for diagnosis of traumatic PNIs. These studies can be used to locate lesions, diagnose the underlying pathology, assess the extent and severity of axonal degeneration, and monitor recovery^[8]. However, electrodiagnosis has a limited capacity to pinpoint the exact site and degree of nerve damage in certain situations, notably in the early post-injury period, because some electrodiagnostic alterations take time to manifest^[9]. These studies also are restricted in their capability to detect morphological changes associated with a specific type of nerve injury^[5].

In comparison to electrodiagnostic examinations, the US can reliably offer this information in a painless manner. In the overall clinical workup, the US plays a complementary, growing role. Not only does it provide a cost-effective and time-efficient method of interrogating long segments of peripheral nerves, but it also has special advantages in terms of its dynamic, real-time nature, as well as a limited number of clinical contraindications and limitations. The role of the US is expanding thanks to advancements in high-frequency transducers and post-processing techniques^[5].

The present study aimed to evaluate the role of high-frequency US in the diagnosis of traumatic PNIs and to compare US findings with the electrodiagnostic findings. Therefore, we conducted a cross-sectional study that included 30 patients with 42 nerve injuries. These cases include 24 (80%) men and 6 (20%) women, with a mean age of 36.3 ± 13.5 years old. These findings are consistent with the evidence in the literature from Ciaramitaroi^[10] and Bösch^[11], which found that PNIs more often occur in male versus female individuals, and the most common age group to be affected by PNIs is patients aged 30–40 years.

Kamble^[12] reported that among 1–2% of individuals with PNIs associated with central nervous system damage, 60% of cases were spinal injuries, fractures, and dislocation of adjacent bones. About 10–34% of patients hospitalized in the rehabilitation center have related nerve damage. Also, early diagnosis and care are vital to enhancing the functional prognosis in these individuals; consequently, it is necessary to identify the related nerve damage.

A study by Martins^[13] showed that the degree and type of nerve injury involved in traumatic neuropathy must be determined because therapy for different degrees and types of traumatic neuropathy may vary. In cases of severe nerve damage (Sunderland classification group 5, neurotmesis), surgical intervention is essential, and it may also be required in cases with Sunderland groups 3 and 4. In the current study, the most common cause of PNI was crush injury and cut wounds in 54 of 69 cases, followed by the other types of injuries. In contrast, Uzun [8] and Elfayoumy^[14] reported that cut injury by sharp objects was the most common cause of PNI. The site of laterality of the injured nerves in our study was more often on the left side (71.9%). Ferrante^[15] reported that the most frequent nerve injury in the upper limbs is the radial nerve, followed by the median and ulnar nerves, whereas our study showed the median nerve was the nerve most often injured (56.3%), followed by the ulnar (25%) and radial (18.8%) nerves.

Koenig^[16] divided the nerve injury findings in the US into five groups: 1) normal, 2) epineurial fibrosis, 3) intraneural fibrosis, 4) neuroma/partial neuroma, and 5) transection^[17] examined 36 patients with nerve injuries in the upper limbs and showed the excellent capability of US in determining the type of injury and detection of proximal and distal nerve stump, foreign particles, stump neuroma, and perilesional excessive scar tissue formation. In our study, neuroma was detected in 21 (21.9%) nerves, and 33 nerves (34.4%) showed an increase in CSA of the injured nerve. In addition, 48 patients had associated fibrosis. These findings almost paralleled those of the study by Elfayoumy^[14], who divided high-resolution US findings into three groups: fusiform in shape (increased CSA) (36.7%), fibrosis (36.7%), and neuroma (26.6%).

In the present study, no statistically significant difference was revealed between US and NCS for diagnosis of PNI ($p = 0.054$; sensitivity 100%; specificity 95.8%; NNP 100%; PPV 95.8%; overall diagnostic accuracy 97.8%). These findings have high agreement with those by Assy [18], who assessed 30 patients with suspected traumatic nerve

injury by higher resolution US and NCS. Their results for detection on US of neuroma, and local thickening were 100% and 83.3% sensitivity, 94.4% and 100% specificity, 96.7% and 90% accuracy, 92.8% and 100% PPV, and both 100% NPV, respectively. Another study by Cartwright [19] in cadaveric nerves confirmed that US can detect transactions with 89% sensitivity and 95% specificity. Zhu [20] compared the severity of nerve injuries in US with surgical findings and reported the sensitivity of US was 93.2%.

The limitations of this study are the relatively small number of patients and the limited availability of postoperative findings in all patients.

CONCLUSION

Results showed that the US can be used efficiently as a complementary diagnostic tool in the assessment of PNIs based on its capability not only to detect the type of injury and morphological abnormalities of affected nerves, but because it also allows proper assessment of the surrounding tissue, including swelling, scar tissue and neuroma formation, thus altering management decisions as needed. Availability of data and material

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Abbreviations

B-mode . . . Brightness mode

CI	Confidence interval
CSA	Cross-sectional area
CTS	Carpal tunnel syndrome
CMAPs	Compound motor action potentials
EDX	Electrodiagnostic studies
EMG	Electromyography
FCU	Flexor carpi ulnaris
FDP	Flexor digitorum profundus
FPL	Flexor pollicis longus
ME	Medial epicondyle
LS	Longitudinal section
MUAPs	Motor unit action potentials
NCS	Nerve conduction study
NCV	Nerve conduction velocity
NMJ	Neuromuscular junction
NPP	Negative predictive value
PPV	Positive predictive value
TS	Transverse section
RNS	Repetitive nerve stimulation
SD	Standard deviation
SNAPs	Sensory nerve action potentials
US	Ultrasound

REFERENCES

- Wijnjes J, Borchert A, van Alfen N (2021) Nerve ultrasound in traumatic and iatrogenic peripheral nerve injury. *Diagnostics* 11:30. <https://doi.org/10.3390/diagnostics11010030>
- Robinson LR (2000) Traumatic injury to peripheral nerves. *Muscle Nerve* 23(6):863–873
- Robinson L (2015) How electrodiagnosis predicts clinical outcome of focal peripheral nerve lesions. *Muscle Nerve* 52:321–333
- Mostafa MA, Haitham AD, Maïem K (2020) The value of ultrasound in preoperative diagnosis of traumatic nerve injuries in forearm and hand. *Zagazig Univ Med J*. <https://doi.org/10.21608/ZUMJ.2019.19002.1612>
- Mallon S, Starcevic V, Rheinboldt M, Petraszko A (2018) Sonographic evaluation of peripheral nerve pathology in the emergency setting. *Emerg Radiol* 25(5):521–531
- Sunderland S (1990) The anatomy and physiology of nerve injury. *Muscle Nerve* 13:771–784
- Eser F, Aktekin LA, Bodur H, Atan C (2009) Etiological factors of traumatic peripheral nerve injuries. *Neurol India* 57(4):434
- Uzun N, Tanrıverdi T, Savrun FK, Kiziltan ME, Sahin R, Hani-moglu H, Hanci M (2006) Traumatic peripheral nerve injuries: demographic and electrophysiologic findings of 802 patients from a developing country. *J Clin Neuromuscul Dis* 7(3):97–103
- Suk JJ, Walker FO, Cartwright MS (2013) Ultrasonography of peripheral nerves. *Curr Neurol Neurosci Rep* 13(2):328. <https://doi.org/10.1007/s11910-012-0328-x>
- Ciaramitaro P, Mondelli M, Logullo F, Grimaldi S, Battiston B, Sard A, Scarinzi C, Migliaretti G, Faccani G, Cocito D (2010) Traumatic peripheral nerve injuries: epidemiological findings, neuropathic pain and quality of life in 158 patients. *J Peripher Nerv Syst* 15(2):120–127
- Bösch F, Angele MK, Chaudry IH (2018) Gender differences in trauma, shock and sepsis. *Mil Med Res* 5(1):1–10
- Kamble N, Shukla D, Bhat D (2019) Peripheral nerve injuries: electrophysiology for the neurosurgeon. *Neurol India* 67(6):1419
- Martins RS, Bastos D, Siqueira MG, Heise CO, Teixeira MJ (2013) Traumatic injuries of peripheral nerves: a review with emphasis on surgical indication. *Arq Neuropsiquiatr* 71:811–814
- Elfayoumy NM, Elgendy HH, Saad MSA, Labib AA (2020) Role of nerve ultrasound versus electrophysiological studies in the evaluation of nerve injuries. *Egypt J Neurol Psychiatry Neurosurg* 56(1):1–6
- Ferrante MA (2018) The assessment and management of peripheral nerve trauma. *Curr*

Treat Options *Neurol* 20(7):1–18

- Koenig RW, Schmidt TE, Heinen CP, Wirtz CR, Kretschmer T, Antoniadis G, Pedro MT (2011) Intraoperative high-resolution ultrasound: a new technique in the management of peripheral nerve disorders. *J Neurosurg* 114(2):514–521
- Extremite Ü, Sinir P, Cerrahi Y, Intraoperatif T, Kulllanmas U (2007) Ultrasound examination in the surgical treatment for upper extremity peripheral nerve injuries: part I. *Turk Neurosurg* 17:277–282
- Assy MM, Dawood HA, Khamis M (2020) The value of ultrasound in preoperative diagnosis of traumatic nerve injuries in forearm and hand. *Zagazig Univ Med J* 26(5):853–861
- Cartwright MS, Chloros GD, Walker FO, Wiesler ER, Campbell WW (2007) Diagnostic ultrasound for nerve transection. *Muscle Nerve* 35(6):796–799
- Zhu J, Li D, Shao J, Hu B (2011) Preliminary study of the types of traumatic peripheral nerve injuries by ultrasound. *Eur Radiol* 21(5):1097–1101