



BACTERIOLOGICAL SPECTRUM IN NECROTISING FASCIITIS

General Surgery

Dr. Ranjan Kumar Senior Resident, Department of General Surgery, PMCH Patna.

Dr. Binod Kumar Professor, Department of General Surgery, PMCH Patna.

ABSTRACT

Necrotising fasciitis is a relatively rare but seriously fatal soft tissue infection. Prompt treatment with extensive surgical debridement and antibiotics is the main principle. The wound swab culture sensitivity reports of patients (n=50) from July 2022 to December 2023 were analysed. Type I infection was dominating. The involvement of anaerobic bacteria was associated with an increase in the number of surgical revisions. In severe cases the combination of necrotic skin and soft tissue gas facilitates the correct diagnosis, which then should be followed by immediate and repeated debridement. If anaerobes are isolated then wound should be frequently reassessed.

Aim: This study was undertaken to get an idea about the organisms involved in the pathogenesis of necrotising fasciitis, that will help in starting antibiotic therapy before arrival of Culture and sensitivity report.

KEYWORDS

INTRODUCTION:

Necrotising fasciitis is a rapidly progressive infection primarily involving the fascia and subcutaneous tissue. This infection which is usually induced by virulent, toxin producing bacteria can occur in any region of the body but is predominantly located in the abdominal wall, perineum and extremities[1,2]. Giuliano and colleagues[3] classified it into two distinct groups. Type I infections are polymicrobial and involve non group A streptococci, aerobes and/or facultative anaerobes. Type II is usually caused by beta hemolytic group A streptococci alone or in combination with staphylococci. Predisposing factors of the disease are diabetes, alcohol and intravenous drug abuse as well as immunosuppression and peripheral vascular disease[4,5]. However it has been also reported in young and previously healthy individuals.

An early sign of necrotising fasciitis is local erythema and pain out of proportion to findings of physical examination as well as fever. Skin necrosis due to thrombosis of blood vessels at the fascial level is sequel of the initiating process and is often associated with severe sepsis[6-9]. Histomorphological hallmarks are necrosis of superficial fascia with blood vessel thrombosis, severe inflammation of the dermis and subcutaneous fat as well as subcutaneous fat necrosis. Sometimes myonecrosis of underlying skeletal muscle is obvious[10-12]. Diagnostic clues include clinical findings and detection of soft tissue gas or in less obvious cases edema and stranding of fascial layers by computed tomography scanning as well as soft tissue fluid by magnetic resonance imaging[13-16].

This study was conducted to determine microbial flora involved in necrotising fasciitis so that it will help in starting empirical antibiotic therapy before culture and sensitivity report arrives.

MATERIAL AND METHODS:

The data for the study was obtained from the patients hospitalised with a provisional diagnosis of necrotising fasciitis on clinical evaluation and are admitted at Patna Medical College and hospital. Patients presenting with signs and symptoms of necrotising fasciitis admitted during July 2022 to December 2023 at Patna medical college and hospital were counselled for investigations and treatment of necrotising fasciitis and its complications. 50 patients were treated in the above hospital during this period. Wound swabs of these patients were sent for culture and sensitivity.

RESULTS:

Type of organism	Number of patients	Percentage
Aerobic	32	64
Anaerobic	1	2
Polymicrobial	7	14

Type of organism	No. of patients	Mortality	Percentage
Gram +ve cocci	18	1	5.55%
Gram +ve bacilli	1	1	100%
Gram -ve bacilli	24	2	8.33%
Polymicrobial	7	2	28.57%

ANTIBIOTICS	NUMBER OF PATIENTS	PERCENTAGE
Ceftriaxone	18	36%
Cefotaxime	11	22%
Ciprofloxacin	9	18%
Amikacin	6	12%
Gentamycin	4	8%
Meropenem	2	4%

Antibiotic Sensitivity

Outcome:

Organisms	Number of patients	Percentage
Klebsiella	13	26
Streptococcus	10	20
Staphylococcus	8	16
Polymicrobial	7	14
E. Coli	7	14
Proteus	3	6
Pseudomonas	1	2
Clostridium	1	2

DISCUSSION:

In our study the most common infection was a type I-polymicrobial infection. In the present study predominant organism was Klebsiella seen in 13 cases (26%), followed by Staphylococcus aureus seen in 10 cases (20%). Polymicrobial infection was seen in 7 cases (14%). Most isolates were sensitive to ceftriaxone(36%).

CONCLUSION:

The most common bacterial isolate found was Klebsiella (26%) and Polymicrobial infections were seen in 7 cases (14%). As most isolates were sensitive to ceftriaxone, it can be started empirically before knowing culture and sensitivity report

REFERENCES:

- Singh G, Sinha SK, Adhikary S, Babu KS, Ray P, Khanna SK. Necrotising infections of soft tissues-a clinical profile. Eur J Surg. 2002;168(6):366-71.
- Green RJ, Dafoe DC, Raffin TA. Necrotizing fasciitis. Chest. 1996 Jul;110(1):219-29.
- Giuliano A, Lewis F Jr, Hadley K, Blaisdell FW. Bacteriology of necrotizing fasciitis. Am J Surg. 1977 Jul;134(1):52-7.
- Chen JL, Fullerton KE, Flynn NM. Necrotizing fasciitis associated with injection drug use. Clin Infect Dis. 2001 Jul 1;33(1):6-15.
- Roujeau JC. Necrotizing fasciitis. Clinical criteria and risk factors. Ann Dermatol Venerol. 2001 Mar;128(3 Pt 2):376-81. Review.
- McHenry CR, Piotrowski JJ, Petrinic D, Malangoni MA. Determinants of mortality for necrotizing soft-tissue infections. Ann Surg. 1995 May;221(5):558-63.
- Wong CH, Chang HC, Pasupathy S, Khin LW, Tan JL, Low CO. Necrotizing fasciitis: clinical presentation, microbiology, and determinants of mortality. J Bone Joint Surg Am. 2003 Aug;85-A(8):1454-60.
- Elliott DC, Kufera JA, Myers RA. Necrotizing soft tissue infections. Risk factors for mortality and strategies for management. Ann Surg. 1996 Nov;224(5):672-83.
- Wong CH, Wang YS. The diagnosis of necrotizing fasciitis. Curr Opin Infect Dis. 2005 Apr;18(2):101-6. Review.
- Umbert IJ, Winkelmann RK, Oliver GF, Peters MS. Necrotizing fasciitis: a clinical, microbiologic and histopathologic study of 14 patients. J Am Acad Dermatol. 1989 May;20(5 Pt 1):774-81.
- Barker FG, Leppard BJ, Seal DV. Streptococcal necrotising fasciitis: comparison between histological and clinical features. J Clin Pathol. 1987 Mar;40(3):335-41.
- Bakleh M, Wold LE, Mandrekar JN, Harmsen WS, Dimashkieh HH, Baddour LM. Correlation of histopathologic findings with clinical outcome in necrotizing fasciitis.

- Clin Infect Dis. 2005 Feb 1;40(3):410-4.
13. Wysoki MG, Santora TA, Shah RM, Friedman, AC. Necrotizing fasciitis: CT Characteristics. Radiology. 1997 Jun;203(3):859-63.
 14. Fugitt JB, Puckett ML, Quigley MM, Kerr SM. Necrotizing fasciitis. Radiographics. 2004 Sep-Oct;24(5):1472-6.
 15. Schmid MR, Kossmann T, Diewell S. Differentiation of necrotizing fasciitis and cellulitis using MR imaging. AJR Am J Roentgenol. 1998 Mar;170(3):615-20.
 16. Brothers TE, Tagge DU, Stutley JE, Conway WF, Del Schutte H Jr, Byrne TK. Magnetic resonance imaging differentiates between necrotizing and non-necrotizing fasciitis of the lower extremity. J Am Coll Surg. 1998 Oct;187(4):416-21.