



PERSISTENT ACUTE KIDNEY INJURY IN CRITICAL ILLNESSES; CONNECTING THE MISSING LINKS.

Medical Science

Dr. Ajay Gupta	Hod and Director, Deptt. of Critical care, Max Super Speciality Hospital, Shalimar Bagh, New Delhi.
Dr. Surinder Singh Sodhi	Attending Consultant, Deptt. of Critical care, Max Super Speciality Hospital, Shalimar Bagh, New Delhi.
Dr. Vinod Kumar Jaiswal	Senior medical officer, Deptt. of Nephrology, Max Super Speciality Hospital, Shalimar Bagh, New Delhi.

ABSTRACT

Acute kidney injury (AKI) is now recognized as a heterogeneous syndrome that not only affects acute morbidity and mortality, but also a patient's long-term prognosis. In this narrative review, an update on various aspects of AKI in critically ill patients will be provided. Focus will be on prediction and early detection of AKI (e.g., the role of biomarkers to identify high-risk patients and the use of machine learning to predict AKI), aspects of pathophysiology and progress in the recognition of different phenotypes of AKI, as well as an update on nephrotoxicity and organ cross-talk. In addition, prevention of AKI (focusing on fluid management, kidney perfusion pressure, and the choice of vasopressor) and supportive treatment of AKI is discussed. Finally, post-AKI risk of long-term sequelae including incident or progression of chronic kidney disease, cardiovascular events and mortality, will be addressed.

KEYWORDS


INTRODUCTION

The availability of a consensus definition of acute kidney injury (AKI) [1] has been an important step in establishing AKI epidemiology. AKI affects 30–60% of critically ill patients and is associated with acute morbidity and mortality [2]. Evidence is also accumulating that the burden of AKI extends beyond the acute phase with progression to chronic kidney disease (CKD), increased risk of cardiovascular complications, recurrent episodes of AKI and long-term mortality [3]. Prevention of development and/or progression is currently limited to hemodynamic and fluid status optimization and avoidance of nephrotoxins. Search for a specific pharmacologic treatment is hampered by the late diagnosis and the complex and incompletely elucidated pathophysiology. Progress in the management of AKI is to be expected from the recognition that AKI is a very heterogeneous syndrome with variable etiology, pathophysiology and clinical presentation. CKD, characterized by a decreased number of functioning nephrons [4], is probably the most important factor that defines the susceptibility of a subject to developing AKI under nephrotoxic exposure [5]. Furthermore, baseline renal function is the most significant variable in assessment of patients' AKI risk; therefore, it is a component of several risk scores used in different clinical settings [6, 7]. However, AKI is a recognized risk factor for CKD [8], by damaging nephrons and reducing first renal functional reserve (RFR) and then glomerular filtration rate (GFR) [9]. In 2012, the Kidney Disease: Improving Global Outcomes (KDIGO) AKI workgroup defined AKI and CKD as syndromes mainly differentiated by the timing of kidney function impairment (i.e. a period of ≤ 7 days for AKI and > 90 days for CKD) and its reversibility [10]. Additionally, KDIGO also defined the persistence of kidney functional impairment for a duration of between 7 and 90 days as acute kidney disease (AKD). In 2016, the Acute Disease Quality Initiative (ADQI) group further defined AKI as persistent if variations of serum creatinine (sCr) and/or urine output, according to KDIGO criteria, last beyond 48 h from AKI onset. On the contrary, a rapid reversal of AKI within 48 h has been defined as transient AKI [11]. However, persistent AKI needs to be consistently defined in the literature. While AKI and CKD are well-described syndromes, transient or persistent AKI and AKD are relatively new entities. This review aims to highlight the critical phases of persistent AKI in ICU patients in terms of both clinical and mechanistic features to offer to clinicians and researchers an updated basis from which to start improving patients' care and direct future research.

Before An Aki Episode: Risk Factors For Persistent Aki In Critically Ill Patients -

The traditional classification by which transient AKI is due to a brief kidney injury while persistent AKI is due to a prolonged kidney injury is partially obsolete. Depending on patients' susceptibility, a short

kidney injury may evolve into persistent AKI and AKD, triggering the AKI-to-CKD transition. While patient-related risk factors determine the individual patient's susceptibility to developing AKI, procedure-related risk factors are the exposure of a patient to potentially nephrotoxic agents. Susceptibility and exposure together define each patient's risk of AKI. According to this concept, it is reasonable to assume that persistent versus transient AKI occurs when a patient has a high susceptibility to developing kidney damage and/or undergoes intensive, repeated potentially nephrotoxic exposures. The first case is due to patient-related risk factors that limit the adaptability to conditions potentially harmful for the kidneys (i.e. pre-existing CKD or low/absent RFR, comorbidities and clinical conditions that may contribute to AKI development). The exposures to which the patient can be subjected vary in intensity, number and frequency (e.g. nephrotoxic agents, emergency surgery, liver failure, sepsis, etc.) [12]. As patients' susceptibility increases, the intensity of exposure that can lead to persistent AKI decreases. On the contrary, in the presence of severe nephrotoxic exposure (e.g. sepsis), even a lowly susceptible subject may develop persistent AKI, AKD and CKD. Critically ill patients are exposed to various potential nephrotoxic substances, including prescribed therapeutic agents, that could be responsible for underrecognized persistent AKI. In addition, although several studies have identified conditions that predispose to persistent AKI [13, 14], risk factors are currently under investigation limiting the designing of simplified risk prediction models.



The Role of Mitochondria in Acute Kidney Injury and Chronic Kidney Disease and Its Therapeutic Potential

During An Aki Episode: Clinical And Mechanistic Features Of Persistent Aki In Critically Ill Patients -

Predictive and early diagnostic criteria of persistent versus transient AKI The early diagnosis of transient/persistent AKI has a key role in

critical care management. Assessing the variation of sCr for AKI diagnosis requires ascertaining its change from a known base-line. In literature, four different baseline sCr definitions are described: (i) pre-admission sCr: sCr measured in a time-period of a maximum of 365 days and a minimum of 7 days from the moment of hospital admission; (ii) admission sCr: the first sCr measured at hospital admission; (iii) nadir sCr: the lowest measured value of sCr during the first 3 days of the ICU stay; (iv) estimated sCr: calculated using a back-estimation equation, starting from an assumed GFR of 75 or 100 mL/min, or assigning GFR from age- and sex-standardized reference tables [15, 16]. In clinical practice, the most used definitions are pre-admission sCr and admission sCr, while nadir sCr and estimated sCr are mostly used in experimental settings. However, it needs to be clarified which is the best method for assessing baseline sCr. Based on the increase in sCr levels that starts when > 50% of nephrons are impaired, it is evident that sCr is a late marker of kidney dysfunction [17]. Nevertheless, equations for GFR estimation have a limited use in critically ill patients in which sCr is not at a steady state [18]. Numerous other methods for renal function assessment have been proposed, such as the Jelliffe equation [19], the kinetic eGFR [20] and the point-of-care determination of GFR through a fiberoptic ratiometric fluorescence analyzer [21], but they are still under investigation, while more established methods, such as GFR measurement with inulin or iothexol, are cumbersome and time-consuming. However, measured GFR would allow a rapid diagnosis of AKI and quantification of the extent of injury without the limits due to sCr and urine output. In addition to the late increase of sCr during a kidney damage, both sCr and urine output can be influenced by non-renal factors. This is particularly obvious in ICU patients with frequent muscle wasting syndrome, hyperhydration, hypovolemia and diuretic use [15, 22]. Recently, the 23rd ADQI expert group proposed expanding the AKI definition and staging by considering both kidney dysfunction and early kidney damage [23]. Although new biomarkers, functional tests [e.g. RFR assessment, furosemide stress test (FST), etc.] and ultrasonographic markers have been widely investigated with the aim to improve early diagnosis of AKI, they are still underused, even in critical care settings. Among ultrasonographic markers, the renal resistive index (RRI) value, measured in a segmental artery, is commonly used to assess blood flow in renal intra-parenchymal vessels. RRI is calculated using the formula: $RRI = (\text{peak systolic velocity} - \text{end-diastolic velocity}) / \text{peak systolic velocity}$, thus being a valuable indicator of the resistance to flow within the kidney. In addition, studies suggest that RRI value may predict persistent AKI in some clinical settings [24]. Like CKD, AKI does not have early specific symptoms. Beyond the issue related to “when” to use markers, another critical concern is “which” marker/s to use. In fact, AKI is a very heterogeneous syndrome with different etiologies, pathophysiological pathways and clinical features, thus it is difficult to generalize. Therefore, for instance, RRI is a good predictor of persistent cardiac surgery-associated AKI [25], but its role is less defined in sepsis-associated AKI [26, 27]. Then, the predictive capability of different biomarkers varies depending on the etiology of AKI and the group of patients considered. In addition to all these reasons, the main issue concerning the scarce clinical use of biomarkers is the need for more awareness of the importance of early AKI diagnosis and treatment among healthcare team members, as revealed by a recent Italian survey [28, 29]. Table 1 summarizes potential markers of persistent vs transient AKI in different settings, some of which are not yet available for clinical use.

Cellular And Molecular Differences Between Persistent And Transient AKI-

Immediately after a kidney injury, repair processes are activated. Some of them allow a rapid return of renal function, and others lead to a persistence of renal dysfunction. The 13th ADQI Consensus Conference defined as “adaptive repair” as those processes that lead to a “resolution of the renal structure free of long-term sequelae” and as “maladaptive repair” a “process that results in a durable reduction in kidney function usually associated with a change in renal structure” [30]. Adaptive and maladaptive repair processes together with the injury mechanisms determine the number of irreversibly lost nephrons and then the long-term prognosis of kidney function. Several pathophysiological processes that promote maladaptive repair and AKI-to-CKD transition have been identified and gained growing attention, such as microvascular damage and loss of peritubular capillaries, systemic inflammation and pro-fibrotic cytokine secretion, cell cycle arrest and cellular senescence [31], pericyte activation with myofibroblasts generation [32] and metabolic re-

programming of tubular epithelial cells (TECs). Some of these mechanisms, such as cell cycle arrest and metabolic reprogramming of TECs, start as adaptive repair processes, but, if persistent, become maladaptive [33]. When not completely repaired, TECs undergo cell cycle arrest at G2/M, as a protective mechanism for maintaining genomic stability. Nonetheless, TECs with arrested cell cycle acquire a pro-fibrotic secretory phenotype known as senescence-associated secretory phenotype, which promotes the development of kidney fibrosis [34]. It has been demonstrated that AKI leads to a permanent loss of TECs even when full clinical renal recovery occurs (see below). Kidney function can be augmented by polyploidization of TECs in unaffected nephrons (i.e. compensatory hypertrophy) [35]. However, TECs polyploidy promotes TECs senescence, progressive interstitial fibrosis and AKI-to-CKD transition. Although the exact mechanisms are still unknown, the metabolic reprogramming of TECs during an AKI episode seems to play a pivotal role in inhibiting AKI-to-CKD transition. In the early 2000s, activated T cells were found to switch their metabolism to glycolysis, thus adjusting their energetic and biosynthetic needs in response to changing conditions [37]. Some data suggest that, like immune cells, TECs can also reprogram their metabolism in response to kidney injuries. In particular, investigations focused on sepsis-associated AKI [38, 39]. It has been hypothesized that proximal TECs act as danger sensors by responding to pathogen-associated molecular patterns (PAMPs), secreting pro-inflammatory cytokines into the tubule, and activating TECs in other nephron segments, thus amplifying tubular inflammation [40]. These findings suggest the presence of an early proinflammatory phase followed by a later catabolic phase, probably related to mitochondrial functions. They are respectively characterized by a shift in metabolism to aerobic glycolysis, generating appropriate inflammatory responses, limiting oxidative damage and rearranging substrates to prevent cell death, and to oxidative phosphorylation, promoting renal function recovery [41] (Fig. 1). Mitochondria regulate intracellular calcium homeostasis and reactive oxygen species (ROS) levels, and control cell cycle, cellular differentiation and death through their biogenesis, dynamics, mitophagy, bioenergetics and mitochondrion endoplasmic reticulum cross-talk. During the early glycolytic phase, TECs remove dysfunctional mitochondria from the cytosol via mitophagy (i.e. a form of autophagy that selectively eliminates damaged mitochondria) to avoid ROS generation. In experimental models of AKI, impaired mitophagy has been associated with a reduced renal recovery [42] while a diminished AKI progression and improved survival have been found in conditions of increased mitophagy [43]. After the first proinflammatory phase, the restitution of a functioning mitochondria pool via biogenesis (i.e. the generation of new mitochondrial mass) is mandatory to cope with the increased metabolism and energy demands, thus promoting renal recovery and survival. Recent findings suggest that impaired biogenesis leads to a maladaptive repair after AKI and renal fibrosis [44] (Fig. 1). As the glycolytic phase is fundamental for appropriate responses to kidney injuries, the capacity to turn off the inflammation is required for renal recovery. In fact, cytokines and damage-associated molecular patterns (DAMPs), released by injured TECs and endothelial cells in the proinflammatory state, participate in the development of a pro-fibrotic milieu. This activates pericytes to proliferate and evolve into myofibroblasts, thereby inducing matrix deposition and subsequent decreased capillary density, thus leading to maladaptive repair, renal fibrosis and AKI-to-CKD conversion [45]. In immune cells the early switch to glycolysis is also required to develop a trained immunity (i.e. the process by which the innate immune system develops memory and adjusts its response to future insults). The fascinating hypothesis of a biological memory that drives the kidney to respond to future insults in humans has been postulated, but it is still a subject of debate.

After An AKI Episode: Prognosis Of Persistent AKI In Critically Ill Patients:

Persistent AKI has been correlated with higher morbidity and mortality, and a higher progression rate to AKD and CKD than transient AKI [47]. It seems that prompt interventions can change the course of AKI and improve patients' outcomes [47]. The continuum of AKI-to-CKD proposed by 23rd ADQI consensus group is based on the AKD definition that can be considered the evolution of a persistent AKI episode that has not been promptly and appropriately treated or, more generally, in which the abovementioned mechanisms (and probably many others) have reduced functioning nephrons, thus affecting RFR and/or GFR [11]. Unfortunately, like persistent AKI, the AKD definition also depends on sCr level variations and its incidence

is probably underestimated because of muscle wasting, hyperhydration, hyperfiltration, etc. Plasma cystatin C has been proposed in critical settings as an alternative marker to assess renal recovery. Despite promising preliminary results, it is still far from being routinely used in clinical practice [15, 48]. Moreover, even in the case of sCr levels returning to baseline, a complete recovery of baseline renal conditions, in terms of both kidney structure and function, may not occur. A subclinical damage may be identified by new-onset or worsening proteinuria, arterial hypertension and/or a drop in RFR [11, 49]. Moreover, patients with AKD in whom sCr levels returned to baseline have been demonstrated to be more prone to further kidney damage and other adverse events [50].

Persistent AKI In Transplanted Patients -

The classification of transient vs persistent AKI has also been used in the setting of liver and lung transplantation. In these settings, AKI has a high incidence, reaching more than the half of transplanted patients (studies have reported an incident of AKI up to 64% in the liver [51] and 69% in lung transplant recipients [52]) and a worse prognosis in terms of morbidity and mortality [51, 53]. In kidney transplantation, AKI, manifesting as either delayed graft function (DGF) or de novo post-transplant acute deterioration of graft function, occurs in 30% of kidneys coming from deceased donors and in 50% of those coming from deceased donors after cardiac death [54], and affects short- and long-term transplant outcomes [55]. In transplanted patients, multifarious risk factors contribute to the development of AKI. They can be classified as pre-transplant comorbidities, related to both donor and recipient, perioperative problems and post-transplant factors, mainly associated with exposure to nephrotoxic agents, such as calcineurin inhibitors and antimicrobial drugs [56]. There is growing evidence demonstrating that persistent AKI after lung transplantation is associated with worse renal and non-renal outcomes [57], including a higher mortality [57, 58]. A recent study on liver-transplanted patients showed that persistent AKI is associated with an increased incidence rate of graft failure and a decreased graft survival time [59]. According to recently published data, a linear association between duration of DGF and graft loss has been found. In particular, a prolonged DGF beyond 7 days posttransplant has been associated with a > 40% greater risk of death-censored graft loss [60]. These data confirm the results of prior studies [61, 62] and underline the importance of clinical interventions that aim to prevent or reduce the duration of DGF.


Clinical Approach To Critically Ill Patients With Persistent AKI Prevention And Diagnosis -

According to the varied pathophysiology of AKI, the choice of which clinical parameters and risk scores, biomarkers and functional tests to use should be considered and tailored to the AKI setting, clinical phase and center availability or local expertise. The rapid diagnosis of persistent AKI may allow reconsideration of all causes of AKI, their correction when possible and remodulation of the therapy [11]. In critically ill individuals, AKI is frequently multifactorial and depends on both patient- and agent-related factors. Although patient-related factors are commonly unmodifiable, the ones related to nephrotoxic agents, when possible, must be reassessed according to the literature evidence. The timely recognition of persistent AKI potentially enables the physician to promptly withdraw potential nephrotoxic drugs and to interrupt the pathogenetic mechanisms of kidney injury, thus preventing further renal impairment [63]. AKI awareness programs could help to improve AKI awareness and recognition, and they should include a broader and more appropriate use of early biomarkers of kidney damage [23] and functional tests in critical care settings and the involvement of all healthcare team members, each for their own competence [28].

Treatment And Follow-up -

The treatment of persistent AKI largely depends on its cause(s) and the setting in which it occurs, and varies according to the clinical phase. Therapeutic indications for AKI due to specific kidney diseases, such as acute glomerulonephritis, acute renal vascular disease, etc., are beyond the scope of this review. At the beginning of an AKI episode, therapeutic indications for persistent AKI do not differ from the ones for transient AKI. They mainly concern the optimization of fluid management and hemodynamic support and the withdrawal of nephrotoxic agents, when it is possible. In cases in which agents potentially harmful for the kidneys are required, benefits and risks should be balanced and they should be administered for as long as needed with a close monitoring of renal function [22]. The risk of drug

(or metabolites) accumulation and toxicity due to kidney dysfunction, altered protein binding and variable volume of distribution should be considered [5]. Simultaneously, the risk of therapeutic failure due to drug under-dosing should be balanced, particularly in patients undergoing renal replacement therapy (RRT). As an accelerated RRT strategy in critically ill patients with AKI has failed to demonstrate a lower risk of death [64], RRT should be considered when metabolic and fluid demands exceed the kidney's capacity to meet them [10]. The choice of modality depends on the patient's clinical status, the center's available resources and the expertise of personnel. Although conclusive randomized clinical trials are missing, in hemodynamically unstable patients, continuous RRT is more appropriate than intermittent RRT. As a consequence, also for maintenance hemodialysis and peritoneal dialysis patients, modality transitions should be considered based on their status [5]. As early diagnosis is fundamental to addressing preventive strategies, transitioning care from ICU to other hospital departments and from hospital to community is crucial to defining optimal follow-up care. It should be personalized according to patients' overall risk (e.g. pre-existent CKD, age, etc.), AKI severity and duration (e.g. need for RRT, persistent vs transient AKI) and renal recovery [65]. Despite several medical advances meaning that critically ill patients with AKI increasingly have better survival [66], they are nevertheless at risk of adverse outcomes, including renal failure [2], progression to CKD, and cardiovascular events such as new-onset arterial hypertension or myocardial infarction [67]. Unfortunately, few patients benefit from early nephrological evaluation that could decrease mortality among survivors of dialysis-receiving AKI [68]. The improvement of care includes appropriate risk stratification, monitoring of kidney function, the management of CKD complications, blood pressure control, medication reconciliation and education [69]. Risk stratification and a kidney function monitoring plan at discharge are needed to improve post-AKI care. In their randomized controlled trial, Silver et al. [70], compared structured nephrologist follow-up versus usual care in survivors of AKI and showed that the enrollment of fully eligible patients was a barrier, primarily due to the requirement for post discharge, in-person visits. The reason was that eligible patients had long hospital travel times/hospital-related fatigue. Additionally, they did not identify any difference in major adverse kidney events at 1 year. However, the high frequency of events highlights the vulnerability of this population and the urgent need for feasible and effective interventions for survivors of AKI. Strategies for optimal follow-up care of patients post-AKI are still under investigation.



The continuum of AKI and AKD in patients with and without RFR. Before an AKI episode, patient-related risk factors define the kidney's susceptibility to kidney damage. Based on this, a different intensity of exposure is required to lead to subclinical to clinical AKI. Residual renal function, including RFR, is critical among patient-related risk factors for AKI. In the presence of an intact RFR, even in case of intensive, repeated potentially nephrotoxic exposures, the damage to the kidney may remain subclinical. In contrast, if RFR is lost, even on mild exposures, the kidney damage may become clinically manifest.

ONGOING RESEARCH ON PERSISTENT AKI Prevention And Diagnosis-

Several studies have assessed promising tools as prediction models for AKI based on machine-learning methodologies [71]. However, the performance of models may vary in different scenarios, and the abovementioned studies do not focus on the critical care setting. Furthermore, few studies involving comparisons of models have concentrated on AKI prediction in the ICU and the majority of them lack external and prospective validation [72]. Liang et al. built models to predict AKI within 48 h in critically ill patients by using three transcontinental databases, and then they evaluated the clinical effect of the model through a 1-year prospective validation. A total of 2532 patients were admitted to the center for prospective validation; 358 positive results were predicted and 344 patients were diagnosed with severe AKI, with the best sensitivity of 0.72, specificity of 0.80 and area under the receiver operating characteristic of 0.84. However, the AKI predictions using artificial intelligence is still controversial. The assessment of RFR, i.e. the capability of the kidney to increase GFR

under conditions of physiologic or pathologic higher functional demand, thus being an indicator of renal function- ing mass, allows better definition of susceptibility to developing acute damage. Different methods to measured RFR have been proposed, such as oral protein load and infusion of amino acids. In the presence of an intact RFR, even in case of intensive, repeated potentially nephrotoxic exposures, the damage to the kidney may remain subclinical. On the contrary, if RFR is progressively lost, the susceptibility of the kidney progressively in- creases and even in the presence of a mild exposure, the kidney damage may become clinically manifest [9]. However, although RFR is an important tool to assess the kidney's capacity to respond to conditions of higher functional demand, the exam is challenging and may be stressful for some patients. For this reason, based on the pathophysiological hypothesis of kidney damage during intra-abdominal hypertension, the intra- parenchymal renal resistive index variation (IRRV) test was de- signed to predict the presence of RFR [74]. Samoni et al . found a correlation between IRRIV test and RFR measured by using an oral protein loading test in healthy volunteers and in cardiac surgery patients [75]. However, these findings must be confirmed in large patient cohorts and different settings. Among functional stress tests to early diagnose persistent versus transient AKI, the FST, based on the pharmacokinetic properties of furosemide, assesses the functional integrity of the renal tubule, and it may be a reliable predictor of AKI progres- sion to severe stages and requirement of dialysis in ICUs [80]. It could help clinicians identify patients with tubular injury and at higher risk of AKI or CKD progression. Chen et al . [76]. in their systematic review and meta-analysis showed that the FST is a simple tool for the identification of AKI populations at high risk of AKI progression and the need for RRT, and the diagnostic per- formance of FST in RRT prediction is better in early AKI popula- tion. Koyner et al . [82]. showed that, in the setting of early AKI, FST urine output outperformed biochemical biomarkers for pre- diction of progressive AKI, need for RRT and inpatient mortality. Using an FST in patients with increased biomarker levels im- proves risk stratification. However, several aspects can limit FST such as the retrospective nature of most studies, variability of doses and timing. Additionally, some of the published studies solely rely on AUC values to define the predictive capacity of the test, while actual or absolute predicted risk could be more ap- propriate to assess risk prediction models [83]. During an AKI episode, accurate ascertainment of GFR is cru- cial for the diagnosis of early kidney disease [84]. Patients at risk of early kidney disease when GFR is normal-to-elevated would benefit from having their GFR measured using more accurate and precise techniques. Recent advances in technology to mea- sure fluorescent compounds through the skin are providing a new approach for real-time monitoring of GFR [77]. After the in- jection of an ideal GFR marker, its concentration in the plasma reaches a peak, followed by an exponential decay due to its diffu- sion into the extracellular space. Using transdermal sensors, the plasma disappearance rate can be measured rather than their steady state concentration. This eliminates the delay inherent in using an endogenous marker of filtration and permits continuous monitoring of GFR.

Treatment And Follow-up -

Without effective pharmacologic therapies, severe AKI is usually managed through RRT. Unfortunately, no specific therapy is currently available to prevent or treat AKI. However, various poten- tial agents, including novel compounds, re-purposed drugs and cell-based therapies, are in ongoing early-phase clinical trials. Various pathways have been targeted for this purpose, includ- ing oxidative and mitochondrial stress, cellular metabolism and repair, inflammation, apoptosis and hemodynamics [63, 86]. After an AKI episode, the need for an appropriate follow- up of patients recovering from AKI or AKD is increasingly be- ing recognized [65]. Kidney function and proteinuria monitor- ing, new biomarkers of renal tubular fibrosis measurement [e.g. Chemokine (C-C motif) ligand 14 (CCL14) [87], Dickkopf-3 (DKK3) [88], etc.] and functional tests application (e.g. RFR assessment) are some of the tools proposed for a better risk stratification and prevention of new episodes of AKI. Simultaneously, drugs that act on the renin-angiotensin-aldosterone system, such as sodium-glucose co-transporter-2 inhibitors, have been shown to improve clinical outcome in CKD patients [89] and open a poten- tial therapeutic strategy in post-AKI patients.

CONCLUSIONS

Persistent AKI has been associated with worse outcomes than transient AKI, including a higher rate of AKI-to-CKD transition. As prompt

interventions can change its course, persistent AKI should be diagnosed early by measuring new biomarkers and applying functional tests. In the absence of effective pharmacological therapies, the prevention of AKI has a crucial role, as well as the recon- sideration of all causes of AKI, their correction when possible and modulation of the therapy. Furthermore, the follow-up post-AKI should be personalized according to patients' overall risk, AKI severity and duration, and renal recovery, considering that, also in case of full clinical recovery, subclinical damage may be present and predispose to further kidney damage and other adverse event(s). Ongoing research on persistent AKI may lead to a better as- sessment of kidney susceptibility, to an earlier and more accu- rate diagnosis, and to a pharmacological therapy, thus improving the care of AKI and AKD patients

Conflict Of Interest Statement None declared.

Data Availability Statement No new data were generated or analyzed in support of this research.

REFERENCES:

- Hoste EAJ, Bagshaw SM, Bellomo R et al. Epidemiology of acute kidney injury in critically ill patients: the multinational AKI-EPI study. *Intensive Care Med* 2015; 41 :1411–23. <https://doi.org/10.1007/s00134-015-3934-7>.
- Coca SG, Singanamala S, Parikh CR. Chronic kidney disease after acute kidney injury: a systematic review and meta- analysis. *Kidney Int* 2012; 81 :442–8. <https://doi.org/10.1038/ki.2011.379>.
- Chawla LS, Eggers PW, Star RA et al. Acute kidney in- jury and chronic kidney disease as interconnected syn- dromes. *N Engl J Med* 2014; 371 :58–66. <https://doi.org/10.1056/NEJMr1214243>.
- Denic A, Lieske JC, Chakkera HA et al. The substantial loss of nephrons in healthy human kidneys with aging. *J Am Soc Nephrol* 2017; 28 :313–20. <https://doi.org/10.1681/ASN.2016020154>.
- De Rosa S, Samoni S, Villa G et al. Management of chronic kidney disease patients in the intensive care unit: mixing acute and chronic illness. *Blood Purif* 2017; 43 :151–62. <https://doi.org/10.1159/000452650>.
- Thakar CV, Arrigain S, Worley S et al. A clinical score to pre- dict acute renal failure after cardiac surgery. *J Am Soc Nephrol* 2005; 16 :162–8. <https://doi.org/10.1681/ASN.2004040331>.
- Parolari A, Pesce LL, Pacini D et al. Risk factors for perioper- ative acute kidney injury after adult cardiac surgery: role of perioperative management. *Ann Thorac Surg* 2012; 93 :584–91. <https://doi.org/10.1016/j.athoracsur.2011.09.073>.
- Kurzhaen J, Dellepiane S, Cantaluppi V et al. AKI: an increasingly recognized risk factor for CKD development and progression. *J Nephrol* 2020; 33 :1171–87. <https://doi.org/10.1007/s40620-020-00793-2>.
- Sharma A, Mucino MJ, Ronco C. Renal functional reserve and renal recovery after acute kidney injury. *Nephron Clin Pract* 2014; 127 :94–100. <https://doi.org/10.1159/0003637>.
- Kellum JA, Lameire N, Aspelin P et al. KDIGO clinical practice guideline for acute kidney injury. *Kidney Int Suppl* 2012; 2 :1–141.
- Chawla LS, Bellomo R, Bihorac A et al. Acute kidney dis- ease and renal recovery: consensus report of the Acute Dis- ease Quality Initiative (ADQI) 16 Workgroup. *Nat Rev Nephrol* 2017; 13 :241–57. <https://doi.org/10.1038/nrneph.2017>.
- Silver S, Nadim MK, O'Donoghue D et al. Community health care quality standards to prevent acute kidney injury and its consequences. *Am J Med* 2020; 133 :552–60. <https://doi.org/10.1016/j.amjmed.2019.10.038>.
- Brown J, Kramer R, MacKenzie T et al. Determinants of acute kidney injury duration after cardiac surgery: an externally validated tool. *Ann Thorac Surg* 2012; 93 :570–6. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3624763/pdf/nihms412728.pdf>.
- Basu RK, Wang Y, Wong HR et al. Incorporation of biomark- ers with the renal angina index for prediction of severe AKI in critically ill children. *Clin J Am Soc Nephrol* 2014; 9 :654–62. <https://doi.org/10.2215/CJN.09720913>.
- De Rosa S, Samoni S, Ronco C. Creatinine-based definitions: from baseline creatinine to serum creatinine adjustment in intensive care. *Crit Care* 2016; 20 :1–5.
- Cooper DJ, Plewes K, Grigg MJ et al. An evaluation of com- monly used surrogate baseline creatinine values to clas- sify AKI during acute infection. *Kidney Int Rep* 2021; 6 :645–56. <https://doi.org/10.1016/j.ekir.2020.12.020>.
- Delanaye P, Schaeffner E, Ebert N et al. Normal reference val- ues for glomerular filtration rate: what do we really know? *Nephrol Dial Transplant* 2012; 27 :2664–72. <https://doi.org/10.1093/ndt/gfs265>.
- Hoste EAJ, Damen J, Vanholder RC et al. Assessment of renal function in recently admitted critically ill patients with nor- mal serum creatinine. *Nephrol Dial Transplant* 2005; 20 :747–53. <https://doi.org/10.1093/ndt/gh707>.
- Jelliffe R. Estimation of creatinine clearance in patients with unstable renal function, without a urine specimen. *Am J Nephrol* 2002; 22 :320–4. <https://doi.org/10.1159/000065221>.
- Pianta TJ, Endre ZH, Pickering JW et al. Kinetic estimation of GFR improves prediction of dialysis and recovery after kid- ney transplantation. *PLoS One* 2015; 10 :1–15. <https://doi.org/10.1371/journal.pone.0125669>.
- Wang E, Meier DJ, Sandoval RM et al. A portable fiberop- tic ratiometric fluorescence analyzer provides rapid point-of-care determination of glomerular filtration rate in large animals. *Kidney Int* 2012; 81 :112–7. <https://doi.org/10.1038/ki.2011.294>.
- Goldstein SL, Kane-gill SL, Liu KD et al. Controversies in acute kidney injury: conclusions from a Kidney Disease: Improving Global Outcomes (KDIGO) Conference. *Kidney Int* 2021; 98 :294–309.
- Ostermann M, Zarbock A, Goldstein S et al. Recommenda- tions on acute kidney injury biomarkers from the Acute Dis- ease Quality Initiative Consensus Conference: a consensus statement. *JAMA Netw Open* 2020; 3 :e2019209. <https://doi.org/10.1001/jamanetworkopen.2020.19209>.
- Faubel S, Patel NU, Lockhart ME et al. Renal relevant ra- diology: use of ultrasonography in patients with AKI. *Clin J Am Soc Nephrol* 2014; 9 :382–94. <https://doi.org/10.2215/CJN.04840513>.
- Guinot PG, Bernard E, Abou Arab O et al. Doppler-based renal resistive index can assess progression of acute kidney injury in patients undergoing cardiac surgery. *J Cardiothorac Vasc Anesth* 2013; 27 :890–6. <https://doi.org/10.1053/j.jvca.2012.11.024>.
- Fu Y, He C, Jia L et al. Performance of the renal resistive index and usual clinical indicators in predicting persistent AKI. *Ren Fail* 2022; 44 :2028–38. <https://doi.org/10.1080/0886022X.2022.2147437>.

27. Zhang Y, Zhu J, Zhang C et al. Non-invasive early prediction of septic acute kidney injury by doppler-based renal resistive indexes combined with echocardiographic parameters: an experimental study. *Front Med* 2021; 8 : 1–12. <https://doi.org/10.3389/fmed.2021.723837>.
28. De Rosa S, Marengo M, Romagnoli S et al. Management of acute kidney injury and extracorporeal blood purification therapies during the COVID-19 pandemic: the Italian SIN-SIAARTI Joint Survey (and Recommendations for Clinical Practice). *Front Med* 2022; 9 : 1–10. <https://doi.org/10.3389/fmed.2022.850535>.
29. Weiss R, Meersch M, Pavenstädt HJ et al. Acute kidney injury a frequently underestimated problem in perioperative medicine. *Dtsch Arztebl Int* 2019; 116 : 833–42.
30. Basile DP, Bonventre JV, Mehta R et al. Progression after AKI: understanding maladaptive repair processes to predict and identify therapeutic treatments. *J Am Soc Nephrol* 2016; 27 : 687–97. <https://doi.org/10.1681/ASN.2015030309>.
31. Franzin R, Stasi A, Ranieri E et al. Targeting premature renal aging: from molecular mechanisms of cellular senescence to senolytic trials. *Front Pharmacol* 2021; 12 : 1–22. <https://doi.org/10.3389/fphar.2021.630419>.
32. Ferenbach DA, Bonventre J. Mechanisms of maladaptive repair after AKI leading to accelerated kidney ageing and CKD. *Nat Rev Nephrol* 2015; 11 : 264–76. <https://doi.org/10.1038/nrneph.2015.3>.
33. Kellum JA, Chawla LS. Cell-cycle arrest and acute kidney injury: the light and the dark sides. *Nephrol Dial Transplant* 2016; 31 : 16–22. <https://doi.org/10.1093/ndt/gfv130>.
34. Castellano G, Franzin R, Sallustio F et al. Complement component C5a induces aberrant epigenetic modifications in renal tubular epithelial cells accelerating senescence by Wnt4/β catenin signaling after ischemia/reperfusion injury. *Aging (Albany NY)* 2019; 11 : 4382–406. <https://doi.org/10.18632/aging.102059>.
35. Lazzeri E, Angelotti ML, Peired A et al. Endocycle-related tubular cell hypertrophy and progenitor proliferation re-cover renal function after acute kidney injury. *Nat Commun* 2018; 9 : 1–18. <https://doi.org/10.1038/s41467-018-03753-4>.
36. De Chiara L, Conte C, Semeraro R et al. Tubular cell polyploidy protects from lethal acute kidney injury but promotes consequent chronic kidney disease. *Nat Commun* 2022; 13 : 1–19. <https://doi.org/10.1038/s41467-022-33110-5>.
37. Frawirth KA, Riley JL, Harris MH et al. The CD28 signaling pathway regulates glucose metabolism. *Immunity* 2002; 16 : 769–77. [https://doi.org/10.1016/S1074-7613\(02\)00323-0](https://doi.org/10.1016/S1074-7613(02)00323-0).
38. Smith JA, Jay Stallons L, Schnellmann RG. Renal cortical hexokinase and pentose phosphate pathway activation through the EGFR/akt signaling pathway in endotoxin-induced acute kidney injury. *Am J Physiol Ren Physiol* 2014; 307 : F435–44. <https://doi.org/10.1152/ajprenal.00271.2014>.
39. Waltz P, Carchman E, Gomez H et al. Sepsis results in an altered renal metabolic and osmolyte profile. *J Surg Res* 2016; 202 : 8–12. <https://doi.org/10.1016/j.jss.2015.12.011>.
40. Kalakeche R, Hato T, Rhodes G et al. Endotoxin uptake by S1 proximal tubular segment causes oxidative stress in the downstream S2 segment. *J Am Soc Nephrol* 2011; 22 : 1505–16. <https://doi.org/10.1681/ASN.2011020203>.
41. Tran M, Tam D, Bardia A et al. PGC-1α promotes recovery after acute kidney injury during systemic inflammation in mice. *J Clin Invest* 2011; 121 : 4003–14. <https://doi.org/10.1172/JCI58662>.
42. Howell GM, Gomez H, Collage RD et al. Augmenting autophagy to treat acute kidney injury during endotoxemia in mice. *PLoS One* 2013; 8 : 4–11. <https://doi.org/10.1371/journal.pone.0069520>.
43. Figueiredo N, Chora A, Raquel H et al. Anthracyclines induce DNA damage response-mediated protection against severe sepsis. *Immunity* 2013; 14 : 874–84. <https://doi.org/10.1016/j.immuni.2013.08.039>.
44. Lu W, Karuppagounder SS, Springer DA et al. PGC1α-dependent NAD biosynthesis links oxidative metabolism to renal protection. *Nature* 2015; 531 : 528–32.
45. Lee SA, Noel S, Sadasivam M et al. Role of immune cells in acute kidney injury and repair. *Nephron* 2017; 137 : 282–6. <https://doi.org/10.1159/000477181>.
46. Zager RA. “Biologic memory” in response to acute kidney injury: cytoresistance, toll-like receptor hyper-responsiveness and the onset of progressive renal disease. *Nephrol Dial Transplant* 2013; 28 : 1985–93. <https://doi.org/10.1093/ndt/gfr101>.
47. Kellum JA, Sileanu FE, Bihorac A et al. Recovery after acute kidney injury. *Am J Respir Crit Care Med* 2017; 195 : 784–91. <https://doi.org/10.1164/rccm.201604-0799OC>.
48. Martensson J, Martling CR, Oldner A et al. Impact of sepsis on levels of plasma cystatin C in AKI and non-AKI patients. *Nephrol Dial Transplant* 2012; 27 : 576–81. <https://doi.org/10.1093/ndt/gfr358>.
49. Prowle JR, Kolic I, Purdell-Lewis J et al. Serum creatinine changes associated with critical illness and detection of persistent renal dysfunction after AKI. *Clin J Am Soc Nephrol* 2014; 9 : 1015–23. <https://doi.org/10.2215/CJN.11141113>.
50. Husain-Syed F, Ferrari F, Sharma A et al. Preoperative renal functional reserve predicts risk of acute kidney injury after cardiac operation. *Ann Thorac Surg* 2018; 105 : 1094–101. <https://doi.org/10.1016/j.athoracsurg.2017.12.034>.
51. Klaus F, Keitel Da Silva C, Meinerz G et al. Acute kidney injury after liver transplantation: incidence and mortality. *Transplant Proc* 2014; 46 : 1819–21. <https://doi.org/10.1016/j.transproceed.2014.05.053>.
52. Jing L, Chen W, Guo L et al. Acute kidney injury after lung transplantation: a narrative review. *Ann Transl Med* 2021; 9 : 717–. <https://doi.org/10.21037/atm-20-7644>.
53. Lertjitbanjong P, Thongprayoon C, Cheungpasitporn W. Acute kidney injury after lung transplantation: a systematic review and meta-analysis. *J Clin Med* 2019; 8 : 1–19. <https://doi.org/10.3390/jcm8101713>.
54. Wang CJ, Wetmore JB, Israni AK. Old versus new: progress in reaching the goals of the new kidney allocation system. *Hum Immunol* 2017; 78 : 9–15. <https://doi.org/10.1016/j.humimm.2016.08.007>. Downloaded from <https://academic.oup.com/ckj/article/16/11/1813/1760476> by guest on 03 November 2023
55. Palmisano A, Gandolfini I, Delsante M et al. Acute kidney injury (AKI) before and after kidney transplantation: causes, medical approach, and implications for the long-term outcomes. *J Clin Med* 2021; 10 : 1484. <https://doi.org/10.3390/jcm10071484>.
56. Dong V, Nadim MK, Karvellas CJ. Post-liver transplant acute kidney injury. *Liver Transpl* 2021; 27 : 1653–64. <https://doi.org/10.1002/lt.26094>.
57. Jing L, Chen W, Zhao L et al. Acute kidney injury following adult lung transplantation. *Chin Med J (Engl)* 2022; 135 : 172–80. <https://doi.org/10.1097/CM9.0000000000001636>.
58. Dorcic J, Greite R, Vijayan V et al. Kidney injury after lung transplantation: long-term mortality predicted by post-operative day-7 serum creatinine and few clinical factors. *PLoS One* 2022; 17 : 1–17. <https://doi.org/10.1371/journal.pone.0265002>.
59. Hussaini T, Yoshida EM, Partovi N et al. Early persistent progressive acute kidney injury and graft failure post liver transplantation. *Transplant Direct* 2019; 5 : 1–5. <https://doi.org/10.1097/TXD.0000000000000868>.
60. Lim WH, Johnson DW, Teixeira-Pinto A et al. Association between duration of delayed graft function, acute rejection, and allograft outcome after deceased donor kidney transplantation. *Transplantation* 2019; 103 : 412–9. <https://doi.org/10.1097/TP.0000000000002275>.
61. De Sandes-Freitas TV, Felipe CR, Aguiar WF et al. Prolonged delayed graft function is associated with inferior patient and kidney allograft survivals. *PLoS One* 2015; 10 : 1–13.
62. Giral-Classe M, Hourmant M, Cantarovich D et al. Delayed graft function of more than six days strongly decreases long-term survival of transplanted kidneys. *Kidney Int* 1998; 54 : 972–8. <https://doi.org/10.1046/j.1523-1755.1998.00071.x>.
63. Pickkers P, Darmon M, Hoste E et al. Acute kidney injury in the critically ill: an updated review on pathophysiology and management. *Intensive Care Med* 2021; 47 : 835–50. <https://doi.org/10.1007/s00134-021-06454-7>.
64. The STARTRT-AKI Investigators, for the Canadian Critical Care Trials Group, the Australian and New Zealand Intensive Care Society Clinical Trials Group, the United Kingdom Critical Care Research Group, the Canadian Nephrology Trials Network and the ICCTG. Timing of initiation of renal-replacement therapy in acute kidney injury. *N Engl J Med* 2020; 383 : 240–51. <https://doi.org/10.1056/NEJMoa2000741>.
65. Macedo E, Bihorac A, Siew ED et al. Quality of care after AKI development in the hospital: consensus from the 22nd Acute Disease Quality Initiative (ADQI) conference. *Eur J Intern Med* 2020; 80 : 45–53. <https://doi.org/10.1016/j.ejim.2020.04.056>.
66. Waikar SS, Curhan GC, Wald R et al. Declining mortality in patients with acute renal failure, 1988 to 2002. *J Am Soc Nephrol* 2006; 17 : 1143–50. <https://doi.org/10.1681/ASN.2005091017>.
67. Hsu CY, Hsu RK, Yang J et al. Elevated BP after AKI. *J Am Soc Nephrol* 2016; 27 : 914–23. <https://doi.org/10.1681/ASN.2014111114>.
68. Harel Z, Wald R, Bargman JM et al. Nephrologist follow-up improves all-cause mortality of severe acute kidney injury survivors. *Kidney Int* 2013; 83 : 901–8. <https://doi.org/10.1038/ki.2012.451>.
69. Silver SA, Siew ED. Follow-up care in acute kidney injury: lost in transition. *Adv Chronic Kidney Dis* 2017; 24 : 246–52. <https://doi.org/10.1053/j.ackd.2017.05.008>.
70. Silver SA, Adhikari NK, Bell CM et al. Nephrologist follow-up versus usual care after an acute kidney injury hospitalization (FUSION): a randomized controlled trial. *Clin J Am Soc Nephrol* 2021; 16 : 1005–14. <https://doi.org/10.2215/CJN.17331120>.
71. Kiers HD, Van Den Boogaard M, Schoenmakers MCJ et al. Comparison and clinical suitability of eight prediction models for cardiac surgery-related acute kidney injury. *Nephrol Dial Transplant* 2013; 28 : 345–51. <https://doi.org/10.1093/ndt/gfs518>.
72. Parreco J, Soe-Lin H, Parks JJ et al. Comparing machine learning algorithms for predicting acute kidney injury. *Am Surg* 2019; 85 : 725–9. <https://doi.org/10.1177/000313481908500731>.
73. Wilson T, Quan S, Cheema K et al. Risk prediction models for acute kidney injury following major noncardiac surgery: systematic review. *Nephrol Dial Transplant* 2016; 31 : 231–40.
74. Liang Q, Xu Y, Zhou Y et al. Severe acute kidney injury predicting model based on transcontinental databases: a single-centre prospective study. *BMJ Open* 2022; 12 : e054092. <https://doi.org/10.1136/bmjopen-2021-054092>.
75. Husain-Syed F, Ferrari F, Sharma A et al. Persistent decrease of renal functional reserve in patients after cardiac surgery-associated acute kidney injury despite clinical recovery. *Nephrol Dial Transplant* 2019; 34 : 308–17. <https://doi.org/10.1093/ndt/gfy227>.
76. Villa G, Samoni S, De Rosa S et al. The pathophysiological hypothesis of kidney damage during intra-abdominal hypertension. *Front Physiol* 2016; 7 : 1–4. <https://doi.org/10.3389/fphys.2016.00055>.
77. Samoni S, Nalesso F, Meola M et al. Intra-parenchymal renal resistive index variation (IRIV) describes renal functional reserve (RFR): pilot study in healthy volunteers. *Front Physiol* 2016; 7 : 1–6. <https://doi.org/10.3389/fphys.2016.00286>.